

THE INTERNATIONAL LILY REGISTER & CHECKLIST

TWENTY-FIRST SUPPLEMENT

The following list contains names registered between 1 October 2001 and 30 September 2002.

Raisers and introducers are urged to ensure that all their plants have been registered and are reminded that registration should take place *before* a plant is released. Registration forms can be obtained from the International Lily Registrar, Royal Horticultural Society's Garden, Wisley, Woking, Surrey, England GU23 6QB. The form may also be downloaded from the RHS website (<http://www.rhs.org.uk>)

Forms may also be obtained from a number of regional representatives. The International Cultivar Registration Authority is indebted to these individuals for the considerable effort they make to obtain and pass on all information about lily cultivars. They are as follows:

Regional representative, Australia

Neil Jordan, 27 Scarfe Street, Camdale, Tasmania 7320, Australia.

Regional representative, Czech Republic

Ing. Břetislav Mičulka, 687 07 Velehrad 225, Czech Republic.

Regional representative, The Netherlands

Dr Johan van Scheepen, Royal General Bulbgrowers' Association,
Postbus 175, NL-2180 AD Hillegom, The Netherlands

Regional representative, New Zealand

Mrs Merle L Lepper, PO Box 1394, Christchurch, New Zealand

Regional representative, North America

Michael Homick, RR no.2, Tillsonburg, Ontario, Canada N4G 4G7

In addition the International Cultivar Registration Authority (ICRA) is grateful to a great many other correspondents who have continued to bring new lilies to its attention.

This Supplement continues the use of a dual subdivision for every entry. Thus both flower habit (up-, out- or down-facing) and flower form (trumpet, bowl, flat or recurved) are indicated for each entry when this information is available. The existing lettering is retained, with flower habit given first, separated from the flower form by a slash mark (/). An entry such as I(b/c) 'Aloft' therefore indicates this is an out-facing, flat-flowered asiatic lily. It is not a requirement that dual subdivisions should be used at all times. The way in which the information is presented means that where, for a particular purpose it is necessary, one (or both) can be ignored. Their inclusion, however, together with the division number helps to present in an abbreviated form a quick 'picture' of the flower concerned. For a full account of any plant it is necessary to refer to the description.

This Supplement also includes accounts, for information only, of a number of cultivars whose epithets are not registered. In most cases these concern lily names which are, for a variety of reasons, unacceptable but where it is now unlikely that the epithet will be changed. These form part of the Checklist function of this work.

Dr Alan C Leslie, International Lily Registrar, RHS Garden, Wisley, Woking, Surrey GU23 6QB, England (Fax: 01483 211750, email: alanl@rhs.org.uk).

I (a/c) 'Ace of Spades'

cv. (('Sun Ray' × 'Windahloo') × ('Edith' × 'Fudge Sundae')) H (1989), G (1993), N (1996) & REG (2001); E.E. Fox I: Fox Lily Ranch, 2002. Inside vivid yellow (12A), with spear-point-shaped, dark red (187A) brushmarks, throat brilliant yellow-green (150B); outside vivid yellow (9A); unspotted; nectaries lime, frosted white; pollen vivid orange-yellow (23A); stigma off white towards light yellow-green (145D). Fls 160mm wide; petals 80 × 40mm, margins slightly ruffled, tips slightly recurved. Lvs alternate, 80 × 12mm. Stems 0.6m, pale green, with 9-12 fls. Late July. (Etymology: the name suggested by the spade-like shape of the brushmark.)

I (a-b/-) 'Albatros'

cv. (('Juliana' × unknown) × (un-named seedling no. 8759 × 'Orlito')) H (1992) & G (1994); World Flower B.V. N, I & REG: 2000. Inside white (155D) with a green throat; outside white (155D); no spots but papillae present; stigma green, spotted purple; pollen brown. Fls 130mm wide; petals 70 × 45mm, margins ruffled, tips recurved. Lvs alternate, 100 × 20mm. Stems 1.35m, green, with dark spots and stripes and c.12 fls. Early. (The spelling with one 's' is intentional.)

I (a/b) 'Alfred Weinreich'

cv. ('Heidi III' × 'Sonnenwunder') H (1992) & G (1995); G. Steinbrück N, I & REG: 2002. Inside pale yellow (18D) overlaid strong orange-yellow (22A) – especially in lower ½, throat pale yellow-green (157C) with delicate papillae; outside orange-yellow; numerous, moderate-sized spots and very reduced brushmark streaks of dark purple (79A); nectaries pale yellow-green (157A), green-white hoary; pollen dark greyish reddish brown (200A); stigma dark red (187A), style greenish yellow. Fls 150mm wide; petals 90 × 42mm, not ruffled, tips slightly recurved. Lvs alternate, 60 × 20mm. Stems 1.4m, dark green, with c.11 fls. Mid to late July. (Standard: colour print provided by registrant (WSY 0032242). Etymology: named for Alfred Weinreich a lily breeder from Wolmirstedt, Germany.)

I (c/d) 'Alkor'

cv. ('Rondo' × Sutter's Gold Group) R & N: K. Braun I: 2001 REG: 2002. Inside yellow, throat green; outside yellow, with yellowish brown ribs; moderate-sized, numerous, dark brown spots over ¾ of each petal; papillae pale yellow; nectaries green; pollen orange; stigma green. Fls 80mm wide; petals 60 × 26mm; pedicels 65mm long, brown dotted. Lvs alternate, 60 × 10mm. Stems 0.8m, brown dotted, with a moderately long inflorescence. Early July.

VII (-/d) Allegro Group

grp (*auratum* var. *virginale* × 'Allegra') L: Blackthorne Gardens, 1982. Fls white, with a few pink spots; petal margins ruffled. Stems 1.2-1.5m, with up to 20 fls. August.

I (a/-) 'Alwin'

cv. ('Yellow Star' × ('Tabasco' × *L. × hollandicum*)) H (1977) & G: G. & S.E. Bottema N: G. Bottema. Inside vivid to strong reddish orange (34A/C), with orpiment orange blotches, dark brown spots and violet red edges in the throat; outside moderate red (179A/B), the outer petals moderate orange (168C) edged burnt brownish red and tipped saffron yellow; nectaries greyish red (184A); pollen dark reddish orange (172B). Fls 170mm wide; petals 85 × 40mm, not ruffled. Lvs alternate, 190 × 12.5mm, mid green. Stems 1.25m, green, tinged red, with c.4 fls. June to July.

VII (a-bl-) 'Amazing'

cv. ((*speciosum* 'Brabander' × ('Star Gazer' × 'Early Splendor')) × 'Elysee') H (pre-1996) & G: Bischoff Tulleken Lelies B.V. REG: H.J.A. Rooyakkers B.V., 2002. Inside white (155D), with brilliant yellow (9C) along midvein and base of midvein yellow; outside white (155D); nectaries yellow-green; yellow spots present; papillae also present; pollen orange-red; stigma purple. Petal margins ruffled, tips recurved. Early to midseason.

VIII (b/b) 'Amber Ace'

cv. (Trumpet seedling (with Golden Splendor Group and a multiple cross in its parentage) × 'Canterbury Belle') H (1995) & G (1998): A.E. Alexander N, I & REG: 2002. Inside apricot, with an apricot throat; outside apricot with some wine red stripes; unmarked; nectaries green; pollen orange; stigma frosty green. Fls 130mm wide; petals 90 × 50mm, not ruffled, tips slightly reflexed. Lvs alternate, 90 × 15mm. Stems 1.3m, green, with purple flecks and c.8 fls. January (New Zealand) (Standard: colour print provided by registrant (WSY 0032050). An orientpet lily.)

I (a/-) 'Amber Dream'

cv. (Un-named seedling (no. 89-68-03) × un-named seedling (no. 89-6-21)) H (1992), G (1995), N (2000), I & REG (2002): World Flower B.V. Inside strong orange (24A/25A) with a reddish glow at the apex; outside vivid orange-yellow/strong orange (23A/24A) with a yellowish tinge at apex; not spotted, but papillae present; stigma and pollen orange. Stems 1.1m, green, with 8-10 fls (16-18cm bulb). Very early to early.

VIII (c/c) 'Amber Star'

cv. ((*henryi* × un-named trumpet) × 'Canterbury Belle') H (1992) & N (1995): A.E. Alexander N, I & REG: 2002. Inside apricot; outside apricot marked with pink and green; nectaries green; pollen brown; stigma grey. Fls 140mm wide; petals 110 × 50mm, margins not ruffled, tips slightly recurved. Lvs alternate, 150 × 20mm. Stems 1.8m, black, flecked with green, with 10-12 fls in a racemose inflorescence. January (New Zealand). Ryrirube Group. (Standard: colour print provided by registrant (WSY 0032055). An orientpet lily.)

I (a/b) 'Ambos Mundos'

cv. ('Alexis' × mixed pollen of 'Limpopo', 'Brushstroke' and 'Impact') H (1991) & G (1994): G. Steinbrück N, I & REG: 2002. Inside moderate reddish orange (35A) throughout; outside yellowish white with pale green midribs; unspotted; nectaries pale yellow-green (196B); pollen deep red (185A) on thin, short, light orange-yellow (19A) filaments; stigma moderate yellowish white. Fls 150mm wide; petals 80 × 30mm, not ruffled, not recurved. Lvs alternate, 70 × 17mm. Stems 1.0m, mid green, with 8-10 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032244). Etymology: Ambos Mundos is a place in Havana, Cuba where Ernest Hemingway worked.

I (-/-) Antique Ivory Group

grp (Selections from Futura Group) R, N & I (1982): Blackthorne Gardens. Fls yellowish white.

VII (-/-) Appleblossom Time Group

grp (Parentage unknown) R (pre-1982) & N: Blackthorne Gardens. Fls pastel pink. Stems up to 1.0m. Early.

I (a/-) 'Apricot'

cv. ('Connecticut King' × 'Enchantment') R (pre-1981) & N: M. Rohrmeier. Fls light orange, unspotted. (Unacceptable name (Art. 17.2).)

I (-/-) Apricot Glory Group

grp (Selections from Futura Group) R, N & I (1982): Blackthorne Gardens. Fls apricot.

I (a/b) 'Aquino'

cv. (Parentage unknown) R: W. Paauw & Zn B.V., pre-1987 REG: Peacock B.V., 2002. Inside vivid yellow (12A), throat yellow; spots black with red margins; pollen dark brown; stigma brown. Petals slightly ruffled, tips recurved. Stems brown/brown purple.

I (a/-) 'Arancia'

cv. (Parentage unknown) R (pre-1989), N & REG (2002): P.M. Hoogland. Inside a little darker than strong orange (24A); outside strong orange (24A), with strong orange-yellow (24B) midveins; markings dark red-purple; spots present; papillae also present; pollen dark brown; stigma purple. Petal margins ruffled, tips recurved. Stems with dark markings. Early to midseason.

I (b-c/-) 'Arctica'

cv. (('Stroinaja' × *davidii* var. *willmottiae*) × *cernuum* hybrids) R (c1983) & N: M.F. Kireeva. Fls white, with brown spots; large. Stems 1.3m, with bulbils. July. (Sometimes seen as 'Arktika'; unacceptable name (Art. 17.9).)

I (a/c) 'Arctic Sunrise'

cv. (Un-named seedling × un-named seedling) H (1990), G (1993), N (2002) & REG (2002): R.H. Bazett I: The Lily Nook, 2002. Inside white, with a pale orange flush at base extending up c½ of each petal, throat pale orange shading to green at nectaries; outside white, with grey-green midribs; numerous, small, maroon spots on lower ⅓ of each petal; nectaries green; pollen burnt orange; stigma green. Fls 160mm wide; petals 87 × 42mm, not ruffled, tips slightly recurved. Lvs alternate, 113 × 18mm. Stems 0.9m, dark green, with c17 fls, in a racemose inflorescence (including some secondary buds). Late June to early July. (Standard: colour reproduction provided by registrant (WSY 00351212). Etymology: descriptive of flower colouration.)

IX (b/b) *auratum* 'Cruentum'

cv. (Selection from *auratum*) I: E. Pynaert-van Geert, c1891. Fls 25-30cm wide, creamy white, lightly tinted nankeen yellow near edges, with a very broad central band of chocolate purple. Lvs dark green. Stems 60-80cm, with 6-8 fls. (Published, with colour illustration in *Rev. Hort. Belg.* 17: 193 (1891).)

IX (-/-) *auratum* Kimono Group

grp (Seed-raised selections from *auratum* var. *platyphyllum*) R (pre-1992) & N: Fairdale Nursery. Fls white, with gold bands and gold throats; spotted. (Unacceptable name (Art. 26.1).)

I (c/-) 'Baby Pink Bells'

cv. (Parentage unknown: may be a *cernuum* hybrid) H (c1980), G & N: F. Fellner REG: 2002. Inside pink; outside pinkish; unmarked; nectaries pink; pollen brown; stigma cream, on a very short style. Fls unusual in being more-or-less campanulate

(i.e. petals not spreading out), 50-70mm wide; petals 55 × 20mm, not ruffled, tips not recurved. Lvs alternate, 60-70 × 10mm. Stems 0.75m, red-black, with 10-30 fls in a tall inflorescence. Mid July. (Standard: colour print provided by registrant (WSY 0035251).)

I? (b/-) 'Baby Ribbon Pink'

cv. (Parentage unknown) **I**: Blackthorne Gardens, 1985. Fls semi-double, baby ribbon pink, with a few darker spots in the centre; pollen absent. Stems 0.6m. July.

I (a/b) 'Baltic Star'

cv. (Parentage unknown) **H** (1991), **G** (1994), **N** (1997) & **I** (1997): J. Vasarietis **REG**: 2001. Inside white, with a very pale pink band around centre of flower at level of nectaries; outside white, with very slight pink shading; c20 tiny, dark spots scattered level with nectaries; nectaries white; pollen brown; stigma greyish pale yellow. Fls 180mm wide; petals 85 × 37mm, margins not or very slightly ruffled, tips slightly recurved. Lvs alternate, 75 × 14mm, dark green. Stems 1.0m, green, with c6 fls. Early July. (Standard: colour print provided by registrant (WSY 0030818). Published in *Lilium Balticum* Lily Register, 2000.)

I (a/-) 'Bananarama'

cv. (Un-named seedling × un-named seedling) **H**, **G**, **N** & **I**: Hoffgaard B.V. **REG**: 2001. Inside vivid yellow (12A), throat green; outside vivid yellow (9A); nectaries green; a few small spots and some papillae present; pollen orange-brown; stigma dark purple. Fls 130mm wide; petals not ruffled, tips slightly recurved. Stems 0.9m, with c5 fls. Very early.

VIII (a/b) 'Barbina'

cv. ('Snow Cap' (OBF) × 'Longicanda') **R** & **N**: K. Braun **I**: 2001 **REG**: 2002. Inside yellowish cream, centre and throat yellow; outside creamy, with cream and green ribs; unspotted; nectaries light green; pollen reddish brown; stigma cream. Fls 140mm wide; petals 80 × 37mm, moderately recurved; pedicels 70mm long, green. Lvs alternate, 80 × 10mm. Stems 0.75m, green. Early July. (A *longiflorum*/asiatic hybrid.)

I (a/-) 'Bariton'

cv. (Parentage unknown) **H** (1994), **G** (1996), **N** (2001) & **REG** (2002): Sande B.V. Inside vivid orange (28A/B); outside vivid orange (28B); nectaries orange; spots and papillae absent; pollen red-brown; stigma purple-red. Petal margins slightly ruffled, tips recurved. Stems 1.2m. Early.

I (a/b) 'Basra'

cv. (Parentage unknown) **R** (pre-1987), **N** & **REG** (2002): Fa. Jaap Boon. Inside light orange (26C), strong orange (28B) at base, light orange in throat, markings dark purple; spots present; pollen red-brown; stigma light orange. Petal margins not ruffled, tips recurved. Stems mid green, with little darker marking.

VIII (a-b/b) 'Becia'

cv. (Unknown × Longistar Group) **H** (1995), **G** (1997), **N** (2001) & **I** (2001): J. Kapias **REG**: 2002. Inside vivid yellow (15B), throat strong yellow-green (144C); outside light greenish yellow (4C); brownish orange (165B) spots on basal ½ of each petal; nectaries pale greenish yellow (2D); pollen greyed orange (165); stigma vivid yellow (15B). Fls 220mm wide; petals 120-135 × 38-50mm, margins slightly waved, tips slightly recurved. Lvs alternate, 150 × 30mm. Stems 1.0m (under glass), mid

green, with c15 fls (including secondaries). Mid July. Reported to be polyploid. (A *longiflorum*/asiatic hybrid. Standard: colour print provided by registrant (WSY 0031318).)

I (a/b) 'Bel Ami'

cv. (Parentage unknown) R & N: The Netherlands, pre-1996. Fls pure white, unspotted; stigma green; pollen dark. Petal not ruffled, not recurved. Stems 60cm. Midseason. Reported to be tetraploid.

I (a/b) 'Belanna'

cv. (Parentage unknown) H (1990), G (1993), N (1996) & I (1996): J. Vasarietis REG: 2001. Inside strong purplish red (64B) throughout (fading to deep purplish pink (64B)); outside strong purplish red (64B) with a band (c10mm wide) of deep purplish pink (64D) around the midvein; no noticeable spots; nectaries white on a greenish ground, conspicuous; pollen brownish orange; stigma light green. Fls 190mm wide; petals 95 × 35mm, relatively rather narrow, not or very narrowly waved at margins, tips slightly recurved. Lvs alternate, 100 × 8mm. Stems 0.9m, dark green, with c11 fls. Mid to late July. (Standard: colour print provided by registrant (WSY 0030817). Etymology: combination of belle (= beautiful) and Anna.)

I (a/-) 'Belinda'

cv. (('Harmony' × *dauidii*) × orange seedlings) R (c1977) & N: De Jong Lelies Holland B.V. Inside vivid reddish orange (30B), tips vivid yellowish orange (30C), throat vivid reddish orange (30A-B); outside strong yellowish pink (32C); a few brown spots; pollen dark brown. Petals 55 × 28mm, not ruffled, slightly recurved. Lvs alternate, 100 × 15mm. Stems 0.8m, dark, with c10 fls. July.

II (-/-) 'Bendick'

cv. (*martagon* origin) L: Honeywood Lilies, 1984. Fls mauve pink, with sparse spots. Stems 1.2m.

I (a/b) 'Big Mac'

cv. (Parentage unknown) R: unknown, pre-1992. Inside orange-red at tips, apricot in centre, with numerous, large, brown-purple spots over ½ of each petal; nectaries white granulated; pollen orange. Fls 115mm wide; petals 70 × 30mm, very slightly recurved. Lvs alternate, 70 × 8mm. Stems 0.7m, dotted with violet black at the base. Early June. (Has been grown in the Czech Republic.)

I (b/b) 'Bílá Opava'

cv. ('Feuer und Rauch' × 'Sterling Star') R & N: J. Dudek I & REG: 2001. Inside creamy white, tips paler, throat yellowish; outside creamy, with purple ribs at the base; numerous, moderately-sized, purple spots over ½ of each petal; pollen and stigma brown. Fls 190mm wide; petals 90 × 46mm, tips slightly recurved; pedicels 100mm long, green. Lvs alternate, 130 × 12mm, glossy green. Stems 1.4m, thick, brown. Early July.

I (a/b) 'Bingo'

cv. (Selection from Pastel Group) R: Bischoff Tulleken Ltd N & I: 1979. Inside creamy white, throat creamy yellow; outside creamy yellow; numerous small, brown spots on basal ½ of each petal; nectaries white; pollen orange-brown. Fls 100mm wide; petals 80 × 20mm, not ruffled, tips slightly recurved. Lvs alternate, 90 × 8mm, dark green. Stems 0.75m, dark, with c6 fls. June. (Unacceptable name (Art. 17.2))

I (-/-) 'Bingo'

cv. (*leichtlinii* var. *maximowiczii* hybrid) R (pre-1962), N & I: Oregon Bulb Farms. Buds green soon turning lemon yellow, tinted bronze at base, opening yellow inside and flushed bronze at base outside; spotted maroon; pollen light orange. Fls 125mm wide. Stems 1.0m, brownish purple, with numerous stem bulbils. August.

I (b/d) 'Blackberry Intrigue'

cv. (('Condamine' × 'Summer Night') × un-named red seedling) H (1989) & G (1991): A.J. Burnett N (1997) & REG (2001): E.E. Fox I: Fox Lily Ranch, 2002. Inside dark red (187A), with a dark red (187A) throat; outside dark red (187A); a few dark greyish purple (202A) raised spots; nectaries and stigma dark red (187A); pollen strong brown (172A). Fls 130mm wide; petals 65 × 35mm, not ruffled, tips strongly recurved. Lvs alternate, 110 × 12mm. Stems 0.9m tall, heavily mottled (appears black), with 12-18 fls. Late July. (Etymology: name descriptive of colour.)

I (a/b-c) 'Blast Off'

cv. (Un-named seedling × *dauricum*) H (1992), G (1994), N (1995) & I (1999): M.E. Rollinson REG: 2002. Inside vivid orange (28A) throughout, with large, darker spots over ½–¾ of each petal; nectaries strong orange (25A); pollen moderate reddish orange (171A); stigma deep red (185A). Fls 135mm wide; petals 68 × 30-40mm, not ruffled, tips very slightly recurved. Pedicels 110mm long. Lvs in whorls of 3, 37 × 13mm, rather short and recurved, green, with brown shading. Stems 0.94m, moderate olive green (137A), mottled brown, with 8-15 fls. December to January (New Zealand). (Standard: colour print provided by registrant (WSY 0032036).)

VI (b/a) 'Bobo'

cv. (*regale* × unknown) × unknown) R & N: J. Dudek I & REG: 2001. Inside white, with pinkish edges, throat yellow-green; outside white, yellow-green at base, the ribs greenish with pinkish bases; unspotted; nectaries green; pollen yellow-orange; stigma green. Fls 120mm wide; petals 120 × 52mm, margins waved, tips recurved; pedicels 70mm long, spotted with brown. Lvs alternate, glossy green. Stems 1.3m, rigid, with fls in an umbellate inflorescence. Mid July.

I (a/b) 'Brandenburger Rotgold'

cv. ('Impala' × un-named seedling (no. AS12, from Weidemann)) H (1988), G (1991), N (1999) & I (2000): G. Steinbrück REG: 2002. Inside strong orange (168B) throughout, with scattered, tiny, dark reddish orange (175B) spots on basal ¼ of each petal; nectaries yellowish grey (156C), tinged soft pink at first; pollen vivid reddish orange (34A); stigma strong reddish orange (169A). Fls 120mm wide; petals 82 × 47mm, not ruffled, not recurved. Lvs alternate, 50 × 24mm. Stems 0.9m, dark green, with c7 fls. Mid June. (Standard: colour print provided by registrant (WSY 0032246).)

VIII (a-b/-) 'Bravoure'

cv. ((Un-named seedling × 'Sahara') × (*longiflorum* × 'Aristo')) H (1989), G (1993) & N (1999): Bischoff Tulleken Lelies B.V. I (1999) & REG (2001): Kwekersvereniging "Het Vrije Westen". Inside brilliant yellow (20A), fading to about pale yellow (18D) with age, base of midveins orange-green to orange; outside light yellow (18A); spots and papillae present; nectaries orange-green; pollen reddish brown; stigma orange-yellow. Fls 220mm wide; petals 110 × 50mm, margins ruffled, tips recurved. Lvs alternate, 180 × 20mm. Stems 80cm, with dark spots and stripes and c4-5 fls. Early. (A *longiflorum*/asiatic hybrid.)

I (c/-) 'Bronzine'

cv. (Parentage unknown) N & I (1980): B.G. Hayler. Inside cream, tips coppery bronze; spotted. Petal tips recurved. Stems 1.0m. December (Australia).

VI (b-c/b) 'Buchlov'

cv. (('Black Dragon' × 'Damson') × (unknown × 'Damson')) R & N: B. Mičulka I & REG: 2002. Inside white, without spots, centre and throat apricot; outside with apricot shining through, midveins green; pollen apricot; nectaries dark green; stigma greyed brown. Fls 130mm wide; petals 115 × 50mm, moderately recurved, slightly ruffled; pedicels 120mm long, thick, slightly spotted. Lvs alternate, 130 × 14mm, dark green. Stems 1.15m, spotted brown-red and with a short inflorescence. Mid July.

VI (-/a) Burnished Gold Group

grp (*regale* hybrids × Royal Group) R (pre-1966) & N: Blackthorne Gardens. Fls old gold, 150 × 150mm. Stems to 1.5m, with up to 17 fls. July.

I (c/d) Butterball Group

grp (Hybrids of *dauidii* var. *willmottiae*) R (pre-1978) & N: Blackthorne Gardens. Shades of chartreuse and gold. Petals not ruffled, strongly recurved. Lvs alternate. Stems 1.0-1.5m. June to July. (Previously listed in error as Buttercup Group (1982 Register, p. 45); unacceptable name (Art. 26.1).)

I (a/b) 'Buttercup'

cv. (Parentage unknown) R (pre-1985), N & I: Oregon Bulb Farms. Inside creamy yellow, with moderate-sized, brownish spots over basal ⅓ of each petal. Fls 125mm wide; petals broad, not ruffled, not recurved. Stems 0.6-0.9m. Late June. (Unacceptable name: Art. 17.2).

VII (-/d) Candy Frost Group

grp (Hybrids involving *rubellum* and 'Manawatu') H: L. Woodriff, pre-1966 S: L. Woodriff & J.M. Shaver. Inside white, spotted with pink and crimson. Fls 125-150mm wide; petals broad, recurved. Stems 0.9-1.9m. August to September.

I (a/-) Cape Cod Moonglow Group

grp (Parentage unknown) H: R.W. Bemis, 1960s N & I: Blackthorne Gardens, c1982. Fls yellow, tinged green. Stems 1.2-1.5m. July to September.

I? (a/-) 'Cape Cod Sunray'

cv. (Parentage unknown) I: Blackthorne Gardens, 1985. Fls golden yellow, with golden bands. Stems 0.6m. June.

I (b/-) 'Cape Cod Sunshine'

cv. (Parentage unknown) H: R.W. Bemis, 1960s N & I: Blackthorne Gardens, c1982. Fls yellow, faintly spotted; no pollen. Stems 0.6m. July onwards.

I? (-/-) 'Cape Cod Sunstar'

cv. (Parentage unknown) R (pre-1983) & N: Blackthorne Gardens. Fls yellow, slightly overlaid green, with gold bands along midribs. Stems 0.45-0.6m. July.

I (a/-) 'Cape Cod Yankee'

cv. ('Connecticut Yankee' × unknown) H: R.W. Bemis, 1960s N & I: Blackthorne Gardens, c1982. Fls apricot coral, unspotted. Stems 1.2m. July to August.

I (a/-) 'Captain Marvel'

cv. (Hybrid involving 'Red Torch', 'Rose Cup', 'Red Carpet' and 'Ming') R (1982) & N: A. Scharf. Red, with small yellow stripes. Stems 0.8m.

I (a/b) 'Carmen Cita'

cv. (Un-named, red, up-facing, asiatic seedling (no. 359/1) × 'Mont Blanc') H (1987), G (1989), N & I (1999): G. Thürmer REG: 2001. Inside light orange (28C); outside of outer petals deep pink (51B); outside of inner petals light orange (28D); numerous, dark red (187B) spots on basal ½ of each petal; nectaries orange-green; pollen deep red (185A). Fls 120mm wide; petals 65 × 30mm, not ruffled, tips slightly recurved. Lvs alternate, 70 × 10mm. Stems 0.8m, rather dark green, with stem bulbils and with 6-12 fls from very dark buds. Early to mid July. (Standard: colour print provided by registrant (WSY 0030945).)

I (a/c) 'Čaro'

cv. (('Wattle Bird' × ('Impact' × unknown)) × ('Accent' × 'Windahloo')) R & N: B. Mičulka I & REG: 2002. Inside pink with a narrow whitish rim, centre yellowish creamy, throat pale pink; small, dark brownish crimson brushmarks (smaller on outer petals) and moderately numerous, large, dark crimson brown spots on 2/5 of each petal; outside pale pink, midveins pale yellow and brownish; nectaries partially open, pink granulated; pollen ochre; stigma creamy. Fls 135mm wide; petals 75 × 40mm, slightly recurved, not ruffled; pedicels 60mm long, spotted. Lvs alternate, 90 × 21mm, dark green. Stems 0.85m, finely spotted brownish and with a moderately long inflorescence. Mid June.

VIII (b/a) 'Casa Rosa'

Syn. of 'Rote Horn' (see Ninth Supplement (1991) p. 25) (A *longiflorum*/asiatic hybrid resembling a pink *longiflorum* and sometimes treated as Div. V.

I (a-b/-) Chartreuse Giant Group

grp (*lanceifolium* var. *flaviflorum* × *wilsonii* var. *flavum*) R (pre-1982), N & I: Blackthorne Gardens. Yellow-green, with a few maroon spots in centre of flower. Fls c150mm wide. Lvs dark green. Stems 1.2-1.5m.

I (a/b) 'Citronkoks'

cv. (Parentage unknown) H (1991), G (1994), N (2001) & I (1997): J. Vasarietis REG: 2002. Inside brilliant yellow (7A); outside with a 10mm wide edge of brilliant yellow (7A) shading into brilliant yellow (8A) along the midveins; unspotted; nectaries light yellow; pollen brown; stigma greyish. Fls 160mm wide; petals 85 × 45mm, not ruffled, tips slightly recurved. Lvs alternate, 110 × 14mm. Stems 1.2m, green, with c10 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032315).)

VIII (b-c/b) 'Clearbrook'

cv. (('henryi' × un-named trumpet) × 'Canterbury Belle') H (1999) & G (2001); A.E. Alexander N, I & REG: 2002. Inside white, with a golden throat; outside white, flushed green at base; unmarked; nectaries green; pollen orange; stigma dull yellow. Fls 150mm wide; petals 100 × 35mm, not ruffled, tips recurved. Lvs alternate, 110 × 15mm. Stems 1.1m, black, with c8 fls in a racemose inflorescence. January (New Zealand). Ryrube Group. (Standard: colour print provided by registrant (WSY 0032056). Etymology: Clearbrook is a local place name where registrant lives in Christchurch, New Zealand.)

I (a/d) 'Collonia'

cv. ('Exception' (syn. 'Bakker's Pride') × 'Heidi') H (1986), G (1988), N & I (2002): G. Thürmer REG: 2001. Inside light orange (28C), tips light orange (28D), throat light orange (28C); outside light orange (28D); numerous, uniformly sized, dark red (183A) spots over at least basal ½ of each petal; nectaries greenish; pollen deep red (185A). Petals 70 × 30mm, not ruffled, tips strongly recurved. Lvs alternate, 110 × 10mm. Stems 1.0m, almost black, with c20 fls. Mid to late July. (Standard: colour print provided by registrant (WSY 0030949).)

VII (a/-) 'Colourful'

cv. (Parentage unknown) H, G, N (1993), I & REG (2002): J.W.A van der Wereld. Inside deep red (60A), towards apex deep purplish pink (64D), edged white (155D), midvein yellow, changing to red; outside pale purplish pink (62D); spots and papillae present; pollen orange-brown; stigma green; petal margins ruffled, tips recurved. Stems green, with dark stripes and spots. Midseason.

VII (-/b) Coral Mist Group

grp (Parentage unknown) R (pre-1982) & N: Strahm Bulb Farm. Pastel, coral-like colours: mostly pinks and crimsons, some lavenders and whites. Stems 1.0-2.2m. July-August.

I (a/-) 'Corona'

Syn. of 'Maureen' (Laan).

VII (b/-) 'Crazy Horse'

cv. (Parentage unknown) R: S. Hogewoning, (pre-1993), REG (2002): D. Kromhout & C. Hogewoning. Inside of inner petals very pale purple (73D), inside of outer petals light purplish pink (62C), base of midvein strong purplish red; outside white (155D); spots and papillae present; pollen light brown; stigma green. Petal margins ruffled, tips recurved. Stems with dark markings. Midseason to late.

I? (-/b) 'Crimson Chalice'

cv. (Parentage unknown) R (pre-1982) & N: Blackthorne Gardens. Garnet red.

VIII (-/a) 'Crimson Glory'

cv. (Aurelian seedling × sister seedling of 'Crimson Lady') R (pre-1982) & N: Y. Asano. Deep red (60A), edged white. Fls large, funnel-shaped. Late July. Triploid. (Unacceptable name (Art. 17.2).)

I (-/l) Crown Jewel Group

grp (Selections from Futura Group) R (pre-1982), N & I: Blackthorne Gardens. Colours range from gold to blood red. Stems to 1.0m. Early.

IX (b/b) 'Cruentum'

See *auratum* 'Cruentum'.

I (a/-) 'Darwin'

This name, now included as part of an EU PBR (no. 8445) has to take precedence over the previously registered 'Dublin' as the accepted name for this cultivar cf. *The International Lily Register, Nineteenth Supplement* (2001), p. 16.

I (b/-) 'Delight'

cv. (Parentage unknown) R (pre-1997) & N: Columbia-Platte Lilies. Inside deep

magenta rose; petals loosely reflexed. Stems 1.0m. Late June to July. (Unacceptable name (Art. 17.2).)

I (a-b/-) 'Delta'

cv. (Sundrop Group × 'Sterling Star') × yellow seedling with a brushmark)) H (1978), G (1981), N (1986) & I (1988): Bischoff Tulleken Lelies B.V. Inside vivid yellow (15A), with a small, spotted dark red (187B) brushmark and a vivid yellow (9B) throat; outside vivid yellow (15A), flamed light greenish yellow (3C); stigma dark red (59A); pollen brownish orange (165B); nectaries green. Fls 150mm wide; petals 75 × 25mm, not ruffled, tips slightly recurved. Lvs alternate, 125 × 16mm. Stems 0.9m, green, with darker markings and 6-10 fls. July. (Unacceptable name (Art. 17.2).)

I (a/-) 'Diana'

cv. ('Stroinaja' × mixed Mid-Century Group pollen) R (1975) & N: M.F. Kireeva. Fls black-red; pollen smoky violet. Stems 0.7-0.9m. June. (Unacceptable name (Art. 17.2).)

I (b/-) 'Diana'

cv. (*cernuum* × *lancifolium*) R (pre-1980) & N: R. Simonet & F. Fellner I: Borbeleta, 1980. Inside ivory white, with brown spots on the basal ½ of each petal; outside brownish; pollen brown. Petals broad, slightly recurved. Inflorescence large, pyramidal. (Published in Borbeleta catalogue, 1980. Unacceptable name (Art. 17.2).)

I (a/c) 'Doris Wharton'

cv. ('Geneve' × 'Merryspring') H (1997), G (1999) & N (2001): D.J. Hay REG: 2001. Inside pink with an orange gold blush from centre of each petal to base; fading to produce a cream stripe down centre of each petal; throat pink; outside pink; a few dark spots near the throat; nectaries pink; pollen rust orange; stigma pink. Fls 131mm wide; petals 63 × 35mm, not ruffled, tips very slightly recurved. Lvs alternate, 83 × 15mm, glossy. Stems 0.78m, dark green, with c7 fls. Midseason. (Standard: colour print provided by registrant (WSY 0031600).)

I (a/b) 'Dover'

cv. (Parentage unknown) R (pre-1988), N & REG (2002): M.A. van den Berg. Inside vivid orange (28A) at base, becoming more vivid reddish orange (30A) towards apex, throat orange; outside orange, becoming yellow towards apex, mid-veins green; markings black with purple-red margins; spots and papillae present; pollen brown; stigma purple. Petal margins ruffled, tips recurved. Stems dark green above with dark spots, base brown with green spots. Early.

I (a/b) 'Dresdener Gelbe'

cv. ('Connecticut King' × 'Redbird') H (1980), G (1982), N & I (1986): G. Thürmer REG: 2001. Inside vivid yellow (12A), with a reddish throat; outside vivid yellow (12A), inner petals with reddish midribs, outer petals darker; unmarked; nectaries greenish; pollen strong brown (172A). Petals 90 × 40mm, not ruffled, tips quite strongly recurved. Lvs alternate, 120 × 22mm, glossy. Stems 1.2m, rather dark green, with 12-15 fls. Early to mid July. (Standard: colour print provided by registrant (WSY 0030946).)

I (a/-) 'Dublin'

See note under 'Darwin'.

I (-/-) 'Duet'

cv. (Parentage unknown) **R** (pre-1996) & **N**: Columbia- Platte Lilies. Deep rose in bud, opening white with a deep rose centre; very few darker spots. Petals not ruffled, not recurved. Stems 1.0-1.2m. Late June. (Unacceptable name (Art. 17.2).)

VII (b-c/b) 'Earl's Pleasure'

cv. (*auratum* var. *platyphllum* × 'Joy' (syn. 'Le Rêve')) **H** (1993) & **G** (1999): K. Swoszowski-Tran **N**, **I** & **REG**; 2002. Inside pink rose fading to near white at tips, each petal with a deep magenta red band along midvein; outside pale pink to yellowish white; numerous, maroon, raised spots and some papillae on basal $\frac{1}{2}$ - $\frac{2}{3}$ of each petal; nectary chartreuse green; pollen brick red; stigma pale green; strongly scented. Fls 150mm wide; petals 110-130 × 45mm, margins slightly ruffled, tips slightly recurved. Lvs alternate, 140 × 40mm. Stems 1.3m, green, slightly marked purple, with 5-8 fls. Midseason to late. (Standard: colour print provided by registrant (WSY 0033599).)

IX (-/a) 'Easter Parade'

See *longiflorum* 'Easter Parade'.

VIII (b-c/-) 'Elegant Lady'

cv. (Tetraploid *longiflorum* 'Gelria' × *rubellum*) **H** (1994) & **G** (1996): CPRO **N** (2000), **I** & **REG** (2002): Coop. Kwekersver "The Originals" U.A. Inside light purplish pink/very pale purple (73C/D), base of midvein green changing to pink, throat pink; outside very pale purple (73D); no spots or papillae; nectaries green; pollen brown; stigma green. Fls 140mm wide; petals 150 × 60mm, margins slightly ruffled. Lvs alternate, 150 × 20mm. Stems 0.8m, green, with 4-6 fls. Early to midseason. (A *longiflorum*/oriental hybrid.)

I (a/c) 'Elise'

cv. (Parentage unknown) **R** (pre-1985) & **N**: A. Weinreich. Inside cream white, with small red spots and dashes; pollen brown. Pedicels long. Stems 1.2m. Mid July. (Published in German Lily Group sales list, 1985; unacceptable name (Art. 17.12).)

I (b/-) 'Elite'

cv. (Parentage unknown) **R** (pre-1974) & **N**: N. & B. Laan Bros. Inside vivid orange-yellow (21A), tips brilliant orange-yellow (21B), throat strong orange (24A); outside strong orange-yellow (24A); nectaries strong orange (24A); heavily spotted dark red (187B) over $\frac{1}{2}$ of each petal; pollen strong brown (172A). Fls 145mm wide; petals 85 × 30mm, tips slightly recurved, margins ruffled. Lvs alternate, 65 × 6mm. Stems 1.2m, dark green, with c10 fls. June.

I (a-b/b-c) 'El Moro'

cv. (Un-named seedling (no. LV15 ex Matthe) × 'Luxor') **H** (1991) & **G** (1994): G. Steinbrück **N**, **I** & **REG**; 2002. Inside vivid yellow (14A) with a strong, cream-coloured midrib, tips white (155A), throat vivid yellow (14A); outside white and yellow-white; numerous, small, rounded, strong red (46A) spots on the basal $\frac{1}{2}$ of each petal; nectaries light yellow-green (150D); pollen strong red (46A), filaments light orange-yellow (19A); stigma light yellowish pink (159A), style moderate yellow (161B). Fls 150mm wide; petals 87 × 40mm, not ruffled, tips slightly recurved. Lvs alternate, 90 × 15mm. Stems 1.1m, dark green with darker markings and c16 fls. Late June. (Standard: colour print provided by registrant (WSY 0032247). Etymology: El Moro is a fortification in Havana, Cuba and relates to the tower-like nature of the plant.

VI (-c) 'Eruption'

cv. ('Bright Star' × Golden Sunburst Group) R (pre-1979) & N: J.-M. Kaiser. Green at base, shading into orange and white. Stems 2.0m. Late July to August.

I (a/-) 'Erythro'

cv. (Parentage unknown) R (pre-1981) & N: M. Rohrmeier. Fls dark red. Late. (Published in German Lily Group sales list, 1982; unacceptable name (Art. 17.9).)

III (-/-) 'Europa'

cv. (*pyrenaicum* × *pomponium*) R (pre-1978) & N: C. North. Resembles a tall, vigorous *pomponium* with narrow leaves and scarlet flowers. (Distributed under code number 232/1.)

I (a/b) 'Evelyn'

cv. ('White Tiger' × 'Heidi') H (1986), G (1988), N & I (1999): G. Thürmer REG: 2001. Inside brilliant greenish yellow (6A), tips light greenish yellow (8C), throat brilliant greenish yellow (6A); outside brilliant greenish yellow (6A), near the midribs pale greenish yellow (9D), midribs reddish; deep red (185A), unequally-sized spots on lower ½ of each petal; nectaries greenish; pollen strong brown (172A). Petals 70 × 32mm, tips of outer petals strongly recurved, tips of inner petals only slightly recurved. Lvs alternate, 90 × 13mm. Stems 0.8m, rather dark green, hairy, with 7-10 fls. Early to mid July. (Standard: colour print provided by registrant (WSY 0030947).)

VII (-/b) 'Fairy'

cv. ((*speciosum* var. *rubrum* × *auratum*) × (*speciosum* var. *rubrum* × *auratum*, plus mixed *auratum* pollen)) R (1983) & N: W. Suzuki. Inside white (155D), base strong yellow-green (144B), strong red (53D) from midrib to near base; dark red (187B) spots conspicuous in basal ½ of each petal. Fls 160mm wide, strongly scented. Stems 1.0m. Midseason. (Unacceptable name (Art. 17.2).)

VII (-/d) Fairy Jewel Group

grp (Hybrids involving *rubellum* and 'Manawatu') H: L. Woodriff, pre-1966 S: L. Woodriff & J.M. Shaver. Inside apple-blossom pink to rose red, tips with a narrow white border. Stems 0.9-1.8m. August to September.

VII (-/b) Fairy Pink Group

grp (Parentage unknown) R (pre-1966) & N: L. Woodriff. Pink or rose, spotted dark pink or dark rose. Fls 125-175mm wide. Stems 0.8-1.5m. June to July.

I (a/-) 'Fantasie'

cv. (Parentage unknown) R (pre-1985) & N: VEG Zierpflanzen, Erfurt. Inside orange-red, with brown spots over ½ of each petal. Fls 115mm wide; petals 75 × 38mm. Stems 60cm. (Unacceptable name (Art. 17.2).)

VII (b/-) 'Farolito'

cv. (Un-named seedling × un-named seedling) H: Imanse Lelieveredeling B.V., 1991 N (1996), I (1996) & REG (2002): Peacock B.V. Inside strong purplish pink (73B), base of midvein white (155D), throat rose; outside strong purplish pink (73B); spots and papillae present; nectaries green; pollen orange-brown; stigma grey. Fls 180mm wide; petals 80 × 30mm, margins ruffled, tips recurved. Lvs alternate, 100 × 20mm. Stems 0.7m, green, with 2-6 fls. Very early.

VI (-/a) Fascination Group

grp (Selections from Marble Temple Group) **R** (pre-1966), **N** & **I**: J.M. Shaver. Pastel colours. Stems 1.5-2.1m. July to August. (Unacceptable name (Art. 26.1).)

I (a/b) 'Favola'

cv. (('Korál' × unknown) × 'Blizzard') **R** & **N**: B. Mičulka **I** & **REG**: 2002. Inside white, centre and throat tinged beige, very few spots; outside pale pink, with brownish midveins; nectaries green, white granulated; pollen orange; stigma brownish pink. Fls 125mm wide; petals 75 × 35mm, hardly recurved, not ruffled; pedicels 60mm long, shining brown. Lvs alternate, 95 × 11mm, pale green. Stems 0.75m, with numerous spots and a moderately long inflorescence. Early July.

I (c/d) 'Fedorov'

cv. (Parentage unknown, but close to *dauidii*) **G** (1980), **N** (1985) & **REG** (2002): A.M. Fedorov. Orange-red throughout, with some black spots; nectaries dark; pollen light red and orange; stigma wine red/dark red. Fls 65mm wide; petals 80 × 15mm, not ruffled, strongly recurved; lowest pedicels to 150mm long. Lvs alternate, 100 × 4mm. Stems 0.8m, green with up to 20 fls. Mid June to mid July.

VI (b-c/b) 'Fetiš'

cv. (Golden Sunburst Group × unknown) **R** & **N**: B. Mičulka **I** & **REG**: 2002. Inside white, centre and throat deep apricot, edges slightly veined pink, unspotted; outside with apricot glowing through, midveins slightly pinkish; nectaries green; pollen apricot brown; stigma crimson red. Fls 160mm wide; petals 110 × 33mm, tips very strongly recurved, not ruffled; pedicels 60mm long, green. Lvs alternate, 115 × 20mm, mid green. Stems 0.95m, slightly marked with violet brown and with a short inflorescence. Mid July.

I (b/-) 'Feuerball'

cv. (*lancifolium* hybrid × 'Red Tiger') **R** (pre-1981) & **N**: A. Winkler. Inside brick red, with dark spots. Stems to 1.3m, dark, with up to 15 fls and many bulbils. Late July.

I (a-b/-) 'Fidelity'

cv. (Parentage unknown) **R** (pre-1993), **G**, **N** (1993), **I** & **REG** (2002): J.W.A. van der Wereld. Inside vivid yellowish orange (30C); outside vivid yellowish orange (30C); spots and papillae present; nectaries orange; pollen orange-brown; stigma dark purple. Petal margins ruffled, tips recurved. Stems green, with dark stripes and spots. Midseason to late.

I (a/-) First Thanksgiving Group

grp (Parentage unknown) **R** (pre-1982) & **N**: Blackthorne Gardens. Apricot shades, some gold-banded. Stems 1.2-1.5m. August to October.

IX (-/a) 'Flevo'

See *longiflorum* 'Flevo'.

I (a-b/b) 'Formát'

cv. (Un-named white tetraploid × ('Milano' × 'Puchanta')) **R** & **N**: B. Mičulka **I** & **REG**: 2002. Inside pink, tips salmon pink, centre apricot with pink veining; moderately numerous, small spots over 2/5 of each petal and papillae in the throat; outside pastel pink, with pale brown midveins; nectaries white granulated; pollen red-brown; stigma beige. Fls 150mm wide; petals 100 × 48mm, not recurved, not ruf-

fled; pedicels 60mm long, very thick, strongly spotted dark brown, hairy. Lvs alternate, 130 × 19mm, dark green. Stems 0.9m, very thick, with pronounced dark brown speckling and a short inflorescence. Tetraploid. Mid June.

I (c/-) 'Fortiuna'

cv. ('Fire King' × *dauidii* var. *willmottiae*) R (c1975) & N: V.A. Volkov. Inside orange-red, with brown spots. Stems 1.8m. Late.

I (a/b) 'Fraboja'

cv. ('Milk and Honey' × 'Feuer und Rauch') R & N: J. Dudek I & REG: 2001. Inside lemon yellow, with a yellow throat; outside yellow with yellow-green ribs; very few spots; nectaries green; pollen orange-brown; stigma red-brown. Fls 180mm wide; petals 96 × 45mm, tips slightly recurved; pedicels 40mm long, dotted with brown. Lvs alternate, 140 × 70mm, glossy green. Stems 1.2m, rigid, brown. Mid July.

I (-/-) 'Fripita'

cv. (Parentage unknown) R (pre-1986) & N: F. Piasek. Inside pale yellow, with a red centre. Fs star-shaped. July.

VIII (b/b) 'Futura'

cv. (Un-named seedling (no. E-890-19) × un-named seedling (no. OR 90-058-8)) H, G, N & I (2000): World Breeding. Inside brilliant yellow (7B), with brownish orange (171B) near edges in basal half, tips brilliant yellow (7B), throat light yellow (10C); outside brilliant yellow (9C), with light yellow-green (145C) midveins and pale greenish yellow (9D) beside midveins, base with a deep purplish red (59B) spot; nectaries strong yellow-green (144B); very few spots on the outer petals only; no papillae; pollen strong orange (26A); stigma greyish purplish blue (103A). Fls 200mm wide; petals 140 × 65mm, not ruffled, tips slightly recurved. Lvs alternate, 250 × 25mm. Stems 1.25m, with c7 fls. (An orienpet lily; unacceptable name (Art. 17.2), but used in EU grant of PBR.)

I (-/-) Futura Group

grp (Complex hybrids, including *dauidii*) R (pre-1982), N & I: Blackthorne Gardens. Very varied colours.

VIII (a-b/-) 'Garibaldi'

cv. (Un-named seedling (no. AH 627-35) × un-named seedling (no. AH 3150)) H (pre-1996), G, N (1998), I & REG (2002): J.W.A. van der Wereld. Inside vivid yellowish orange (30C); outside vivid reddish orange (30B/C); unspotted but with papillae present; nectaries orange; pollen brown-red; stigma red-orange. Petal margins ruffled, tips recurved. Stems 1.15m, green, with dark stripes and spots. Midseason. (*A longiflorum*/asiatic hybrid.)

I (c/-) 'Gelber Goliath'

cv. (Unknown × Sutter's Gold Group) R (pre-1981) & N: A. Winkler. Inside gold-en yellow, strongly spotted. Stems 0.7m. Late July.

I (a/-) 'Gelber Stern'

cv. (Hybrid involving *wilsonii* and *lancifolium*) R (pre-1979) & N: G. Köpke. Inside yellow. Stems 0.6m, black, with up to 12 fls. July to August. (Published in German Lily Group sales list, 1980.)

VII (-/b) 'Genteel Lady'

cv. (Parentage unknown) R: unknown, pre-1982. Inside white, sprinkled with lavender spots; outside lavender pink. Fls 200mm wide; petal tips recurved. Stems 1.2m. August. (Listed in Rex Bulb Farms catalogue, 1982/1983.)

I (c/d) Giant Harlequin Group

grp (Parentage unknown) R: unknown, pre-1970. Varying from pink, pale lilac, old rose, red, violet, purple, ivory, white, yellow, salmon, tangerine to amber pink. Stems 1.5m, with up to 30 fls. June to July. (Listed in Rex Bulb Farms catalogue, 1970/1971.)

I (c/-) 'Gisela'

cv. ('Pink Champagne' × 'Connecticut King') R (pre-1989) & N: A. Weinreich. Inside dusky pink, with yellowish white centre and rather large spots over × of each petal and pale papillae on basal ½. Stems 1.5m, with some axillary bulbils. Early July.

I (b/b-d) 'Glockenklang'

cv. ('Dawn Star' × 'Summer Night') H (1989) & G (1992); G. Steinbrück N, I & REG: 2002. Inside brilliant orange-yellow (23B) throughout; outside brilliant orange-yellow (23B), reddish at base, with a strong greyish green midrib; numerous, round or streaked spots of strong red (46A) over basal ½ of each petal; nectaries light olive (152B); pollen strong red (46A); stigma strong red (46A). Fls 130mm wide; petals 75 × 35mm, not ruffled, tips slightly recurved. Lvs alternate, 95 × 14mm. Stems 0.85m, dark green, with 6-10 fls. Late June. (Standard: colour print provided by registrant (WSY 0032249).)

VIII (a/b) 'Glossy Wings'

cv. (Seedling no. S-6-85 × seedling no. 91.20.149-3) H (1993), G (1995), N (2000), I (2001) & REG (2002): World Breeding B.V. Inside strong purplish red (60B) on a strong purplish red (59D) ground, throat strong purplish red (63B); outside moderate purplish red (59C) on a strong purplish red (59D) ground, midveins suffused strong greenish yellow (151A); nectaries strong yellow-green (143C); very few spots and no papillae; pollen moderate reddish brown (166B); stigma moderate purplish red (58A). Fls 240mm wide; petals 130 × 50mm, tips recurved, not ruffled. Lvs alternate, 170 × 20mm. Stems 0.95m, green with darker markings and c7 fls. (A *longiflorum*/asiatic hybrid.)

VI (a-b/a) 'Glow Bug'

cv. (Golden aurelian seedling × apricot and green aurelian) H (1988), G (1990), N (2002) & I (2002): M.E. Rollinson REG: 2002. Inside creamy white (159D), with conspicuous brilliant orange-yellow (21B) throat with a flash of strong yellow-green (143C); outside creamy white (159D) with dark purplish pink (186C) suffusion along midveins; unmarked; nectaries strong yellow-green (145A); pollen deep orange-yellow; stigma greyish yellow-green (191B). Fls 125mm wide; petals 120 × 28-45mm, not ruffled, tips slightly recurved; strongly scented. Lvs alternate, 90 × 6mm. Stems 0.9-1.0m, moderate yellow-green (139C), flecked dark red (187A), with 15-20 fls. December to January (New Zealand). Syn. 'White Delight'. (Standard: colour print provided by registrant (WSY 0032035).)

I (c/d) 'Glutregen'

cv. (Parentage unknown) R (pre-1970) & N: C. Wittneben. Fls dark red, with reflexed petals.

I (-/-) Gold Banded Primrose Glory Group

grp (Selections from Futura Group) R, N & I (1982): Blackthorne Gardens. Fls primrose, banded gold.

VI (-/a) 'Goldcrest'

cv. (Parentage unknown) R (pre-1982) & N: L. Woodriff. Inside apricot, unspotted, the three inner petals with a median crest. Fls 125mm wide. Stems 1.1m with 10-12 fls. Late July. (Unacceptable name (Art. 17.2).)

I (-/-) Golden Goblet Group

grp (Parentage unknown) R (pre-1982) & N: Blackthorne Gardens. Bright yellow shades. Stems 0.8m. Early.

VII (-/-) 'Golden Harvest'

cv. (Parentage unknown) R & N: F. Jury I; M. Jury, 1998. Fls very large, white, each petal with a gold band. (Unacceptable name (Art. 17.2).)

VI (-/a) Golden Pearl Group

grp (*sargentiae* × yellow selections from Temple Group) R (pre-1969) & N: J.M. Shaver. Pastel gold. Stems 1.2-2.0m. Late July.

VI (-/a) 'Golden Spire'

cv. (Selection from Golden Splendor Group) R, N & I (1976): B.G. Hayler. Inside deep yellow; outside wine-coloured. Stems 1.9m.

VII (a-b/-) 'Golden Stargazer'

cv. (Parentage unknown) H: Van der Salm Bulbfarm Inc., pre-1997 REG: Van der Salm Holland B.V., 2002. Inside vivid yellow/light orange-yellow (16A/B); outside pale orange-yellow (16D); spots and papillae present; nectaries green; pollen light brown; stigma purple. Stems 0.9m. Late to very late.

VIII (a-b/-) 'Golden Tycoon'

cv. (Un-named seedling × un-named seedling) H: Testcentrum voor Siergewassen G (1998), N (2000), I (2001) & REG (2002): Marklily. Inside vivid yellow (13A/14A); base of midvein yellow-green shading to yellow; outside vivid yellow (13A/14A); nectaries green; pollen light brown; stigma yellow. Fls 150mm wide; petals 100 × 50mm, margins slightly ruffled, tips slightly recurved. Lvs alternate, 100 × 30mm. Stems 1.0m, green, with 2-6 fls. Early to midseason. (A *longiflorum* asiatic hybrid.)

VII (-/b) Golden Zephyrs

grp (Parentage unknown) R (pre-1979) & N: Strahm Bulb Farm. Inside white, with a gold band and crimson spots. Stems 1.0-1.9m. July to August.

I (a-b/c) 'Goldkleid'

cv. ('Yukatan' × un-named seedling (no. Fe4 ex Feldmaier)) H (1989), G (1992) & N (1998): G. Steinbrück I & REG: 2002. Inside light yellow (21D) throughout; outside light yellow (21D) throughout; not spotted; nectaries moderate yellowish green (138A); pollen dark reddish orange (172B); stigma brownish orange (164A). Fls 160mm wide; petals 80 × 30mm, not ruffled, tips becoming distinctly recurved. Lvs alternate, 90 × 17mm. Stems 1.2m, mid green, with c11 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032250).)

I (a/-) 'Goldmöve'

cv. (Derived from *lancifolium* var. *flaviflorum*, *wilsonii* var. *flavum* and *L. × maculatum*) R (pre-1980) & N: L. Denkewitz. Inside cadmium orange. Stems 1.1m. Mid July.

I (-/-) 'Goldstaub'

cv. (Parentage unknown) R (pre-1972) & N: C. Wittneben. Inside gold, glittering as if dusted with gold.

I (b/d) 'Goldstern'

cv. (Parentage unknown) R (pre-1970), N & I: S. Ziepke. Inside vivid yellow (12A), throat vivid orange-yellow (21A); outside vivid yellow (12A); nectaries white-green; heavily spotted dark red (187A/B) over $\frac{2}{3}$ of each petal; pollen strong orange (168B). Fls 100mm wide; petals 80 × 30mm, strongly reflexed, strongly ruffled. Lvs alternate, 100 × 13mm, dark green. Stems 0.85m, green, overlaid with purple and with c10 fls. July to August.

I (a/-) 'Governor Winslow'

cv. (Parentage unknown) H: R.W. Bemis, early 1960s G, N & I (1982): Blackthorne Gardens. Inside bright yellow, unspotted. Stems 1.0m. July to September.

I (a/-) 'Granát'

cv. (Parentage unknown) R (pre-1970) & N: J. Marks. Fls dark red.

I (a/-) 'Grand Commander'

cv. (Parentage unknown) R: unknown, pre-1998. Inside intense apricot, slightly spotted. Fls 150mm wide. Stems 0.9m. Late June. (Published in German Lily Group sales list, 1998; unacceptable name (Art. 17.2).)

I (a-b/-) 'Grease'

cv. (Parentage unknown) R, N (1993) & REG (2002): Hoffgaarde B.V. Inside pale purplish pink (62D), almost white at base, midveins moderate purplish red (184C) changing to very light pink; outside light purplish pink (62C); spots and papillae present; nectaries green; pollen orange-brown; stigma purple. Petal margins ruffled, tips recurved. Stems green, with dark stripes and spots. Midseason to late.

I (-/-) 'Green Wax'

cv. (Parentage unknown) R (pre-1982) & N: Blackthorne Gardens. Inside yellow-green, with a few crimson spots; no pollen. Stems and buds glossy.

VIII (b-c/b-c) 'Grenadine'

cv. (Atomic Hybrid seedling × (*auratum* × *henryi*)) H (1990), G (1993), N (2001) & REG (2001): R.H. Bazett I: B & D Lilies. Inside strong red to deep pink (51A/B), darker towards the midribs, throat strong red (46A); outside strong purplish pink (55B) shading lighter toward outer margins, midribs suffused mid green; numerous, small, dark red papillae covering basal $\frac{2}{3}$ of each petal; nectaries prominent, dark green, narrowly edged yellow; pollen vivid reddish orange (34B); stigma pale green; slightly scented. Fls 200mm wide; petals 110 × 50mm, margins ruffled, tips slightly recurved. Lvs alternate, 190 × 55mm. Stems 1.83m, grey-green, with c16 fls in a racemose inflorescence, including secondaries on lower pedicels. Late July. (Standard: colour print provided by registrant (WSY 0030920). An orienpet lily.)

I (b/-) Gurnet Light Group

grp (Parentage unknown) R (pre-1982) & N: Blackthorne Gardens. Inside golden yellow; pollen rusty red. Stems 0.8m. May to July.

VII (a-b/-) 'Hakuhou'

cv. (*nobilissimum* × *auratum*) R (pre-1989) & N: Japan. Inside white; outside white with green suffusion at base; pollen reddish brown. Petals slightly ruffled, tips slightly recurved. Lvs alternate, broadly lanceolate, dark green. Stems with c4 fls. (Listed in Miyoshi & Co. catalogue no. 87:106 (1989), with colour illustration.)

I (a/c) 'Hammamet'

cv. ('Ancona' × unknown) H (1989) & G (1991): G. Steinbrück N, I & REG: 2002. Inside strong orange (170A) mixed with pale yellow, throat somewhat paler; outside similarly coloured; numerous, moderate-sized, round and elongated, dark purple (79A) spots on all but tips of each petal; nectaries yellowish white (155D); pollen strong reddish orange (169A) in large anthers; stigma strong reddish orange (169A) on a pale green style. Fls 160mm wide; petals 80 × 30mm, not ruffled, tips not or very slightly recurved. Lvs alternate, 72 × 17mm. Stems 0.9m, dark green, with c7-10 fls. Late July to mid August (i.e. late for an asiatic). (Standard: colour print provided by registrant (WSY 0032251). Etymology: Hammamet is a town in Tunisia.)

VI (c/d) 'Havelidol'

cv. ('Henrieta' × 'White Henry') H (1990) & G (1994): G. Steinbrück N, I & REG: 2002. Inside moderate orange (167A) shading to pale orange-yellow (165D) at apex, throat moderate orange (167A); outside pale orange-yellow (165D); numerous, moderate-sized, pale red spots over $\frac{2}{3}$ of each petal and large papillae on basal part of each petal, coloured as the petal; nectaries deep yellowish green (141A); pollen dark reddish orange (175B) on long pale yellow-green (157C) filaments; stigma moderate reddish brown (176B) on a long style. Fls 95mm wide; petals 70 × 24mm, not ruffled, strongly recurved. Lvs alternate, unusually long, 250 × 30mm. Stems 2.0m, dark green with dark markings and c25 fls in a large inflorescence (50-60cm long); the long pedicels often two-flowered. Mid to late July. (Standard: colour print provided by registrant (WSY 0032254).)

I (b/b-d) 'Havelschau'

cv. ('Schwefelblüte' × 'Sinai') H (1989) & G (1992): G. Steinbrück N, I & REG: 2002. Inside vivid orange-yellow (23A) with a slight orange suffusion in the throat; outside vivid orange-yellow (23A), with moderate yellow-green midribs; unspotted; nectaries light greyish olive (195A); pollen brownish orange (166C); stigma light orange (28C). Fls 150mm wide; petals 90 × 40mm, not ruffled, tips strongly recurved. Lvs alternate, 90 × 15mm. Stems 0.6m, mid green, with 7-10 fls. Mid June. (Standard: colour print provided by registrant (WSY 0032255).)

I (c/-) 'Hedi'

cv. (Parentage unknown) R (pre-1977) & N: H. Dierssen. Inside orange. Stems 1.5m. Mid July. (Published in ELG selling list, 1997.)

I (a/b) 'Heidi III'

cv. (Parentage unknown) R (pre-1992) & N: H. Georgi. Fls orange-red.

I (b-c/d) 'Hektor'

cv. ('Pinegos Ares' × 'Tetra Tabasco') R & N: B. Mičulka I & REG: 2002. Inside

deep blood red, throat paler, with numerous, large, brown-red spots over $\frac{2}{3}$ of each petal; outside orange-red in centre of petals, midveins brownish; nectaries pinkish granulated, fairly papillose; pollen and stigma red chocolate. Fls 125mm wide; petals 75 × 40mm, strongly recurved; pedicels 85mm long, with pronounced brown markings, thick, hairy. Lvs alternate, 110 × 11mm, dark green. Stems 1.0m, thick, spotted very dark brown, with a moderately long inflorescence. Tetraploid. Late June.

I (-/d) 'Hel'

cv. (Parentage unknown) R (pre-1972) & N: C. Wittneben. Fls dark black-red; petals reflexed.

I (a/-) Hell's Blazes Group

grp (Parentage unknown) R (pre-1982) & N: Blackthorne Gardens. Shades of apricot. Stems 0.8m. July to August.

I (c/-) 'Henry P'

cv. (Parentage unknown) R: F.H. Payne (pre-1988) N: J. Wadekamper. Very light fawn/salmon tan throughout, with a few brown spots on lower $\frac{1}{2}$ of each petal; nectaries green; pollen light brown. Petals 100 × 25mm, not ruffled, slightly recurved. Lvs alternate, 110 × 15mm, mid green. Stems 0.9m, dark green, with 10-12 fls. July. (Published in Borbeleta Gardens catalogue, 1988.)

I (a/b) 'Hera'

cv. (Parentage unknown) R (pre-1984) & N: V.o.f. S. en S. REG: J.J. Schouten, 2002. Inside light/moderate pink (50D/51D) on a white ground, apex strong pink (54C), aging to light/pale purplish pink (55C/D), throat white (155A); outside of outer petals deep purplish pink (54B), deep/moderate pink (51C/D) along midveins, greenish on midveins; outside of inner petals strong pink (54C), toward apex deep purplish pink (54B), midveins greenish; spots and papillae present; pollen orange-brown; stigma red-purple. Petal margins ruffled, tips recurved. Stems mid green, with brownish purple spots.

I (a/c) 'High Note'

cv. (('Vanguard' × 'Attila') × un-named 1(a) white seedling) H (1994), G (1996), N (1998) & REG (2001): E.E. Fox I: Fox Lily Ranch, 2002. Inside light orange-yellow (22C) with dark red (187A) brushmarks just above the nectaries, throat light orange-yellow (22C); outside light orange-yellow (22D), with moderate orange-yellow (165C) midribs; dark red (187A) spots of moderate size on either side of brushmarks; nectaries frosted white; pollen vivid orange (28A); stigma strong reddish orange (31B). Fls 150mm wide; petals 75 × 22mm, not ruffled, tips not recurved. Lvs alternate, 140 × 22mm. Stems 1.25m, moderate olive green (147A), with 16-18 fls. Mid July. (Etymology: brushmark shape suggests a musical note and the lily is tall.)

VI (b/b) 'Hilton Beauty'

cv. (Pink trumpet seedling × yellow trumpet seedling) H (1996), G (1999), N (1999) & I: M.E. Rollinson REG: 2002. Inside vivid yellow (9A); outside vivid yellow (9A) with edges and midribs deep purplish red (61A); unmarked; nectaries strong yellow-green (144B); pollen vivid orange-yellow (23A); stigma light purple (75B). Fls 185mm wide; petals 123 × 36-55mm, margins slightly ruffled, tips slightly recurved; strongly scented. Lvs alternate, 112 × 15mm. Stems 0.95-1.0m, strong yellowish green (141C), with 10-15 fls. January (New Zealand). (Standard: colour print provided by registrant (WSY 0032038). Etymology: the Hilton Highway, Timaru, is

the location of a property of the registrant's, where most of his lily breeding has been done.

VI (b-c/a) 'Hilton Deluxe'

cv. (Parentage unknown) H, G (1997), N (1998) & I (1999); M.E. Rollinson REG: 2002. Inside dark purplish pink (186C), with a brilliant greenish yellow (2B) throat; outside dark purplish pink (186C); unmarked; nectaries strong yellow-green (144B); pollen strong orange (168A); stigma dark red (187A). Fls 125mm wide; petals 138 × 32-48mm, not ruffled, tips slightly recurved. Lvs alternate, 78 × 8mm. Stems 0.85-0.9m, pale green (189C), speckled dark purplish pink (184C), with 8 or more fls. January (New Zealand). (Standard: colour print provided by registrant (WSY 0032033). Etymology: see note under 'Hilton Beauty'.)

VI (b/b) 'Hilton Dream'

cv. (Un-named pink seedling × Copper King Group) H (1996), G (1999), N (1999) & I (1999); M.E. Rollinson REG: 2002. Inside dark purplish pink (186C), with a brilliant yellow (13B) throat; outside moderate purplish red (186B); nectaries brilliant yellow (13B); unmarked; pollen vivid orange-yellow (23A); stigma moderate purplish red (186A). Fls 174mm wide; petals 135 × 32-52mm, not ruffled, tips slightly recurved, slightly scented. Lvs alternate, 87 × 6mm. Stems 1.7-1.8m, pale green (189C), mottled dark red (59A), with 12-15 fls. January (New Zealand). (Standard: colour print provided by registrant (WSY 0032041). Etymology: see note under 'Hilton Beauty'.)

VI (b-c/a) 'Hilton Green'

cv. (Un-named seedling × 'Black Dragon') H (1994), G (1997), N (2001) & I (2002); M.E. Rollinson REG: 2002. Inside greenish white (155C), with a strong yellow-green (145A) throat; outside greenish white, tinged moderate purplish red (185C) on midveins, especially at base; unmarked; nectaries strong yellow-green (145A); pollen moderate orange (168C); stigma strong yellow-green (145A). Fls 155mm wide; petals 115 × 14-42mm, not ruffled, tips slightly recurved; very faintly scented. Lvs alternate, 85 × 13mm. Stems 1.2m, dark red (59A) with a grey undertone, with 8-15 fls. December (New Zealand). (Standard: colour print provided by registrant (WSY 0032039). Etymology: see note under 'Hilton Beauty'.)

VIII (a-b/-) 'Hitfever'

cv. (Parentage unknown) H (1997), G (1998), I (2001) & REG (2002); H.P. Imanse B.V. N: A.J. van der Velde, 1999. Inside vivid reddish orange/strong reddish orange (34B/169A); outside strong reddish orange (34C); spots and papillae present; nectaries brownish green; pollen orange-brown; stigma orange. Fls 160mm wide; petals 110 × 80mm, margins slightly ruffled, tips recurved. Lvs alternate, 150 × 30mm. Stems 1.0m, with dark spots and stripes and 5-7 fls. Early to midseason. (*A longiflorum*/asiatic hybrid.)

I (-/-) 'Holeas'

cv. (Parentage unknown) R (pre-1993), N & REG (2002); Hoffgaarde B.V. Inside a little stronger coloured than vivid yellow (12A), throat light greenish yellow (5C); outside a little stronger coloured than vivid yellow (12A), midveins brownish orange (166C) at base; dark red (187A) spots on basal part of each petal; pollen moderate reddish brown (175A); stigma dark red (187B). Fls 170mm wide; petals 90 × 40mm. Lvs alternate, 120 × 20mm. Stems 1.15m, pale green, with darker markings and c7 fls. Stems 1.5m. Very late.

I (a/c) 'Hot Tamale'

cv. ('Little Moons' × un-named, short, 1(a), red seedling) H (1994), G (1996), N (1999) & REG (2002): E.E Fox I: Fox Lily Ranch, 2002. Inside with circular vivid orange (28A) marks at mid petal, strong red (46A) tips and vivid red (46B) throat; outside vivid red (46C); very fine dark red (187A) spots over basal 1/3 of each petal; nectaries red, frosted white; pollen vivid orange (28A); stigma strong yellowish pink (32C). Fls 100mm wide; petals 50 × 30mm, not ruffled, tips not recurved. Lvs alternate, 70 × 12mm. Stems 35cm, mid green, with 9-12 fls. Late July. (Etymology: name suggested by hot colours.)

I (a-b/b) 'Hugo'

cv. (Parentage unknown) R (pre-1989), N & REG (2002): V.o.f. S. en S. Inside vivid yellow (12A), darker at base, throat yellow; markings purple-red, paler along margin; spots and papillae present; pollen orange-brown; stigma purple. Petal margins ruffled, tips recurved. Lvs mid green. Stems light green above, base brown and green, with darker markings. Very early to early.

VII (a/-) 'Ibiza'

cv. (Parentage unknown) H (1984), G (1987), N (1990) & REG (2002): Vletter & Den Haan Beheer B.V. Inside moderate to light purplish pink (65A/B), base of midvein white changing to pink; outside pale purplish pink (65D); nectaries green; spots and papillae present; stigma grey; pollen orange-brown. Stems 1.2m. Early to midseason. (The flowers are sometimes between up and out-facing.)

V (a-c/a) 'Imai'

cv. (Selection from Formolongi Group) R & N: Japan, pre-1989. Inside white, with a greenish throat; outside suffused very pale green; pollen pale orange-yellow; stigma white. Fls up- to slightly down-facing; petals not ruffled, tips recurved. Lvs alternate, rather narrow, finely pointed. Stems with 3-4 fls. (Listed in Miyoshi & Co. catalogue, no. 87:161 (1989), plus colour illustration.)

VI (b-c/a) 'Iowa Gold'

cv. (Un-named seedling × un-named seedling) H: Mrs D.B. Schaefer, 1993 G: Cebeco Lilies, 1996 N, I & REG: Mrs J. Venard, 2002. Inside vivid yellow (9A) inside and out, unmarked; nectaries deep yellowish green (141B); pollen light reddish brown (177B); stigma strong yellow-green (143C). Fls 180mm wide; petals 170 × 50mm, not ruffled, tips slightly recurved; strongly scented. Lvs alternate, 200 × 15mm. Stems 1.5m, moderate olive green (137B), with 8-10 fls in a racemose inflorescence. July. Tetraploid. (Standard: colour print provided by registrant (WSY 0035292). Etymology: refers to the State of origin and the colour of the flowers.)

VI (b-c/a) 'Iowa Orange-glow'

cv. (Un-named seedling × un-named seedling) H: Mrs D.B. Schaefer, 1993 G: Cebeco Lilies, 1996 N, I & REG: Mrs J. Venard, 2002. Inside white (155B) at tips shading to brilliant orange-yellow (21B) in throat; outside white (155B) with stripes of green and light orange; unspotted; nectaries strong yellowish green (135C); pollen moderate reddish brown (177A); stigma light greyish olive (197C). Fls 170mm wide; petals 130 × 40mm, not ruffled, tips slightly recurved; strongly scented. Lvs alternate, 185 × 15mm. Stems 1.7m, moderate yellowish green (139B) with 9-14 fls in a racemose inflorescence. July. Tetraploid. (Standard: colour print provided by registrant (WSY 0035288). Etymology: refers to the State of origin and the colour of the flowers.)

VI (b-c/a) 'Iowa Pink Blush'

cv. (Un-named seedling × un-named seedling) H: Mrs D.B. Schaefer, 1994 G: Cebeco Lilies, 1997 N, I & REG: Mrs J. Venard, 2002. Inside white (155D) edged with a blush of moderate purplish pink (186D), throat brilliant greenish yellow (6A); outside white (155D), with stripes of moderate purplish pink (186D) and moderate yellow-green (139D); unmarked; nectaries strong yellow-green (143C); pollen moderate reddish brown (166B); stigma light olive grey (197B). Fls 190mm wide; petals 140 × 50mm, margins not ruffled, tips slightly recurved. Lvs alternate, 200 × 15mm. Stems 1.0m, moderate yellow-green (138B), with 8-10 fls in a racemose inflorescence. July. Tetraploid. (Standard: colour provided by registrant (WSY 0035290). Etymology: refers to the State of origin and the colour of the flowers.)

VI (b-c/a) 'Iowa Sunbeam'

cv. (Un-named seedling × un-named seedling) H: Mrs D.B. Schaefer, 1993 G: Cebeco Lilies, 1996 N, I & REG: Mrs J. Venard, 2002. Inside brilliant yellow (11A), throat brilliant yellow (11A); outside brilliant yellow (10A), with green and white stripes; unmarked; nectaries strong yellow-green (141D); pollen moderate brown (200D); stigma strong yellow-green (143C). Fls 200mm wide; petals 170 × 55mm, margins not ruffled, tips slightly recurved; strongly scented. Lvs alternate, 230 × 15mm. Stems 1.9m, moderate yellowish green (137D), with 9-12 fls. July. Tetraploid. (Standard: colour print provided by registrant (WSY 0035291). Etymology: refers to the State of origin and the flower colour.)

VI (b-c/a) 'Iowa Sunset'

cv. (Un-named seedling × un-named seedling) H: Mrs D.B. Schaefer, 1993 G: Cebeco Lilies, 1996 N, I & REG: Mrs J. Venard, 2002. Inside light greenish yellow (4C) shading through dark yellow then to brilliant yellow (21C) in the throat; outside light greenish yellow (4C), with stripes of light orange and yellow; unmarked; nectaries strong yellowish green (141C); pollen brownish orange (165B); stigma moderate yellow-green (138C). Fls 170mm wide; petals 155 × 55mm, margins not ruffled, tips slightly recurved; strongly scented. Lvs alternate, 125 × 17mm. Stems 1.9m, moderate yellowish green (139B), with 8-12 fls. July. Tetraploid. (Standard: colour print provided by registrant (WSY 0035289). Etymology: refers to the State of origin and colour resemblance to an Iowa sunset!)

I (b-c/d) 'Irasmol'

cv. (*davidii* × 'Feuer und Rauch') R & N: J. Dudek I & REG: 2001. Inside red, centre of flower blackish red, with a red blotch, throat red; outside dark red, with red ribs; moderately numerous, small spots over ½ of each petal; pollen orange; stigma red. Fls 80mm wide; petals 65 × 25mm, strongly recurved; pedicels 120mm long, spotted with brown. Lvs alternate, 120 × 8mm. Stems 1.4m, brown, with a long inflorescence. Mid July.

I (a-b/b) 'Irene'

cv. (Parentage unknown) R, N (1990) & REG (2002): Peacock B.V. Inside vivid yellowish orange (30C), throat vivid reddish orange (30B); markings dark purple; outside of outer petals vivid reddish orange (30A); outside of inner petals vivid yellowish orange (30C); nectaries orange; spots and papillae present; pollen red-brown; stigma purple-red. Petals margins ruffled, tips recurved. Stems dark green with brown spots and some darker markings. Midseason.

I (b/-) 'Isabell'

cv. (Parentage unknown) R (pre-1982) & N: J. Klein. Salmon pink. Stems 0.8m.

Early July. (Published in German Lily Group sales list, 1986; unacceptable name (Art. 17.2).)

IX (-/a) 'Jean Paul'

See *longiflorum* 'Jean Paul'.

I (a/a) 'Jette'

cv. (Parentage unknown) R (pre-1989) & N: G. Steinbrück. Inside golden yellow, with red-brown spots and brushmark lines.

VI (b/a) 'Jimbo'

cv. (Parentage unknown) H (1990), G (1992) & N (2001): J. Ball REG; D. Nelson, 2001. Inside light yellow (12C), tips soft yellow (13D), throat green; outside light yellow (12C) with soft brown alongside the green midribs; unspotted; pollen bright orange. Fls 200mm wide; petals 160 × 80mm, not ruffled, tips slightly recurved, very glossy. Lvs alternate, 160 × 30mm. Stems 1.2-1.4m, dark green, with 3-6 fls. Mid July to mid August. (Standard: colour print provided by registrant (WSY 0031594).)

I (a/-) 'Jolly Green Giant'

cv. (Selection from Chartreuse Giant Group) R (pre-1982) & N: Blackthorne Gardens. Inside yellow-green, with darker spots; nectaries green. Fls up to 200mm wide. Lvs broad. Stems to 1.5m. August to September.

I (a/b) 'Jonithe'

cv. (Parentage unknown) R (pre-1986) & N: Gebr. Mak B.V. REG; Mak Lelieveredeling, 2002. Inside moderate red (179A)/dark red (187A), washed over a strong reddish orange (169A) ground in centre of petals and with paler margins, markings dark brown, throat moderate red (184B); outside moderate red (179A)/greyish red (178A); spots and papillae present; pollen dark orange-brown; stigma dark purple. Petal margins ruffled. Stems mid green at base, with some darker markings.

I (a/-) 'Juanita'

cv. ('Showboat' × ('Sun Ray' × un-named yellow seedling)) R (pre-1997) & N: Mak-Leek Inc. Inside burnt red-orange. Stems 0.6m, dark, with high bud count. Early. (Unacceptable name (Art. 17.2).)

I (a-b/b) 'Kadet'

cv. (('Ohnivák' × Red Flare Group) × 'Saulé') R & N: B. Mičulka I & REG; 2002. Inside deep crimson red, centre and throat very dark, narrow white edges; outside crimson red with centre of petals orange-red and midveins pale greenish yellow; sparse, small, black spots over ½ of each petal; nectaries partially open, white granulated; pollen ochre orange; stigma brown-black. Fls 125mm wide; petals 70 × 38mm, slightly recurved (outer more so), not ruffled; pedicels 80mm long, green. Lvs alternate, 80 × 17mm, pale green. Stems 0.85m, thick, green, with a moderately long inflorescence. Late June.

I (b-c/d) 'Kadina'

cv. ('Connecticut King' × 'Červená Karkulka') R & N: J. Dudek I & REG; 2001. Inside dark red, with an orange tinge, tips paler; outside dark red with red ribs; small, dense, black spots over ¾ of each petal; nectaries red; pollen brown; stigma dark red. Fls 85mm wide; petals 65 × 30mm, strongly recurved; pedicels 85mm

long, dotted with brown. Lvs alternate, 140 × 6mm, shiny dark green. Stems 1.3m, spotted with brown, hairy. Mid July.

I (c/d) 'Karibik-Traum'

cv. ('Sinai' × 'Avento') H (1989) & G (1992); G. Steinbrück N, I & REG: 2002. Inside light yellow, paling to yellowish white (158D) at tips; outside yellowish white (158D), with a soft green tinge; numerous, moderate-sized, dark red (59A) spots on basal ½ of each petal; nectaries moderate yellow-green (146D); pollen deep purplish red (187D) on long, pale greenish yellow (1D) filaments; stigma large, pale orange-yellow, on a thick, pale greenish yellow (1D) style. Fls 170mm wide; petals 90 × 42mm, not ruffled, tips strongly reflexed. Lvs alternate, 71 × 11mm. Stems 1.0m, dark green with black markings and c10 fls. Early July. (Standard: colour print provided by registrant (WSY 0032256).)

I (b/-) 'Karnaval'

cv. ('Aelita' × mixed pollen of 'Svetlana' and 'Zabava') R (c1982) & N: V.A. Volkov. Inside ruby red, with small spots. Stems 0.6m, with bulbils. June. (Unacceptable name (Art. 17.2).)

I (a/b) 'Karnevāls'

cv. ('Metelj' × 'Brushstroke') H (1993), G (1995) & N (1998); D.V. Škimele. Inside pale yellow with creamy tips and small, crimson red brushmarks, throat greenish yellow; outside light yellow, with light brownish red midveins; a few dark red spots; nectaries green; pollen light brown; stigma greenish brown. Fls 140mm wide; petals 75 × 40mm, not ruffled, tips slightly recurved. Lvs alternate, 70 × 10mm. Stems 0.8m, red-brown, with c5 fls. Mid July. (First published in *Lilium Balticum Lily Register*, 1998. Unacceptable name (Art. 17.12).)

I (a/-) 'Kassandra'

cv. ('Byam's Yellow' × 'Alpenglow') R (pre-1981) & N: V. Strasser. Inside golden yellow, spotted. Lvs dark green. Stems 0.8-0.9m, brown, with c25 fls. Early July.

I (c/-) 'Käthe'

cv. (Parentage unknown) R: J. Petruske N: P. Geiser, late 1970s. Fls creamy yellow. (Unacceptable name (Art. 17.12).)

I (b/-) 'Katharina'

cv. (Parentage unknown) R (pre-1994) & N: G. Patzold. Inside yellow, with an orange stripe. Stems 0.8m. (Published in European Lily Group sales list, 1994.)

I (b-c/d) 'Kimono'

cv. (Parentage unknown) R (pre-1987) & N: A. Weinreich. Inside golden yellow, heavily flushed reddish brown, pale throat and edges; outside reddish brown, outer petals with a flame of golden yellow towards apex, inner petals with a paler midvein; nectaries golden yellow; a few inconspicuous spots on basal ½ of each petal; stigma dark, style yellow; pollen dark brown, filaments yellow. Petals very slightly waved, tips distinctly recurved. Pedicels almost black. Stems with c5 fls.

IX (b/b) Kimono Group

See *auratum* Kimono Group.

I (a/-) 'King Phillip'

cv. (Parentage unknown) H: R.W. Bemis, 1960s N & I (c1982): Blackthorne

Gardens. Inside apricot/melon-coloured, with few spots. Lvs dark green. Stems 1.2m, mahogany. July to September.

VI (b/b) 'Kiri'

cv. (Parentage unknown) R (pre-1986) & N: G. Geering. Inside pale salmon orange, fading slightly towards base; outside orange, with green midribs, purplish at base; prominent green nectaries; anthers purplish, pollen orange; stigma pale green. Petals not ruffled, not reflexed. Stems with up to 6 fls. (Unacceptable name (Art. 17.2).)

I (a-b/-) 'Kirschroter Stern'

cv. (Parentage unknown) R (pre-1980) & N: C. Feldmaier. Inside dark red, with few spots. Lvs dark green. Stems 0.6m. Late June.

VIII (a/-) 'Kiss Me Kate'

cv. (Parentage unknown) L: Riverside Gardens, 2000. Fls glowing salmon rose, with a gold star centre. Stems 0.7m. Midseason. (A *longiflorum*/asiatic hybrid; unacceptable name (Art. 17.2).)

I (c/b-d) 'Kleine Eva'

cv. (Un-named seedling (no. AS12, ex Weidemann) × (yellow brushmark seedling × orange brushmark seedling)) H (1989), G (1992), N (1997) & I (2001): G. Steinbrück REG: 2002. Inside strong orange (170A), with a strongly developed double central rib; outside rather paler orange, with greyish yellow midribs; unspotted; nectaries yellowish grey (156A); pollen strong reddish orange (31A), filaments yellow-white; stigma dark purple (83A), on a yellow-white style. Fls 90mm wide; petals 70 × 24mm, not ruffled, tips strongly recurved. Lvs alternate, 70 × 24mm. Stems 0.8m, mid green, with c9 fls. Early to mid July. (Standard: colour print provided by registrant (WSY 0032257).)

I (a/-) 'Kleine Freude'

cv. (Parentage unknown) R (pre-1992) & N: G. Steinbrück. Fls blood red.

VI (-/a) 'Kognac'

cv. (Parentage unknown) R (pre-1995) & N: H. Böhm. Cognac-coloured trumpets. Stems 1.4m. Early July. (Published in German Lily Group sales list, 1995: unacceptable name (Art. 17.12).)

VIII (b/b) 'Kryisia'

cv. (Unknown × Longistar Group) H (1995), G (1997), N (2001) & I (2001): J. Kapias REG: 2002. Inside strong orange (25B), throat strong yellow-green (144B); outside moderate orange (168C); dark red (183B) spots on basal ½ of each petal; nectaries strong orange (25B); pollen dark greyish reddish brown (200A); stigma light orange-yellow (22D). Fls 220mm wide; 140 × 40-48mm, not ruffled, tips slightly recurved. Lvs alternate, 150 × 30mm. Stems 1.3m (under glass), greenish brown, with c15 fls. Mid July. Reported to be polyploid. (A *longiflorum*/asiatic hybrid. Standard: colour print provided by registrant (WSY 0031317).)

I (a/c) 'Kvaker'

cv. ('Ranger' × 'Amorette') R & N: B. Mičulka I & REG: 2002. Inside deep crimson red, centre and throat apricot; outside with green midveins; very numerous, large, dark chocolate spots over 3/5 of each petal and papillae in the throat; nectaries white frosted; stigma tinted violet. Fls 140mm wide; petals 75 × 42mm, moderately recurved; pedicels 100mm long, thick, pale brown. Lvs alternate,

105 × 16mm. Stems 1.0m, dark green, with a short, umbellate inflorescence. Late June.

I (a-b/b) 'Kvintán'

cv. (((('Quinella' × unknown) × ('Juno' × unknown)) × 'Majáles') × ('Windahloo' × 'Endeavor')) R & N: B. Mičulka I & REG: 2002. Inside lemon yellow, tips paler and tinted pale red; outside tinged pinkish and with deep yellow-green midveins; small brushmarks, like diffused mahogany blotches and numerous, large, dark brown-red spots over $\frac{2}{3}$ of each petal; nectaries green, white granulated; pollen orange-brown; stigma pinkish beige. Fls 105mm wide; petals 65 × 37mm, not recurved; pedicels 60mm long, strongly spotted. Lvs alternate, 80 × 8mm, dark green. Stems 0.45m, thick, slightly spotted and with a moderately long inflorescence. Early June.

VII (-/-) 'Kylie'

cv. ('Miss Bowen' × Bridal Bouquet Group) R (pre-1996) & N: V.E. Bowen. Inside white, with a sulphur yellow ray on basal $\frac{1}{2}$ of each petal above the green nectaries, continued apically as a purplish pink ray for a short distance; some purplish pink suffusion especially on outer petal edges; small, scattered, rounded spots over basal \times of each petal; pollen orange; stigma purplish. Petals strongly ruffled. (Etymology: named for the raiser's granddaughter.)

I (a/-) 'Kyoto'

cv. (Parentage unknown) R (pre-1992) & N: A. Weinreich. Inside pink, with very pale edges and a narrow, pale flash above nectaries; dark spots on basal $\frac{1}{3}$ of each petal. Petals slightly reflexed. (Unacceptable name (Art. 17.2).)

VI (b/a) 'Ladaho'

cv. ((*regale* × unknown) × unknown) R & N: J. Dudek I & REG: 2001. Inside creamy white, throat orange-yellow; outside creamy, with pink ribs; unspotted; nectaries green; pollen brown. Fls 120mm wide; petals 120 × 48mm, margins wavy, tips recurved; pedicels 90mm long, spotted brown. Lvs alternate, 72 × 8mm, dull green. Stems 1.45m, spotted with purple and with an umbellate inflorescence. Mid July.

VIII (b-c/b) 'Lady Di'

cv. ('Casa Blanca' × ('Starburst Sensation' × ('White Henry' × 'Dominique')))) H (1997), G (1999), N (2001) & REG (2001): R.H. Bazett I: B & D Lilies. Inside pale purplish pink (65C/D), shading to white at tips and margins, throat light purplish pink (65B); outside with inner surface pink showing through, midribs creamy white with a touch of green towards the base; none or very few, insignificant, deep pink spots; nectaries pale yellow-green; pollen strong orange (168B); stigma purple-green; slightly scented. Fls 160mm wide; petals 130 × 55mm, margins slightly ruffled, tips slightly reflexed. Lvs alternate, 260 × 37mm. Strong stems to 1.8m, pale grey-green, with up to 23 fls, including secondaries, in a pyramidal inflorescence. Mid July. (Standard: colour print provided by registrant (WSY 0030917). A orientpet lily.)

I (b/-) 'Laetitia'

cv. (Parentage unknown) R (pre-1992) & N: A. Weinreich. Inside yellow throughout, with a few reddish spots on basal $\frac{1}{3}$ of each petal.

VI (-/a) 'Laimrota'

cv. (Golden Clarion Group × mixed pollen of *L. × aurelianense* hybrids) H (1971), G (1975), N (1976) & I (1982): A. Zorģēvic. Inside light apricot, throat yellow-orange; outside light orange; nectaries light yellow; pollen light brown. Fls 120mm wide; petals 120 × 30mm, not ruffled, tips strongly recurved. Lvs alternate, 75 × 12mm. Stems 1.4-1.6m, dark green, with c8 fls. Late July to early August. (Published in *Dārzs un Drava*, no. 7 (1986).)

VII (a-b/-) 'Lapassa'

cv. (Un-named seedling × 'Joy' (syn. 'Le Rêve')) R: Imanse Lelieveredeling B.V., 1992 N (1998), I (1998) & REG (2002): Peacock B.V. Inside rose (moderate purplish pink (65A) on a white (155A) ground), midveins yellow at base, throat rose; outside pale purplish pink (65C); spots and papillae present; nectaries greenish; pollen yellow-brown; stigma dark purple. Fls 200mm wide; petals 120 × 30mm, margins ruffled, tips slightly recurved. Lvs alternate, 120 × 40mm. Stems 0.95m, brown and green, with 2-6 fls. Early to midseason.

I (-/-) Lavender and Lemon Glory Group

grp (Selections from Futura Group) R, N & I (1982): Blackthorne Gardens. Fls lavender and lemon.

I (b/d) 'Lemon Pudding'

cv. (Un-named, tetraploid, yellow seedling × ('Ivory Sands' × 'Donna Sylvester')) H (1991), G (1994), N (1996) & REG (2001): E.E. Fox I: Fox Lily Ranch, 2002. Inside brilliant greenish yellow (6A), throat brilliant greenish yellow (6C); outside brilliant greenish yellow (6C); unspotted; nectaries brilliant greenish yellow (6C), frosted white; pollen moderate reddish brown (174A); stigma dark red (183A). Fls 140mm wide; petals 70 × 40mm, not ruffled, tips strongly recurved. Lvs alternate, 100 × 7mm. Stems 1.0m, mid green, with 12-16 fls. Mid July. (Etymology: name suggested by colour of fls.)

I (a/b) 'Lēnīgā'

cv. (Parentage unknown) H (1991), G (1994), N (1997) & I (1997): J. Vasarietis REG: 2001. Inside deep purplish pink (55A), with a pale purplish pink (55D) throat; outside deep purplish pink (55A) on edges, strong purplish pink (55B) around the midveins; sparse, moderate-sized, dark spots on basal ¼ of each petal; nectaries white; pollen orange brownish; stigma pale greyish. Fls 150mm wide, on long pedicels; petals 80 × 45mm, not ruffled, tips slightly recurved. Lvs alternate, 120 × 8mm. Stems 0.95m, green, with c15 fls. Early July. (Published in *Lilium Balticum Lily Register*, 2000. Standard: colour print provided by registrant (WSY 0030813). Etymology: Lēnīgā in Latvian means a person who is gentle, mild, meek or soft.)

I (-/-) 'Leopard'

cv. (Selection from Egmont Group) R (pre-1970), N & I: R.E. Harrison. Inside soft lemon, spotted brownish maroon. Stems 0.6-1.0m. (Unacceptable name (Art. 17.2).)

I (a/b) 'Liberation'

cv. (Parentage unknown) R: unknown, pre-1985. Inside clear pink, with a few maroon spots around centre of fls. Stems 0.6m. June to July.

I (a-b/b) 'Lidka'

cv. ('Grun-weiss' × ('Blizzard' × ('Leandra' × 'Windahloo'))) R & N: B. Mičulka I

& REG: 2002. Inside creamy, centre yellowish, throat white; outside with greenish and brownish midveins; moderately numerous, large, dark brownish pink spots over $\frac{1}{2}$ of each petal; nectaries green, white granulated; pollen brown; stigma dark brown. Fls 143mm wide; petals 85 × 38mm, very slightly recurved; pedicels 70mm long, slightly brown marked, slightly hairy. Lvs alternate, 70 × 14mm, dark green. Stems 0.85m, thick, with fine brown spots, fairly hairy and with an umbellate inflorescence. Late June.

I (c/-) 'Lilac Wonder'

cv. (Selection from Harlequin Group) R (pre-1972) & N: N. & B. Laan Bros. Inside dark rose.

I (-/-) Lime Glory Group

grp (Selections from Futura Group) R, N & I (1982): Blackthorne Gardens. Fls lime green.

I (a/c) 'Little Moons'

cv. (Un-named red and yellow bicolor seedling × un-named seedling with red fls with a yellow streak by nectary) H (1991), G (1993), N (1996) & REG (2001): E.E. Fox I: Fox Lily Ranch, 2002. Inside with vivid yellow (9B) circular markings at mid petal, with vivid red tips (45B) and a strong red (46A) throat; outside strong red (45D); unspotted; nectaries strong red (46A), frosted white; pollen moderate reddish brown (166B); stigma greyish brown (166A). Fls 120mm wide; petals 60 × 27mm, not ruffled, tips not recurved. Lvs alternate, 100 × 12mm. Stems 1.0m, pale green, with 28-38 fls. Late July. (Etymology: name suggested by the yellow circular markings, like little moons.)

I (-/-) 'Loki'

cv. (Parentage unknown) R (pre-1972) & N: C. Wittneben. Fls dark black-red.

IX (-/a) longiflorum 'Easter Parade'

cv. (Selection from *longiflorum*) R (pre-1982) & N: Blackthorne Gardens. Fls pure white. Stems 0.6m. August.

IX (-/a) longiflorum 'Flevo'

cv. (Selection from *longiflorum* Indian Summer Group) R (c1981) & N: J. Boon. Fls pure white, with lemon yellow anthers.

IX (-/a) longiflorum 'Jean Paul'

cv. (Selection from *longiflorum* Indian Summer Group) R (c1981) & N: J. Boon. Fls pure white, with buttercup yellow anthers. (Sometimes seen as 'Jan Paul'.)

IX (b-c/a) longiflorum 'White Satin'

cv. (Parentage unknown) R, N & REG (2002): J.W.A. van der Wereld B.V. Inside white (155B) with a white (155A) base; outside similarly coloured; nectaries green; no spots or papillae; pollen orange-yellow; stigma light green to almost white. Petal margins slightly ruffled, tips recurved. Late.

VIII (b/b) Longistar Group

grp (*longiflorum* × 'Sterling Star') H, G & N (1983): P.C. Schenk I: Borbeleta Gardens, 1984. Inside greenish cream; outside greenish; a few scattered brownish spots on lower $\frac{1}{3}$ of each petal. Petals not ruffled, tips not recurved, very brittle. Lvs alternate, mid green. Stems 0.9m, mid green, with up to 19 fls. July. (Published in

Borbeleta Gardens catalogue, 1984; individual clones were sold by Borbeleta as 'Longistar I', 'Longistar II' etc.)

VII (a/-) 'Looking Up'

cv. (Parentage unknown) L: Borbeleta Gardens, 1994. Inside soft pink. Fls up-facing. Stems 0.6m. Early.

I (a/b) 'Lovedream'

cv. (Parentage unknown) R: W. Paauw & Zn B.V. pre-1987 REG: Peacock B.V., 2002. Inside strong orange (39A) in centre of petal, apex deep yellowish pink (39B), spot at base strong orange (24A), markings brown, throat pink/orange; spots and papillae present; pollen red-brown; stigma purple. Petal margins not ruffled, tips recurved. Stems brown to mid green, with darker markings.

I (a-b/-) 'Lugos'

cv. (Parentage unknown) R (pre-2000), N, I & REG (2002): World Flower B.V. Inside strong purplish red (54A)/deep purplish pink (58D), paler at apex, base white; outside deep purplish pink (58D); nectaries green; spots and papillae present; pollen brown; stigma purple. Petal margins ruffled, tips recurved. Stems 0.95m, green, with dark stripes and spots. Very early to early.

I (a/b) 'Madeira'

cv. (Parentage unknown) R (pre-1985), N & REG (2002): G.C. van der Salm. Inside vivid reddish orange (30A/33B), with dark brown markings and a brilliant orange (29A) throat; outside strong reddish orange (31A/B), with vivid reddish orange (30A/32A) along margins and green midveins; spots and papillae present; pollen orange-brown; stigma red-purple. Petal margins not ruffled, tips slightly recurved. Stems brown, with mid green spots and darker markings.

I (b-c/d) 'Mahalak'

cv. ('Červená Karkulka' × 'Hornback's Gold') R & N: J. Dudek I & REG: 2001. Inside red, centre of flower and throat orange-red; outside red; moderately-sized, dense, black spots over $\frac{3}{4}$ of each petal; nectaries red; pollen orange-brown; stigma black-red. Fls 95mm wide; petals 68 × 30mm, strongly recurved; pedicels 100mm long, brown, with secondary buds. Lvs alternate, 140 × 10mm, glossy green. Stems 1.4m, rigid, spotted with brown, brown at base. Mid July.

VII (-/b) 'Majja'

cv. (Parentage unknown) R (c1965) & N: K. Zeltan. Inside rose, with white margins and dark raspberry spots; anthers orange. Fls up to 200mm wide, scented. Stems 1.2m. Late August. (Unacceptable (Art. 17.2).)

I (c/d) 'Majaslada'

cv. (*dauidii* × 'Feuer und Rauch') R & N: J. Dudek I & REG: 2001. Inside red, throat orange; outside orange-red, with orange ribs; moderately dense, large, black spots over $\frac{1}{2}$ of each petal; nectaries dark red; pollen red-brown; stigma dark red. Fls 80mm wide; petals 70 × 35mm, strongly recurved; pedicels 110mm long, spotted with brown and with secondary buds. Lvs alternate, 140 × 6mm. Stems 1.4m, rigid, spotted with brown. Mid July.

VII (-/a-b) Majestic Group

grp (Oriental trumpets × *henryi* hybrids) R (pre-1966) & N: R.W. Bemis. Varying from white or yellow to pink or purple. Stems 1.2-1.9m. July. (Listed in Blackthorne Gardens catalogue, 1966/1967; unacceptable name (Art. 26.1).)

I (b/c) 'Make Up'

cv. (Parentage unknown) R (pre-1991) & N: A. Weinreich. Inside bicoloured: the apical half of each petal clearly separated from basal red half; lightly spotted. Petals only slightly reflexed. Stems 1.2-1.5m. Mid July. (Published in German Lily Group sales list, 1991; unacceptable name (Art. 17.2).)

I (-/-) 'Malcolm'

cv. (Believed to be a sister seedling of 'Wonga') R: G.M. Chandler, pre-1983 N & I: M.K. Baldwin. Inside bright yellow, lightly spotted at centre of flower. Late December (Australia).

II (-/-) 'Mandarin'

cv. (*hansonii* × *tsingtauense*) R (pre-1987) & N: O. Beutnagel. Inside orange, with a few spots. Stems 1.2m. Late July. (Published in German Lily Group sales list, 1987; unacceptable name (Art. 17.2).)

VII (a/-) 'Mandela'

cv. (Parentage unknown) R: Gebr. Mak B.V., pre-1990 REG: Van den Bos Flowerbulbs, 2002. Inside strong to moderate purplish pink (68B/C), midveins strong red (53B), markings red; outside very pale (73D); spots and papillae present; nectaries brilliant yellow-green (149B), with yellow edges; pollen orange-brown; stigma purple with grey. Petal margins strongly ruffled, tips recurved. Stems with dark markings. Very late.

I (a/b) 'Mandy'

cv. (Un-named seedling × un-named seedling) H (1992), G (1995), I (2001) & REG (2002): W. Lighthart N: M. van Vlier, 1999. Inside deep yellowish pink (39B), more orange along margins, throat orange-red; outside strong red (39A)/strong yellowish pink (35C); nectaries orange; spots and papillae present; pollen orange-red; stigma orange-yellow. Fls 140mm wide; petals c70 × 40mm, margins slightly ruffled, tips slightly recurved. Lvs alternate, 80 × 30mm. Stems 1.05m, with darker markings and c6 fls. Early to midseason.

VII (a-b/-) 'Manzini'

cv. (Un-named seedling × 'Joy' (syn. 'Le Rêve')) R: Imanse Lelieveredeling B.V., 1993 N (1998), I (1998) & REG (2002): Peacock B.V. Inside strong purplish pink (73B), midveins yellow at base, throat rose/white; outside strong purplish pink (73B); nectaries yellow-green; spots and papillae present; pollen light to dark brown; stigma dark purple. Fls 180mm wide; petals 110 × 40mm, margins ruffled, tips recurved. Lvs alternate, 100 × 30mm. Stems 0.9m, brown and green, with 2-6 fls. Early to midseason.

I (a/b) 'Margaret Hay'

cv. ('Geneve' × 'Merryspring') H (1997), G (1999) & N (2001): D.J. Hay REG: 2001. Inside ivory white, with a soft butter yellow overtone, throat ivory white; outside white with a pink blush; a few, small, dark spots near the throat; nectaries ivory white; pollen rust-coloured; stigma ivory/pink. Fls 142mm wide; petals 72 × 36mm, margins slightly ruffled, tips not or very slightly recurved. Lvs alternate, 73 × 14mm.

Stems 0.8m, green, with c9 fls. Midseason. (Standard: colour print provided by registrant (WSY 0031598).)

I (b-c/-) 'Maria'

cv. (Parentage unknown) R (pre-1989) & N: A. Weinreich. Fls pale cream, with very broad petals; slightly scented. Stems 1.0m. Early July. (Unacceptable name (Art. 17.2).)

I (a/-) 'Marinka'

cv. ('Rubinovaja' × 'Vostok-2') R (pre-1975) & N: M.F. Kireeva. Inside bright orange-red, spotted; pollen orange-brown. Stems 1.0m, with bulbils. Early.

I (a/-) 'Marissa'

cv. (Parentage unknown) R (pre-1998) & N: Van der Salm. Blend of peach, gold and apricot. Stems 1.0m. Midseason. (Unacceptable name (Art. 17.2).)

VII (a-b/-) 'Marrero'

cv. (Un-named seedling × un-named seedling) H & G (1995): Peacock B.V./Imanse Lelieveredeling B.V. N: Peacock B.V., 1997 I (2000) & REG (2002): Marklily. Inside vivid purplish red (57D), fading with age, base of midvein white, throat white; outside moderate purplish pink (66D); spots white/red; papillae present; pollen orange-brown; stigma dark purple. Fls 200mm wide; petals 170 × 50mm, margins ruffled, tips recurved. Lvs alternate, 150 × 50mm. Stems 1.0m, green, with 2-8 fls. Midseason to late.

IX (b/d) *martagon* var. *cattaniae* 'The Moor'

cv. (Selection from *martagon* var. *cattaniae*) G (c1992): M.C. Foster N, I & REG: 2002. Inside blackish maroon (darker than 187A), base moderate purplish red (186A), waxy and glossy; outside reddish purple (darker than 187A), central ridge redder than deep reddish purple (77A); no spots or papillae; nectaries green; pollen brownish orange (172C), filaments more-or-less moderate red (184B); stigma dark red (187A), style dark red (187B); ovary green. Fls 35-40mm wide; petals 27 × 12-15mm, margins slightly waved, strongly reflexed. Lvs whorled, 118-170 × 22-50mm. Stems 1.0-2.0m, mottled and streaked brown, with 9-28 fls. June to July. PC 2002. (Grown from seed germinated c1986. Etymology: an allusion to the black Shakespearean character Othello, occasioned by the unusually deep colour of the flower. Standard: herbarium specimen (WSY 0035276) supplied by registrant.)

VI (b-c/b) 'Matai Delight'

cv. (Un-named seedling × un-named seedling) H (c1993), G (1999), N (2001) & I (2001): M.E. Rollinson REG: 2002. Inside white (155B), well blushed with moderate purplish red (58A), tips strong yellow-green (144B), throat brilliant yellow-green (150A); outside moderate purplish red (58A) with white (155B) showing through; unmarked; nectaries brilliant yellow-green (150C); pollen strong orange-yellow (22A); stigma moderate purple (79D). Fls 185mm wide; petals 148 × 38-56mm, not ruffled, tips slightly recurved; strongly scented. Lvs alternate, 175 × 16mm. Stems 1.5-1.6m, greyish yellow-green (191B), mottled moderate purplish red (58A), with 10-15 fls. December to January (New Zealand). (Standard: colour print provided by registrant (WSY 0032040). Etymology: Matai is the name of the road where the registrant lives in Timaru, New Zealand.)

VI (b-c/b) 'Matai Glow'

cv. (Un-named seedling × Copper King Group) H (1991), G (1994), N (1996) & I (2000): M.E. Rollinson REG: 2002. Inside light orange-yellow (22D), edges

blushed deep pink (48A), the pink colouration fading after 4-5 days; outside strong red (53B) with light orange-yellow (22D) streaking; inside unmarked; nectaries light yellow-green (145D); pollen vivid orange-yellow (23A); stigma strong red (53B). Fls 160mm wide; petals 115 × 55mm, not ruffled, tips strongly recurved; strongly scented. Lvs alternate, 145 × 13mm, dark yellowish green (189A), speckled dark red (187A), with 8-12 fls. December to January (New Zealand). (Standard: colour print provided by registrant (WSY 0032034). Etymology: Matai is the name of the road where the registrant lives in Timaru, New Zealand.)

VII (a-b/-) 'Matera'

cv. (Parentage unknown) R (pre-1988), N & REG (2002): Vletter & Den Haan Beheer B.V. Inside strong purplish red (63B), midveins deep/moderate red (185A/B), markings dark red-purple; outside strong purplish pink (63C); spots and papillae present; pollen light brown; stigma purple with grey. Petal margins strongly ruffled, tips recurved. Stems with darker markings. Late to very late.

I (a/-) 'Maureen'

cv. (Parentage unknown) R (pre-1980) & N: N. & B. Laan Bros. Inside salmon orange. Stems 1.3m. Early July. Syn. 'Corona'.

VII (-/-) 'Milady'

cv. (Parentage unknown) R (pre-1997) & N: F. Jury. Fls very large, very pale lavender pink, with a green centre.

VIII (a/-) 'Milagro'

cv. (Un-named seedling × un-named seedling) H: Testcentrum voor Siergewassen G, N, I & REG (2002): Marklily. Inside strong orange (24A/25A); outside strong orange (25B); nectaries orange; spots and papillae present; pollen orange-brown; stigma orange. Fls 150mm wide; petals 100 × 50mm, margins slightly ruffled, tips slightly recurved. Lvs alternate, 150 × 20mm. Stems 0.85m, green, with 3-6 fls. Early to midseason. (*A longiflorum*/asiatic hybrid.)

VIII (b-c/a) 'Millennian Dawn'

cv. ('Canterbury Belle' × 'Leslie Woodriff') H (1995), G (1998) & N: A.E. Alexander REG: 2002. Inside white, flushed with pink and with a gold throat; outside pink and white; unmarked; nectaries green; pollen orange; stigma frosty grey. Fls 110mm wide; petals 120 × 50mm, not ruffled, tips recurved. Lvs alternate, 150 × 15mm. Stems 1.8m, green, with c14 fls in a racemose inflorescence. January (New Zealand). (Standard: colour print provided by registrant (WSY 0032049). An orienpet lily.)

VIII (b/a) 'Millennian Miss'

cv. ('Canterbury Belle' × 'Leslie Woodriff') H (1995), G (1998), N (2001) & I (2001): A.E. Alexander REG: 2002. Mid pink inside and out, with white tips and a gold throat; unmarked; nectaries green; pollen orange; stigma purple-black. Fls 105mm wide; petals 90 × 45mm, not ruffled, tips recurved. Lvs alternate, 110 × 10mm. Stems 1.3m, light green, overlaid with wine-coloured flecks, with c12 fls in a racemose inflorescence. Mid to late January (New Zealand). (Standard: colour print supplied by registrant (WSY 0032048). An orienpet lily.)

VII (-/c) 'Mirage'

cv. (*speciosum* × *auratum*) × *speciosum* 'Album Novum') R (c1978) & N: A.N. Ungerman. Inside white, slightly spotted pink in centre of flower. Fls up to 180mm wide. Stems 1.0m. August.

I (a/-) 'Miss Blanche'

cv. (Parentage unknown) R (pre-1986) & N: Fa. P. Hoff & Zn REG: Hoffgaard B.V., 2002. Inside light yellow-green/light greenish yellow (2C/3C) at base and in throat, apex paler than pale greenish yellow (2D), markings dark brown; outside of inner petals pale greenish yellow/pale yellow-green (2D/4D) with margins at base light greenish yellow (3C/D); outside of outer petals pale greenish yellow/light greenish yellow (2D/3C), apex light greenish yellow (3C); spots and papillae present; pollen light orange-brown; stigma red-purple. Petal margins not ruffled, petals recurved. Stems light green, with darker markings.

VI (b/a) 'Mizar'

cv. ('Kord' × 'Andrea') R & N: K. Braun I: 2001 REG: 2002. Inside pastel orange with paler edges and a green throat; outside orange, green at base and with orange ribs; unspotted; nectaries green; pollen chocolate; stigma pale green. Petals 130 × 43mm, slightly waved; pedicels 100mm long, pale green. Lvs alternate, 115 × 15mm. Stems 1.25m, brownish green, with a moderately long inflorescence. Mid July.

I (b-c/c) 'Mokka'

cv. (((('Nutmegger' × 'Connecticut Red') × 'Kytice') × unknown) × (('Potstar' × 'Stellar Gleam') × 'Majáles')) R & N: B. Mičulka I & REG: 2002. Inside reddish brown on a bronze ground, narrow bronze rim and bronze around nectaries; outside with brownish white centre and yellowish midveins; numerous, moderately sized, dark red-brown spots over ½ of each petal; nectaries slightly open, finely white granulated; pollen ochre brown; stigma brownish. Fls 100mm wide; petals 58 × 40mm, slightly recurved; pedicels 60mm long, lightly spotted. Lvs alternate, 70 × 10mm, dark green. Stems 0.85m, thick, green, with a moderately long inflorescence. Mid June.

I (a/b-c) 'Monique'

cv. (Un-named seedling × un-named seedling) H (1990), G (1992), N (2002) & REG (2002): R.H. Bazett I: The Lily Nook, 2002. Inside strong purplish red (63B) shading to strong purplish pink (63C), paling in centre towards a pale yellow (11C) throat; outside deep purplish pink (64D) with some green on the midribs; dark purple (79A), elongated spots (c2mm long) on basal 40mm of each petal; nectaries pink to pale green; pollen moderate orange (167A); stigma dark purple. Fls 180mm wide; petals 100 × 50mm, not ruffled, tips slightly recurved. Lvs alternate, 155 × 19mm. Stems 1.4m, dark green, with c10 fls. June to July. Tetraploid. (Standard: colour reproduction (WSY 0035137) provided by registrant.)

I (-/-) Monostrahlen Group

grp (Parentage unknown) R (pre-1970) & N: C. Wittneben. Inside orange-yellow, with or without spots.

I (a/-) Moonsong Group

grp (Parentage unknown) R: unknown, pre-1982. Inside yellow, unspotted. Stems 1.0-1.2m. June to July.

I (a/-) 'Moorhexe'

cv. (Parentage unknown) R (pre-1986) & N: C. Wittneben. Inside bright red. Stems 0.8m. Late July. (Published in German Lily Group sales list, 1986; unacceptable name (Art. 17.2).)

I (a-b/-) 'Morgenrot'

cv. (Parentage unknown) R (pre-1980) & N: C. Feldmaier. Inside vivid red (45A), with brown spots and white nectaries. Stems 0.7m. July.

I (a/b) 'Morgentau'

cv. ('El Paso' × 'Colleen' (syn. 'Jeanne d'Arc')) H (1991), G (1994), N (2001) & I (2001): G. Steinbrück REG: 2002. Inside white (155A), with soft orange suffusion; outside similarly coloured; no spots; nectaries greyish green; pollen dark red (187A). Fls 125mm wide; petals 70 × 39mm, not ruffled, tips not recurved, with a strong double rib down centre line. Lvs alternate, 90 × 12mm. Stems 0.95m, dark green, with 7-10 fls. Mid to late June. (Standard: colour print provided by registrant (WSY 0032258).)

VI (b/a) 'Naomi'

cv. (Parentage unknown) R & N: J. Kovář I & REG: 2001. Inside deep red-brown, throat yellow, tips yellow-brown; outside darker red-brown, with deeper and brown-er ribs; nectaries green; pollen yellow-brown; stigma dark grey. Fls 130mm wide; petals 130 × 50mm, slightly recurved (outer more so); pedicels 65mm long, dotted with brown. Lvs alternate, 90 × 9mm. Stems 1.1m, slender, greyed green, with darker spots. Mid July.

VII (a/b) 'Newcomer'

cv. (Parentage unknown) R (pre-1987), N & REG (2002): Bischoff Tulleken Lelies B.V. Inside white (155D), with brilliant yellow/light greenish yellow (8A/B) along midveins; nectaries light green; pollen dark red-brown; stigma purple; papillae present; strongly scented. Petal margins ruffled, tips recurved. Stems light green, with darker markings.

VI (-c) Niagra Highland Henryi Group

grp (*henryi* hybrids) R (pre-1967) & N: T.R. Martin. Fls apricot orange. Stems up to 2.2m. August.

I (a/c) 'Nisku'

cv. (Un-named, I(a), white seedling × un-named seedling with yellow fls and red brushmarks) H (1988), G (1991), N (1994) & REG (2001): E.E. Fox I: Fox Lily Ranch, 2002. Inside white, with two moderate red (185B) pencil strokes (c20mm long) extending towards tip from nectary furrow on each petal, throat light yellow-green (2C); outside light yellow-green (2C) with brilliant orange-yellow (23B) midribs; large, dark red (183A) spots over basal ½ of each petal; nectaries moderate red (179A); pollen moderate reddish orange (173B); stigma moderate red (179A). Fls 130mm wide; petals 65 × 37mm, not ruffled, tips not recurved. Lvs alternate, 100 × 15mm. Stems 0.85m, pale green, with 12-16 fls. Late July. (Etymology: Nisku is a village 20 miles from Fox Lily Ranch and means "Goose" in Cree dialect.)

I (a/b) 'Nova Lux'

cv. (Parentage unknown) R (pre-1987) & N: Gebr Vletter & Den Haan. Inside pale greenish yellow (2D) and strong orange (24A), markings dark purple, throat light yellow/green; spots and papillae present; pollen light brown; stigma yellow. Petal margins ruffled, tips recurved. Stems mid green with darker markings. (Unacceptable name (Art. 17.9), but accepted in The Netherlands as part of a grant of Plant Breeders' Rights (1988). The account published under this epithet in *The International Lily Register & Checklist (1982), Twentieth Supplement*, p. 36 (2002) was in error and represents the plant registered as 'High Style'.)

I (a/-) 'Old Colony Queen'

cv. (Sister seedling to 'Jolly Green Giant') R (pre-1982) & N: Blackthorne Gardens. Inside bright yellow, tinged green, with tiny deep red spots. August to September.

I (a/-) 'Olga'

cv. (('Rubinovaja' × Hallmark Group) × (*L.* × *maculatum* × *cernuum*)) R (1983) & N: M.F. Kireeva. Inside apricot white, spotted; pollen absent. Stems 1.2m, with bulbils. July. (Unacceptable name (Art. 17.2).)

VI (-/a) 'Olive Christine'

cv. (Parentage unknown) R (pre-1968) & N: H.E. Barry. Fls deep apricot.

I (a/-) 'Ophelia'

cv. (Parentage unknown) R (pre-1992) & N: A. Weinreich. Inside pale yellow, tinged orange at base and dark spotted at base. Petals reflexed. (Unacceptable name (Art. 17.2).)

VI (-/-) Orange Glocke Group

grp (Parentage unknown) R (pre-1982) & N: V. Strasser. Fls golden yellow to orange, with red-brown midribs. Stems to 1.4m.

I (a-b/c-d) 'Orange Idyll'

cv. (Un-named seedling (no. NNC7.44) × 'Rusalka') H (1989) & G (1992): G. Steinbrück N, I & REG: 2002. Inside strong reddish orange (169A), becoming moderate red towards the base; outside similarly coloured; few, small, dark red (187C) spots at the base of each petal; nectaries light purplish pink (63D); pollen strong brown (172A) in thick anthers; stigma greyish red (178A) on a short style. Fls 160mm wide; petals 78 × 28mm, not ruffled, tips slightly recurved. Lvs alternate, 70 × 15mm. Stems 1.1m, dark green, with c8-10 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032259).)

VIII (b/-) 'Orania'

cv. (Un-named seedling (no. 93-50-108-L) × un-named seedling (no. 87-319-1)) H, G, N & I: World Breeding REG: 2001. Inside strong orange (24A), with light yellow (20B) tips and a light yellow (18A) throat; outside strong orange-yellow (24B), with strong purplish red (61B) midveins; spots almost absent; no papillae; nectaries strong yellow-green (143C); pollen dark reddish orange (175C); stigma dark purple (83A). Fls 200mm wide; petals 140 × 60mm, not ruffled. Lvs alternate, 190 × 35mm. Stems 1.25m, green, with some darker markings and c7 fls. (An orienpet lily.)

I (a/-) 'Oranje Zon'

cv. ('Tabasco' × unknown) R: The Netherlands, pre-1977. Inside vivid orange (28B) inside; outside strong orange (25A). Stems bright green, with a little darker spotting at base.

VII (-/-) 'Orchid Blush'

cv. (Selection from Lavender Lady Group) R: unknown, pre-1970. Inside white, with a lavender flush and lavender spots. (Possibly of Australian origin.)

VIII (b/b-d) 'Pacific Prize'

cv. (*henryi* × un-named trumpet) × 'Canterbury Belle') H (1997), G, N & I: A.E. Alexander REG: 2002. Inside white, with a strong apricot throat; outside white,

flushed with pink and green; unmarked; nectaries green; pollen orange; stigma creamy beige. Fls 150mm wide; petals 100 × 40mm, not ruffled, tips quite strongly recurved. Lvs alternate, 150 × 20mm. Stems 1.3m, green, flecked with dark purple and with 6-9 fls in a racemose inflorescence. Late December to mid January (New Zealand). Ryrube Group. (Standard: colour print provided by registrant (WSY 0032052). An orienpet lily.)

I (a/b) 'Papagena'

cv. (Parentage unknown) R (pre-1992) & N: G. Steinbrück. Fls creamy yellow, with many spots.

VI (-/a) 'Parthenon'

cv. (Parentage unknown) R (c1966) & N: C. Feldmaier. Inside white; outside greenish.

I (a/b) 'Peppone'

cv. (Parentage unknown) R (pre-1985) & N: Fa. P. Hoff & Zn REG: Hoffgaarde, 2002. Inside strong reddish orange (169A), with apex and margins moderate red (179A), markings dark brown, throat moderate red (181B); outside moderate red (179A/180A), with dark red/moderate red (183B/18A) towards margins; spots and papillae present; pollen dark brown; stigma orange, with a few purple-red glandular hairs. Petals slightly ruffled, tips recurved. Stems brown, with mid green spots and darker markings.

I (c/b-d) 'Perle der Karibik'

cv. (Un-named seedling (no. Fe4 ex Feldmaier) × un-named seedling (no. NNC7.44)) H (1989) & G (1992): G. Steinbrück N, I & REG: 2002. Inside yellowish white (155D), suffused palest rose; outside similarly coloured and with greenish yellow midribs; numerous, small, round to elongated, deep purplish red (61A) spots on basal ½ of each petal; nectaries grey-white; pollen deep purplish red (61A); stigma deep purplish red (61A). Fls 100mm wide; petals 60 × 30mm, not ruffled, tips strongly recurved. Lvs alternate, 90 × 15mm. Stems 0.75m, dark green, with 7-10 fls. Late June. (Standard: colour print provided by registrant (WSY 0032261).)

I (a/b) 'Pico Bello'

cv. (Parentage unknown) R (pre-1987), N & REG (2002): Fa. J. Boon. Inside pale yellow (8D) on a moderate red (181C) ground, markings dark purple; throat dark pink; spots and papillae present; pollen dark brown; stigma purple. Petals wavy-edged, tips slightly recurved. Stems light green, with darker markings.

I (a/-) Pilgrims Hope Group

grp (Complex origin from hybrids of *dauricum* var. *luteum* and 'Connecticut King') R (pre-1982) & N: Blackthorne Gardens. Inside yellow, overlain gold in the centre of the flower. May to July.

I (a/b-d) 'Pinar del Rio'

cv. ('Aelita' × 'Rubinstern') H (1989) & G (1992): G. Steinbrück N, I & REG: 2002. Inside moderate red (181B), with a narrow, longitudinal, pale orange-yellow (159C) stripe in centre of each petal; outside paler salmon pink, with strong green-yellow midribs; numerous, moderate-sized, round, greyish reddish brown (200B) spots on ¾ of each petal; nectaries white, tinged light violet at end; pollen greyish red (178A) on long, pink filaments; stigma greyish reddish orange (174B) on a 60mm long beige-coloured style. Fls 140mm wide; petals 70 × 25mm, not ruffled,

almost the whole petal distinctly curved. Lvs alternate, 110 × 18mm. Stems 1.1m, black-green, with c6-8 fls and stem bulbils. Late July. (Standard: colour print provided by registrant (WSY 0032262).)

VII (-/-) Pink Auratum Group

grp (*auratum* hybrids) R (pre-1982) & N: Blackthorne Gardens. Inside soft pink, with darker spots. Stems 1.2m. August.

VIII (c/b) 'Pink Bounty'

cv. (Un-named white Aurelian seedling) × ('Starburst Sensation' × ('White Henryi' × 'Dominique')) H (1997), G (1999), N (2001) & REG (2001): R.H. Bazett I: B & D Lilies, 2001. Inside strong to light purplish pink (63C-D), the colour progressively more intense from margins to centre of petals, throat yellow to green-yellow towards the green nectaries; outside pale yellow (158A), blending to yellowish white (158D), colour at margins similar to inside colour, midribs pale creamy pink with a hint of green; unspotted; pollen moderate orange (167B); stigma pale grey-green, dark purple at edges; strongly scented. Fls 160mm wide; petals 145 × 50mm, not ruffled, tips slightly recurved. Lvs alternate, 240 × 30mm. Strong stems to 1.9m, dark green, with up to 34 fls, including tertiaries. Early July. (Standard: colour print provided by registrant (WSY 0030916). An orienpet lily.)

VII (-/-) 'Pink Dawn'

cv. (Parentage unknown) R (pre-1981) & N: H.J. Strahm. Inside rhodonite pink with white edges and fuchsia purple spots. Fls 250mm wide; petals broad, overlapping, wavy-edged. Stems 1.2-1.9m. August. (Unacceptable name (Art. 17.2).)

VII (-/-) Pink Dream Group

grp (Parentage unknown) R (pre-1983) & N: Blackthorne Gardens. Fls soft pink. Stems 1.0m. August.

I (-/-) Pink Glory Group

grp (Selections from Futura Group) R & N: Blackthorne Gardens I: 1982. Fls pink. (Unacceptable name (Art. 26.1).)

VIII (c/a-b) 'Pink Jazz'

cv. ('Heads Up' × ('Starburst Sensation' × ('White Henryi' × 'Dominique')))) H (1997), G (1999), N (2001) & REG (2001): R.H. Bazett I: B & D Lilies. Inside light to pale purplish pink (55C/D), deeper coloured towards centre of petals, throat strong red (53D), blending to yellow near the pale green nectaries; outside light purplish pink (65B), outer petals deeper pink with a green cast over midribs, inner petals paler pink with distinct deep pink midribs; unspotted; pollen vivid orange (28B); stigma pale green; strongly scented. Fls 150mm wide; petals 150 × 50mm, margins slightly ruffled, tips slightly recurved. Lvs alternate, 200 × 20mm. Stems 1.63m, dark green, with c11 fls in a racemose inflorescence. Mid July. (Standard: colour print provided by registrant (WSY 0030919). An orienpet lily.)

VII (b/c) Pink Jewel Group

cv. (Parentage unknown) R (pre-1966) & N: R.W. Bemis I: Blackthorne Gardens. Inside rich pink, edged silver; resembling a large *speciosum*, but less reflexed and tending to face outwards; candy-like scent. Stems 1.2m. Late.

I (a/b-c) 'Pink Joy'

cv. (Parentage unknown) R: unknown, pre-1986. Inside pale rose, centre paler,

throat green at base; outside pale rose with pale green midveins; sparse, tiny, dark pink spots over 1/6 of each petal; pollen apricot. Fls 120mm wide; petals 75 × 35mm, moderately recurved at tips. Lvs alternate, 70 × 8mm. Stems 0.8m, green, tinged brownish at base. Early July.

VII (-/-) 'Pink Lace'

cv. (Selection from Atomic Group) R (pre-1988) & N: B.G. Hayler. Fls pink.

VIII (a-b/-) 'Pink Promise'

cv. (*longiflorum* seedling × oriental seedling) H (c1990) & G (c1992): Takii & Co. Ltd N (1998), I (2000) & REG (2001): De Jong Lelies Holland B.V. Inside strong purplish pink (73B) paling towards base and margins, base of midvein yellow-green changing to purple-red; outside very pale purple (73D) to white; no spots or papillae; nectaries green; pollen red-brown; stigma grey-green. Fls 180mm wide; petals 115 × 45mm, margins ruffled, tips recurved. Lvs alternate, 140 × 17mm. Stems 1.2m, green, with c5 fls (16-18cm bulb). Midseason. (A *longiflorum*/oriental hybrid.)

VII (-/-) 'Pink Spire'

cv. (*auratum* × *speciosum*) R (pre-1970), N & I: R.E. Harrison. Fls soft rose, up to 200mm wide, star-shaped. Stems 1.9-2.5m.

VII (-/b) 'Poise'

cv. (Parentage unknown) R (pre-1978) & N: Strahm Bulb Farm. Inside deep pink, with a deeper pink band, red spots and white tips; nectaries green, edged gold. Stems 0.6-1.2m. August.

VI (b-c/b-d) 'Poklad'

cv. ((Palmer seedling no. 582812 × 'Gold Eagle') × 'Mimosa Star') R & N: B. Mićulka I & REG: 2002. Inside yellow, with fine yellow spots in lines over 2/3 of each petal; outside with green midveins and brownish at base; pollen brown; stigma green. Fls 140mm wide, flat bowl-shaped; petals 85 × 35mm, slightly recurved (outer petals more so), roughly wavy and sometimes twisted; pedicels 100mm long, dark spotted. Lvs alternate, 90 × 12mm, dark green. Stems 1.4m, pale green, slightly brown spotted, with a moderately long inflorescence. Late July.

I (a/b-c) 'Pretty Girl'

cv. (Un-named seedling × un-named seedling) H (1990), G (1992), N (2002) & REG (2002): R.H. Bazett I: The Lily Nook. Inside strong purplish pink (63C), with basal part of each petal pale yellow (11C-D), throat pink; outside strong purplish red (63B), with pale green midribs; numerous small spots over basal (white) 2/3 of each petal; nectaries pink to green; pollen moderate reddish brown (166B); stigma dark purple-brown. Fls 150mm wide; petals 80 × 40mm, not ruffled, tips slightly recurved. Lvs alternate, 180 × 21mm. Stems 1.2m, mid green, with c8 fls. Late June to early July. Tetraploid. (Standard: colour reproduction (WSY 0035139) provided by registrant.)

I (-/-) Primrose Glory Group

grp (Selections from Futura Group) R, N & I (1982): Blackthorne Gardens. Fls primrose yellow.

VI (b/b) 'Quiet Times'

cv. ('Royal Flush' × ('Pamela's Joy' × 'Kiri')) R (pre-1998) & N: G. & A. Geering.

Inside white with a pale yellow throat, green nectaries and unspotted; outside white, with pale red stripes; pollen dark orange, anthers purplish; stigma pale greeny grey, style purplish at apex. Petals not ruffled, not or slightly reflexed. Stems with 6 or more fls.

VII (-/b) Raptura Group

grp (*L. × parkmannii* type hybrids) R (pre-1971) & N: Strahm Bulb Farm. Inside with watermelon red bands, white edges and profuse crimson spots. Fls 225-250mm wide. Stems 1.0-1.9m. August. (Unacceptable name (Art. 19.8).)

I (b-c/c) 'Rasma'

cv. (Parentage unknown) H, G, N (1985) & I: V.P. Orekhov. Golden yellow throughout, centre of fls with numerous dark brown spots; pollen brown. Fls 120mm wide; petals 80 × 35mm, not ruffled, strongly recurved. Lvs alternate. Stems 1.1m, brown.

I (a/b) 'Rebel'

cv. ('Target' × 'Windahloo') R & N: B. Mičulka I & REG: 2002. Inside orange-red on a bronze ground, with a narrow yellow rim, throat yellowish; outside bronze orange, with pale greenish yellow midveins; large, diffused, dark mahogany brushmarks and a few, small, dark red-brown spots over 1/5 of each petal; nectaries white granulated; a few small papillae in the throat; pollen brown ochre; stigma dark crimson. Fls 120mm wide; petals 75 × 38mm, outer petals slightly reflexed; pedicels 20mm long, thick, brown spotted, hairy. Lvs alternate, 65 × 11mm, pale green. Stems 0.85m, thick, slightly hairy, slightly brown spotted, with a short inflorescence. Late June.

VIII (a-b/-) 'Red Alert'

cv. (Un-named seedling × un-named seedling) H: Testcentrum voor Siergewassen G (1996), N (2001), I (2001) & REG (2002): Marklily C.V.. Inside dark red (183B/187B); outside deep red (185A); nectaries purple-red; spots purple-red; papillae present; pollen red; stigma red. Fls 150mm wide; petals 100 × 50mm, margins slightly ruffled, tips slightly recurved. Lvs alternate, 150 × 20mm. Stems 0.9m, green and brown, with 3-6 fls. Early to midseason. (*A longiflorum*/asiatic hybrid.)

I (c/-) 'Red Beauty'

cv. ('Radostnaja' × un-named seedling) R (c1979) & N: De Jong Lelies Ltd. Inside vivid reddish orange (34A) and vivid reddish orange/vivid red (42A/45A); outside moderate red (179A); a few black spots; pollen light brown. Petals 80 × 30mm, not ruffled, slightly recurved. Lvs alternate, 110 × 25mm. Stems 1.2m, brown violet, with 8-12 fls. July.

VII (-/-) 'Red Devil'

cv. (Parentage unknown) R (pre-1997) & N: F. Jury. Fls clear bright red throughout, including nectaries.

I (a/-) Red Flare Group

grp (Parentage unknown) R: unknown, pre-1982. Inside red, with dark spots on basal × of each petal. Petal margins ruffled. (Unacceptable name (Art. 19.8).)

I (-/-) Red Glory Group

grp (Selections from Futura Hybrids) R, N & I (1982): Blackthorne Gardens. Fls red.

I (a/-) 'Redwood'

cv. (Parentage unknown) R & N: E.H. Doerr I: Borbeleta Gardens, 1982. Inside deep red, unspotted; nectar furrows pale; stigma pale. Petals not ruffled, slightly reflexed at tips, rather narrow. Lvs deep green. Stems 1.0m, with up to 30 fls. (Unacceptable name (Art. 17.2).)

I (a/-) 'Reimei'

cv. ('Enchantment' × *L. × maculatum*) R (c1968) & N: S. Fujishima. Fls red.

I (b/-) 'Renaissance'

cv. (Selection from *lancifolium* hybrids) R (early 1970s) & N: Laan Lelie B.V. Fls brick red. Stems 1.2m. Early July.

I (-/-) 'Renoir'

cv. (Parentage unknown) R (pre-1989) & N: A. Weinreich. Fls salmon orange.

VII (a/-) 'Rioe-Kioe'

cv. (Parentage unknown) R (pre-1994), N & REG (2002): Gebr. Vletter & Den Haan. Inside light purplish pink (65B), base of midvein strong purplish red (63A); outside white (155D), with a pink suffusion; nectaries green changing into white; spots and papillae present; pollen orange-brown; stigma purple. Petal margins ruffled, tips recurved. Stems with some darker markings. Midseason.

I (a/-) 'Rosa Hoffnung'

cv. (Parentage unknown) R & N: G. Steinbrück, pre-1992. Fls star-shaped, cream and rose.

I (c/-) Rosette Group

grp (Hybrids between *davidii* and *cernuum*) R (pre-1985) & N: R.W. Bemis. Very varied pastel colours, some bicoloured; very long pedicels. Stems 1.0-1.2m. (Unacceptable name (Art. 19.8).)

I (a/-) 'Roswitha'

cv. (Parentage unknown) R (pre-1986) & N: A. Rech. Inside white. Stems 1.0m, dark green, with c10 fls. Late June. (Published in German Lily Group sales list, 1986.)

VII (-/b) Rosy Treasures Group

grp (Selections from *auratum* × *speciosum* hybrids) R (pre-1982) & N: Strahm Bulb Farm. Fls in various shades of pink. July to August.

VIII (b/b) 'Rote Zunge'

cv. (*regale* seedling × 'Blood Tiger') H (1992), G (1996), N (1998) & REG (2002): P. Müller. Inside deep red (53A) with 53A tips, outside of inner petals dark red (187B), outside of outer petals moderate purplish red (64A); a few thin markings of dark red (187A); nectaries vivid reddish orange (34A); pollen dark red-brown (but anthers often absent); stigma dark purplish pink (186C). Fls 130mm wide; petals 90 × 22-33mm, not ruffled, tips slightly recurved. Lvs alternate, 110 × 7mm. Stems 1.0m, brown with green spots and c12 fls. Midseason. (Standard: colour illustration provided by registrant (WSY 0035253) An asiatic/trumpet hybrid.)

I (a/b) 'Rotterdam'

cv. (Parentage unknown) R (pre-1986), N & REG (2002): Vletter & Den Haan.

Inside of inner petals dark yellowish pink (181D) at apex, moderate red/moderate purplish red (184B/C) at base, inside of outer petals dark pink (182C/D) at apex, moderate red (184B/185B) at base; outside of inner petals moderate red (184B) with moderate red (181C) to moderate purplish pink (54D) at apex; outside of outer petals moderate purplish pink (54D); spots and papillae present; stigma red-purple; pollen orange. Petals not ruffled, tips recurved. Stems brown, with mid green ribs and spots and darker markings.

VI (-/-) 'Royal Flush'

cv. (Derived from Pink Ice Group) R (pre-1987) & N: G. & A. Geering. Fls white to cream; not pendent, not trumpet-shaped.

VIII (a/-) 'Royal Giant'

cv. (Un-named seedling (no. A 412-87) × un-named seedling (no. AH 627-T-444)) H (1991) & G (1993): World Bulbs BVBA N & I: 1999 REG: 2002. Inside brilliant greenish yellow (6C), paler towards tips and edges, base yellow-green; outside light greenish yellow (6D); nectaries green; no spots or papillae; stigma yellow; pollen brown. Stems 1.3m, green. Early to midseason. (A *longiflorum*/asiatic hybrid.)

VIII (a-b/-) 'Royal Humboldt'

cv. (Un-named seedling (no. A-226-83) × un-named seedling (no. 8759-12)) R, N (1998) & I: J.W.A. van der Wereld & Zn B.V. REG: World Flower B.V., 2002. Inside deep purplish pink (54B), paler towards apex, red at base; outside strong pink (54C); nectaries green; spots and papillae present; pollen brown; stigma purple. Petal margins ruffled, tips recurved. Stems 1.3m, green. Early. (A *longiflorum*/asiatic hybrid.)

VIII (a-b/-) 'Royal Inspiration'

cv. (Parentage unknown) R, N & REG (2002): J.W.A. van der Wereld. Inside light orange-yellow (23C), towards the apex; outside light orange-yellow (23C); spots and papillae present; pollen orange-brown; stigma purple. Petal margins ruffled, tips recurved. Stems green, with darker markings. Midseason to late. (A *longiflorum*/asiatic hybrid.)

VIII (a/-) 'Royal Melody'

cv. (Un-named seedling (no. 5804-1) × un-named seedling (no. A-204-7)) R & N: J.W.A. van der Wereld REG: World Flower B.V., 2002. Inside strong orange (26A); outside brilliant orange (25C); nectaries green; no spots or papillae; pollen brown; stigma light brownish orange. Stems green. Very early to early. (A *longiflorum*/asiatic hybrid.)

VIII (a-b/-) 'Royal Rain'

cv. ('Bora' × un-named seedling (no. AL-11-85)) R (pre-1999), N & I: J.W.A. van der Wereld & Zn B.V. REG: World Flower B.V., 2002. Inside moderate yellowish pink (39C), yellow at base; outside light pink (50D); nectaries yellow; spots and papillae present; pollen red; stigma orange. Petal margins ruffled, tips recurved. Stems 0.95m, green, with darker markings. Early. (A *longiflorum*/asiatic hybrid.)

I (a/-) 'Royal Shadow'

cv. ('Monte Negro' × un-named seedling (no. 5866-1)) R, N (1998) & I: J.W.A. van der Wereld B.V. REG: 2002. Inside strong red (46A)/deep red (185A); outside deep red (53A); nectaries green; spots and papillae present; pollen dark brown; stigma red. Petal margins ruffled, tips recurved. Stems 1.25m, green, with darker markings. Early to midseason.

I (b/c) 'Ruby Glow'

cv. (Un-named red seedling × un-named red seedling) **H** (1989) & **G** (1991); A.J. Burnett **N** (1997) & **REG** (2001); E.E. Fox I: Fox Lily Ranch, 2002. Inside vivid red (46B) with a vivid orange-yellow (23A) throat; outside vivid red (46C); fine, dark red (187A) spots on basal $\frac{1}{2}$ of each petal; nectaries strong red (46A); pollen vivid orange (28A); stigma greyish red (176A). Fls 150mm wide; petals 75 × 35mm, not ruffled, tips not recurved. Lvs alternate, 130 × 17mm. Stems 0.85m, mid green, with 18-20 fls. Late July. (Etymology: name descriptive of colour when backlit.)

VI (c/d) 'Rügenflirt'

cv. (*henryi* × mixed pollen of 'White Henryi', 'Mimosa Star' and 'Louise') **H** (1990) & **G** (1994); G. Steinbrück **N**, **I** & **REG**: 2002. Inside white (155A), with a light orange-yellow (24C) throat; outside with bases pale green otherwise greenish white (157D); many, moderate-sized spots and streaks of moderate purplish red (59C) and conspicuous, large, light orange-yellow (24C) papillae; nectaries moderate yellowish green (138A); pollen moderate reddish brown (175A), on very long, moderate yellowish green (138A) filaments; stigma moderate yellow-green (138B) on a long, pale green style. Fls 97mm wide; petals 75 × 25mm, slightly ruffled, strongly reflexed. Lvs alternate, 200 × 25mm. Stems 1.2m, dark green, with c15 fls. Mid to late July. (Standard: colour print provided by registrant (WSY 0032264).)

I (a/b) 'Sachsenflamme'

cv. (Un-named, yellow-orange seedling (no. 308/1) × 'Cabaret') **H** (1989), **G** (1991), **N** & **I** (2003); G. Thürmer **REG**: 2001. Inside deep yellowish pink (39B) in apical half with a very narrow pale margin, giving way to a broad zone of brilliant yellow (12B) and at the base of each petal dark reddish; outside deep yellowish pink (39B); a few deep purplish red (59B) spots inside, above the dark basal area; nectaries greenish; pollen strong reddish orange (169A). Fls 150mm wide; petals 80 × 40mm, not ruffled, tips not or slightly recurved. Lvs alternate, 110 × 12mm. Stems 1.0m, pale green, with stem bulbils and with c10 fls. Mid to late July. (Standard: colour print provided by registrant (WSY 0030948).)

I (a/b) 'Salaška'

cv. (Parentage unknown) **R** & **N**: B. Mičulka **I** & **REG**: 2002. Inside deep yellow, throat greenish; outside with greenish midveins; numerous, small, dark brown spots over 2/5 of each petal; nectaries green, white granulated; pollen pale brown; stigma greenish. Fls 140mm wide; petals 80 × 41mm, not recurved; pedicels 50mm long, thick, green, with secondary buds. Lvs alternate, 90 × 12mm, dark green. Stems 0.9m, thick, green, with a moderately long inflorescence. Early June.

VII (a/c) 'San Pedro'

cv. (Un-named seedling × un-named seedling) **H**: Peacock/Imanse Lelieveredeling B.V. **I**: Marklily C.V., 2000 **G**, **N** (1999) & **REG** (2002): Testcentrum voor Siergewassen. Inside deep to strong red (53A/B), changing to moderate purplish red (59C) on a white ground near edges, edged white (narrowly so at base), throat white; outside moderate purplish red (59C), paler towards apex, narrowly edged white, midvein light yellow-green (150D) with a little strong yellow-green (144B) at base; nectaries strong yellow-green (144B) surrounded by a little light greenish yellow (8B); dark red (187A+B) spots on up to $\frac{1}{2}$ of each petal and dark red (187A+B) papillae in the throat; stigma deep purplish red (59B) on a beige ground; pollen strong brown (172A). Fls 300mm wide; petals 155-160 × 45-75mm, strongly ruffled, tips strongly recurved. Lvs alternate, 150-185 × 25-50mm. Stems 1.0m, light green, with a little darker marking.

I (a/c) 'Santa Clara'

cv. ('Dawn Star' × 'Summer Night') H (1989) & G (1992): G. Steinbrück N, I & REG: 2002. Inside moderate orange-yellow (165C); outside similarly coloured; many, moderate-sized, strong red (46A) spots and streaks on ¾ of each petal; nectaries pale yellow-green (157B); pollen greyish red (184A), filaments moderate yellow (162A); stigma deep red (185A). Fls 150mm wide; petals 82 × 37mm, not ruffled, tips slightly recurved. Lvs alternate, 75 × 15mm. Stems 0.75m, mid green, with 6-8 fls. Mid July. (Etymology: Santa Clara is a town in Cuba.)

VII (a-b/-) 'Sapporo'

cv. (Parentage unknown) H (1994), G, N (2001) & I (2002): Sande B.V. REG: 2002. Inside and outside white (155D); nectaries green; spots absent, papillae present; pollen red-brown; stigma purple. Petal margins strongly ruffled, petals recurved. Stems 1.0m. Midseason to late.

I (a/b) 'Sardegna'

cv. (Parentage unknown) R (pre-1988), N & REG (2002): Vletter & Den Haan. Inside strong pink (50C), shading to deep pink (50B) at base and margins, central patch of pale yellow (11C/D) on each petal, markings purple-red, throat pink; spots and papillae present; pollen orange-brown; stigma light brown. Petals ruffled, tips recurved. Stems light green. Early.

I? (-/-) 'Satanas'

cv. (Parentage unknown) R (pre-1972) & N: C. Wittneben. Fls deep shining red. Very late.

VII (-/-) 'Satan's Breath'

cv. (Parentage unknown) R (pre-1998) & N: F. Jury. Fls pale creamy white, with a very deep mahogany red centre.

I? (-/-) 'Saturnus'

cv. (Selection from Egmont Group) R (pre-1970) & N: R.E. Harrison. Fls orange flame-coloured. Fls 120mm wide. Stems 0.6m.

I (a/b-d) 'Schierke'

cv. ('Cynthia' × ('Alpenglow' × orange brushmark seedling)) H (1989) & G (1992): G. Steinbrück N & I: 2000 REG: 2002. Inside strong yellowish pink (31C); outside similarly coloured; a few, dark red (183A) spots at base of each petal; nectaries moderate olive green (147A); pollen dark red (183A); stigma yellowish grey (156B). Fls 150mm wide; petals 88 × 38mm, not ruffled, tips slightly to strongly recurved. Lvs alternate, 90 × 15mm. Stems 0.8m, dark green with black markings and 8-10 fls. Late June. (Standard: colour print provided by registrant (WSY 0032265). Etymology: Schierke is a town in the Harz region of Germany.)

I (b/-) 'Schwarzer Prinz'

cv. (Parentage unknown) R (pre-1997) & N: Dierssen. Fls black-red, with thick petals. Stems 0.8m. Mid July. (Published in ELG selling list, 1997.)

VII (a/b) 'Seagull'

cv. (Parentage unknown) R (pre-1987), N & REG (2002): Bischoff/Tulleken Lelies B.V. Inside white (155D), with a white/light green throat; papillae present; pollen dark brown; stigma dark purple. Petals ruffled, tips recurved. Stems light green, with darker markings.

VIII (a-b/-) 'Sea Treasure'

cv. (*longiflorum* 'Lorina' × 'Muscadet') H (1993), G (1995), N (2000), I (2002) & REG (2002): De Jong Beheer B.V. Inside white (155C), base of midvein yellow-green shading apically into vivid purplish red (57C); outside white (155B); nectaries green; no spots or papillae; pollen red-brown; stigma grey, suffused purple. Petals 164mm long, slightly ruffled. Lvs alternate, 230 × 30mm, veins more impressed than in 'Wild Treasure'. Stems 1.1m, green, with c5 fls. Late. (*longiflorum*/oriental hybrid.)

I (a-b/b) 'Sevilla'

cv. (Parentage unknown) R (pre-1988), N & REG (2002): Gebr Mak. Inside light orange-yellow (22C), with a strong orange-yellow (22A) glow along veins, throat orange; outside orange, with paler midveins; spots and papillae absent; pollen orange-brown; stigma orange. Petals ruffled, tips recurved. Stems light green, with some darker marking at the base. Very early to early.

VII (-/d) 'Shogun'

cv. (Parentage unknown) R: T. Sakata & Co. pre-1983 REG: Van Til-Hartman B.V., 2002. Inside strong purplish red (60C-D), with deep red (60A) along midveins and white (155D) along margins and at apex; markings red-purple; throat white (155D); outside moderate purplish red (186B), margins and apex white (155D), outer petals also with some moderate purplish red (59C); spots and papillae present; pollen orange; stigma purple. Petals recurved, margins ruffled. Stems light green, with darker markings and at base as well as at nodes a greyish waxy layer.

I (a/b) 'Silver Bunny'

cv. (Parentage unknown) R (pre-1987), N & REG (2002): Gebr. Laan & Zn. Inside creamy white (159D), apically white (155D) with a creamy white (159D) suffusion; throat white; markings red-purple; spots and papillae present; pollen dark orange-brown; stigma dark pink. Petals slightly ruffled, tips not recurved. Stems with darker markings. Very early.

VII (-/-) Silver Dream Group

grp (*rubellum* hybrids) R (pre-1982) & N: R.W. Bemis I: Blackthorne Gardens. Inside white, spotted crimson or rose; nectaries green. Stems 1.2-1.5m. July to August.

VI (-/c) 'Sirius'

cv. (Un-named seedling × 'Solntse Rossii') R (c1984) & N: V.A. Volkov. Inside cream white, with a bright orange star in throat; pollen brown. Stems 1.0m. August. (Unacceptable name (Art. 17.2).)

VIII (b-c/a) 'Skylight'

cv. (((McLaren trumpet × 'Canterbury Belle') × deep pink trumpet) × sibling) H (1996) & G (1999): A.E. Alexander N, I & REG: 2002. Inside golden apricot; outside gold with broad pink bands; unmarked; nectaries green; pollen brown; stigma purple/black. Fls 130mm wide; petals 120 × 40mm, not ruffled, tips slightly recurved. Lvs alternate, 120 × 10mm. Stems 1.2m, light green, flecked purple, with c9 fls in a racemose inflorescence. January (New Zealand). Ryrirube Group. (Standard: colour print provided by registrant (WSY 0032051). An orienpet hybrid.)

VIII (a/b) 'Sonatina'

cv. (Un-named seedling (no. BG 95-71) × un-named seedling (no. OR 93-20)) H,

G, N (1999) & I: Vletter & Den Haan Beheer B.V. Inside deep red (185A) on a moderate red (185B) ground, with a deep orange-yellow (163A) suffusion above nectary for c $\frac{1}{2}$ of petal, throat brilliant yellow-green (150C); outside with moderate yellow (160A) midveins, deep pink (180D) alongside the midveins and becoming moderate red (183D) towards edges; nectaries moderate yellowish pink (179D); dark red (187B) spots but no papillae; pollen strong orange (24A); stigma brilliant orange (25C). Fls 170mm wide; petals 130 × 50-60mm, not ruffled, tips recurved. Lvs alternate, 150-160 × 20-25mm. Stems 1.0-1.3m, green, with darker markings and c4 fls. (*A longiflorum*/asiatic hybrid.)

I (-/-) Sonnenstrahlen Group

grp (Parentage unknown) R (pre-1970) & N: C. Wittneben. Inside orange-yellow, with or without spots.

I (b/b-d) 'Sonnentempel'

cv. ('Yukatan' × un-named seedling (no. Fe4 ex Feldmaier)) H (1989) & G (1992): G. Steinbrück N & I: 2001 REG: 2002. Inside pale orange-yellow (24D); outside pale orange-yellow (24D), with soft pink midribs; many, large, mostly round, deep purplish red (71A) spots over $\frac{2}{3}$ of each petal; nectaries pale orange-yellow (159B), hoary; pollen moderate reddish orange (171A), on soft pale yellow filaments; style thick, soft yellow-green. Fls 160mm wide; petals 81 × 40mm, not ruffled, tips slightly recurved. Lvs alternate, 90 × 16mm. Stems 1.2m, violet black (as are the pedicels), with c15 fls in a pyramidal inflorescence. Early July. (Standard: colour print provided by registrant (WSY 0032267).)

I (a/b) 'Sortini'

cv. (Parentage unknown) H (1990), G (1993), N (1996) & I (1996): J. Vasarietis REG: 2001. Inside with a vivid yellow (12A) ground throughout, above the nectary about $\frac{1}{2}$ of each petal is spotted with red-purple and marked with brushmark lines; outside vivid yellow (12A) at edges, remainder brilliant yellow (12B); nectaries white; pollen orange; stigma pale greyish. Fls 140mm wide; petals 90 × 43mm, not ruffled, tips slightly recurved. Lvs alternate, 140 × 20mm. Stems 0.95m, brownish green, with c8 fls. Early to mid July. (Standard: colour print provided by registrant (WSY 0030815). Etymology: Sortini is locality where the raiser's wife was born.)

I (a/b-d) 'Sousse'

cv. ('Heidi III' × 'Berliner Morgen') H (1992) & G (1995): G. Steinbrück N, I & REG: 2002. Inside strong orange (170A), tips moderate orange (171D), throat brilliant orange-yellow (23B); outside moderate orange; scattered, small, dark reddish orange (175B) spots on basal $\frac{1}{3}$ of each petal; nectaries yellowish white (156D); pollen strong reddish orange (169A); stigma moderate reddish brown (175A). Fls 150mm wide; petals 80 × 35mm, not ruffled, tips recurved. Lvs alternate, 90 × 15mm. Stems 0.75m, dark green, with 5-8 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032268). Etymology: Sousse is a town in Cuba.)

VIII (b/c) 'Southern Song'

cv. ((*henryi* × un-named trumpet) × 'Canterbury Belle') H (1989) & G (1992): A.E. Alexander N, I & REG: 2002. Pale apricot throughout, unmarked; nectaries green; pollen brown; stigma frosty green. Fls 160mm wide; petals 100 × 40mm, not ruffled, tips slightly recurved. Lvs alternate, 110 × 15mm. Stems 1.2m, mid green, flecked black, with c6 fls in a racemose inflorescence. January (New Zealand). Ryrirube Group. (Standard: colour print provided by registrant (WSY 0032054). An orienpet lily.)

IX (c/d) *speciosum* ‘White Orchid’

cv. (Selections from *speciosum* var. *album*) R (pre-1982) & N: Blackthorne Gardens. Fls snow white. Stems 1.0m. September to November. (This cultivar consists of more than one clone.)

I (a-b/c) ‘Spreecharme’

cv. (‘Connecticut Red’ × mixed pollen of ‘Vater Zille’, ‘Fanal’ and ‘Brushstroke’) H (1992) & G (1995): G. Steinbrück N, I & REG: 2002. Inside moderate red (47A) throughout; moderate-sized, round spots and a small brushmark of dark red (187A) on basal ¼ of each petal; nectaries white (155A), fading at apex and becoming pink with a green centre; pollen strong reddish orange (31A) on thin filaments; stigma deep red (185A) on a short style. Fls 150mm wide; petals 82 × 37mm, not ruffled, not recurved, inner rather broadly ovate. Lvs alternate, 80 × 15mm. Stems 0.9m, mid green, with 7-9 fls. Late June. (Standard: colour print provided by registrant (WSY 0032269).)

VII (-/-) Sprinkles Group

grp (Parentage unknown) R (pre-1982) & N: Strahm Lily Farm. Fls white, with pink or lavender spots, 150-225mm wide. Stems 1.2-1.9m. August.

I (a/-) Squanto Group

grp (Parentage unknown) R (pre-1982) & N: Blackthorne Gardens. Fls melon to soft apricot. Stems 0.8m. May to July.

VII (-/b) ‘Starfire’

cv. (Selection from Fire Dancer Group) R (pre-1983) & N: Strahm Lily Farm. Fls crimson red, with darker spots and white margins. Stems 1.0-1.9m. August to September. (Unacceptable name (Art. 17.2).)

VII (b-c/b) ‘Starmaker’

cv. (Parentage unknown) R (pre-1985), N & REG (2002): Sun Valley Bulb Farms. Inside with a central moderate purplish red (184C) area on each petal, midveins moderate/strong red (47A/B), along margins moderate purplish red/deep pink (185C/D), margins white (155A); markings red-purple; throat white (155C); outside white (155D), with a purple-pink suffusion and light green midveins; nectaries green; spots and papillae present; stigma purple; pollen orange-brown. Petal tips recurved, margins ruffled. Stems light green/greyish, with some darker markings at base.

VIII (b/a) ‘Status Hi’

cv. ((‘White Henryi’ × *auratum*) × ((Golden Splendor Group × a 4 way cross) × unnamed pink trumpet)) H (1995) & G (1999): A.E. Alexander N, I & REG: 2002. Inside pinkish white with a conspicuous gold throat; outside with pink and white bands; unmarked; nectaries green; pollen orange; stigma black. Fls 140mm wide; petals 130 × 50mm, not ruffled, tips becoming quite strongly recurved. Lvs alternate, 110 × 15mm. Stems 2.0m, mid green, flecked black, with c12 fls in a racemose inflorescence. January (New Zealand). (Standard: colour print provided by registrant (WSY 0032053). An orientpet lily.)

I (c/d) ‘Stelinka’

cv. (‘Tetra Nutmegger’ × ‘Apricot Supreme’) R & N: B. Mićulka I & REG: 2002. Inside lemon yellow; outside pale pastel yellow with yellow edges and more-or-less brownish midveins; very few, dark brown crimson spots; nectaries yellow granulated; pollen and stigma brown. Fls 130mm wide; petals strongly recurved; pedicels

70mm long, thick, brownish. Lvs alternate, 90 × 14mm, mid green. Stems 0.75m, very thick, slightly spotted brown, with a short inflorescence. Tetraploid. Early July.

VII (-/-) 'Summer Charm'

cv. (Parentage unknown) R: F. Jury, pre-1997. Fls very large, white, with central bands of deep vermilion.

VII (-/d) 'Supernova'

cv. (Parentage unknown) R (c1979) & N: Oregon Bulb Farms. Fls deep red. Stems 1.5-1.9m. August. (Has been listed as a *speciosum* selection, but OBF give it as above.)

I (b/c) 'Sutherland Flamethrower'

cv. ('Merryspring' × 'Sutherland Peach Dream') H (1996), G (1998) & N (1999): D.J. Hay REG: 2001. Inside gold and red bicolor (essentially pink at apex and along margins, gold in basal part of each petal), throat red; outside pink; a red flash runs down centre of each petal from nectary to mid petal, otherwise unspotted; nectaries red. Fls 111mm wide; petals 73 × 34mm, not ruffled, tips slightly recurved. Lvs alternate, 42 × 7mm. Stems 43cm, with c6 fls. Midseason. (Standard: colour print provided by registrant (WSY 0031593).)

I (b/b-c) 'Sutherland Peach Melody'

cv. ('Unique' × 'Merryspring') H (1993), G (1996) & N (1998): D.J. Hay REG: 2001. Inside with tips pink and centre of petals apricot, throat dark pink; outside pink; heavily spotted with wine red on lower half of each petal. Fls 155mm wide; petals 73 × 39mm, margins very slightly ruffled, tips slightly recurved. Lvs alternate, 55 × 9mm. Stems 65cm, green, mottled with brown and with 7-9 fls. Midseason. (Standard: colour print provided by registrant (WSY 0031599).)

II (-/-) Swinging Safari Group

grp (*martagon* var. *album* & var. *cattaniae* × *medeoloides*) F2) R (pre-1981) & N: F. Ewald. Fls ochre yellow, with brown spots. Lvs and stems dark green. Stems 1.6-1.8m, with up to 15 fls. July. (Later into growth than *hansonii* hybrids and therefore less subject to frost damage.)

I (a/b-c) 'Tanz der Feen'

cv. (Un-named seedling (no. LV15, ex Matthe) × 'Luxor') H (1991), G (1994), N (1999) & I (2002): G. Steinbrück REG: 2002. Inside white (155A), with a soft suffusion of pale yellow (20D) in centre towards base, tips pure white; outside white (155A); numerous, moderate-sized, strong purplish red (54A) spots on basal ½ of each petal; nectaries white (155A), hoary; pollen moderate reddish brown (174A); stigma pale yellow (20D). Fls 150mm wide; petals 88 × 42mm, not ruffled, tips not recurved. Lvs alternate, 90 × 15mm. Stems 0.8m, mid green, with c11 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032270).)

VII (b-c/b) 'Tessa'

cv. (Parentage unknown) R (pre-1988), N & REG (2002): M.A. van de Berg. Inside of inner petals white (155D), with a pink suffusion at apex and midveins light yellow at base; inside of outer petals white (155D) at base and margins, apex light purplish pink (62C) on a white ground and with light yellow midveins; throat white/green/light yellow; spots and papillae present; pollen dark brown; stigma purple. Petal margins ruffled, tips recurved. Stems light green, with a grey waxy layer and darker markings. Late.

IX (b/d) 'The Moor'

See *martagon* var. *cattaniae* 'The Moor'.

VII (a-b/-) 'Tosca'

cv. (Un-named seedling × un-named seedling) H: Imanse Lelieveredeling B.V. 1991 N (1995), I (1995) & REG (2002): Peacock B.V. Inside white (155D), base of mid-vein pale yellow (8D) changing into light purplish pink (73C), throat rose; outside white (155D); nectaries green changing into yellow-green; spots and papillae present; pollen red-brown; stigma purple. Fls 200mm wide; petals 110 × 40mm, margins ruffled, tips recurved. Lvs alternate, 100 × 30mm. Stems 85cm, green, with 2-6 fls. Midseason.

VII (a-b/-) 'Tourmalet'

cv. (Parentage unknown) H (1985), G (1988), N (1991) & REG (2002): Vletter & Den Haan Beheer B.V. Inside white (155A); outside white (155D); nectaries green; spots and papillae absent; stigma dark purple; pollen light brown. Stems 1.4m. Early to midseason.

I (b/b-d) 'Trinidad'

cv. ('Rubinstern' × 'Darja') H (1989) & G (1992): G. Steinbrück N, I & REG: 2002. Inside brilliant yellow (7A); outside brilliant yellow (7A), with soft pink midribs; many, large, dark red (187A) spots on more than $\frac{3}{4}$ of each petal and pale papillae all across base of each petal; nectaries white (155B), hoary; pollen brownish orange (171B), on long, slightly reflexed filaments; stigma light yellowish pink (36A) on a long, yellow style. Fls 140mm wide; petals 80 × 41mm, not ruffled, tips strongly reflexed. Lvs alternate, 90 × 17mm. Stems 1.1m, red-brown (like the pedicels), with 7-10 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032271). Etymology: named for the town of Trinidad in Cuba.)

I (a-b/-) 'Trojan'

cv. ('Enchantment' × 'Connecticut King') I: Blackthorne Gardens, 1983. Inside amber pink, with oxblood red brushmarks. Stems 0.75m. May to June. (Unacceptable name (Art. 17.2).)

I (a-b/b) 'Twincolour'

cv. (Parentage unknown) R (pre-1988), N & REG (2002): Vletter & Den Haan Beheer B.V. Inside strong reddish orange (42B) with a central area of vivid yellow (12A) in centre of each petal and dark purple-red markings; throat red-brown; outside strong reddish orange (42B), darker towards margins; spots and papillae present; pollen brown; stigma dark purple. Petals with ruffled margins and slightly recurved tips. Stems light green, with darker markings at base. Very early to early.

VI (-/a) 'Uncle Mark'

cv. (*regale* × *leucanthum* var. *centifolium*) R: Lord Limerick, pre-1984 N & I: J. & E. Parker-Jervis. Fls white inside, with a rose madder exterior; scented. Stems 1.5m, with 6-8 fls.

I (a/b) 'University of Saskatchewan'

cv. ('Geneve' × 'Merryspring') H (1997), G (2000), N (2002) & REG (2002): D.J. Hay. Inside white with a gold blush in middle of each petal; outside white, tinged pink; halo of dark spots surrounding the throat; nectaries green; pollen orange; stigma green. Fls 134mm wide; petals 68 × 37mm, not ruffled, tips very slightly recurved. Lvs alternate, 38 × 10mm, glossy. Stems 45cm, green, with c7 fls.

Midseason. (Standard: colour print provided by registrant (WSY 0035138).
Etymology: named to celebrate the 100th anniversary of the University of Saskatchewan.)

I (a-b/b) 'Ursula'

cv. (Parentage unknown) R (1991), N & REG (2002): Douwe Wierstra B.V. Inside of inner petals vivid orange-yellow (21A) with some dark red (183A), inside of outer petals vivid yellow (15A/17B) with some dark red (183A); dark red-brown markings; throat vivid yellow (12A); outside of inner petals vivid yellow (17B) sometimes with brownish orange (165B) at base, outside of outer petals vivid yellow (15A); spots and papillae present; nectaries yellow-green; pollen orange-brown; stigma purple-red. Petal margins ruffled, tips recurved. Stems brown, with green spots and darker markings. Midseason.

I (a/-) 'Vater Zille'

cv. (Asiatic seedling × mixed pollen of a brushmark seedling and 'Brushstroke') H (1989), G & N: G. Steinbrück. Inside yellow, with a large purple-violet brushmark and striking green nectaries. Fls star-shaped.

I (a/b) 'Vendija'

cv. (Parentage unknown) H (1990), G (1993), N (1996) & I (1996): J. Vasarietis REG: 2001. Inside creamy white at first, fading to white; throat and outside white; c10 tiny, dark spots scattered around each nectary; nectaries white on a greenish ground; pollen brown; stigma greyish brown. Fls 130mm wide, slightly scented especially in evening; petals 70 × 38mm, not ruffled, slightly recurved. Lvs alternate, 80 × 12mm. Stems 0.8m, light green, with c12 fls. Mid July. (First published in *Lilium Balticum Lily Register*, 1996. Standard: colour print provided by registrant (WSY 0030814). Etymology: Vendija is the ancient name of a region in Latvia.)

VII (a-b/-) 'Venere'

cv. (Parentage unknown) R (pre-1993), N, I & REG (2002): Gebr. Vletter & Den Haan Beheer B.V. Inside white (155D), with deep pink (50B) midveins; outside white (155D); nectaries green shading to yellow; spots and papillae present; pollen orange-brown; stigma dark purple. Petals margins ruffled, tips recurved. Stems 1.1m, with dark markings. Very early to early.

VI (-/a) 'Vera'

cv. (Parentage unknown) R (c1980) & N: N.A. Miroshniczenko. Inside white; outside brownish. Stems 1.8m. Early. (Unacceptable name (Art. 17.2).)

I (a/b) 'Verigold'

cv. (Un-named seedling × un-named seedling) H (1990), G (1992), N (2001) & REG (2001): R.H. Bazett I: The Lily Nook, 2003. Inside vivid yellow (17B), throat strong orange-yellow (17A); outside vivid yellow (14B) shading to vivid yellow (14A) at margins; numerous, small, elongated, dark purple spots on basal 1/2 of each petal; nectaries green-yellow; pollen strong brown (172A); stigma red-purple on a green base. Fls 160mm wide; petals 95 × 57mm, not ruffled, tips not recurved. Lvs alternate, 150 × 20mm. Stems 1.1m, dark green, with c15 fls, including secondaries. Late June. Lvs and petals of heavy texture. (Standard: colour print provided by registrant (WSY 0030918).)

I (a-b/b) 'Veronie'

cv. (Un-named seedling × un-named seedling) H: Imanse Lelieveredeling B.V. &

Peacock B.V., 1995 G (1998), N (2001), I & REG (2002): Marklily C.V. Inside vivid reddish orange (32A), throat strong yellowish pink (32D); outside strong reddish orange (31B), edged vivid reddish orange (32A), midveins on outer petals strong yellow-green (143A); a few dark red (187A) spots; papillae present; pollen strong reddish orange (169A); stigma vivid reddish orange (30A). Fls 160mm wide; petals 90 × 50mm, margins not ruffled, tips recurved. Lvs alternate, 190 × 15mm. Stems 1.2m, dark, with c6 fls, from hairy buds. Early.

I (a/c) 'Vikomt'

cv. ('Grun-weiss' × ('Windahloo' × 'Endeavor')) R & N: B. Mićulka I & REG: 2002. Inside pale yellow on opening, fading to yellowish cream, throat paler and greenish; small, inconspicuous, mahogany brushmarks and moderately numerous, moderately large, dark pinkish brown spots on ⅓ of each petal. Fls 150mm wide; petals 90 × 45mm, hardly recurved; pedicels 80mm long, thick, green, hairy. Lvs alternate, 120 × 15mm, dark yellowish green. Stems 0.95m, green, slightly hairy, brown at base and with a short inflorescence. Early June.

I (a/b) 'Vivija'

cv. (Parentage unknown) H (1990), G (1993), N (1996) & I (1996): J. Vasarietis REG: 2001. Inside strong purplish pink (55B) towards apex, gradually shading to light purplish pink (55C) in mid petal with an indistinct white area (c20 × 20mm) above nectary, throat strong purplish pink (55B); outside strong purplish pink (55B); with age all colours fade; numerous, tiny, black spots on the lower ⅓ of each petal; nectaries short, white; pollen brown; stigma greyish brown. Fls 140mm wide; petals 70 × 32mm, slightly waved and noticeably pinched in towards apex, inner petals noticeably broader, tips slightly recurved. Lvs alternate, 90 × 9mm. Stems 0.8m, greenish brown, with c11 fls, including some secondaries. Early to mid July. (Standard: colour print provided by registrant (WSY 0030816). Etymology: 'Vivija' is the pet name of the registrant's wife (Violeta).)

I (a-b/b-d) 'Vlad'ka'

cv. ('Kain' × 'Blizzard') R & N: B. Mićulka I & REG: 2002. Inside white, with a pale beige tint, centre light beige; outside with greenish and slightly pinkish midveins; hint of a brown-red, line-like brushmark on midvein inside, very few spots; nectaries green, white-granulated; pollen brown; stigma pale green. Fls 125mm wide; petals 85 × 45mm, slightly recurved (outer petals more so), slightly ruffled; pedicels 75mm long, glossy brown. Lvs alternate, 80 × 17mm, yellowish green. Stems 1.05m, slightly brown spotted and with a moderately long inflorescence. Mid July.

I (a/-) 'Wendenschloss'

cv. (Parentage unknown) R (pre-1992) & N: G. Steinbrück. Fls carmine red, with an orange sheen.

I (a-b/-) White Glory Group

grp (Selection from Futura Group) R, N & I (1982): Blackthorne Gardens. Inside white, with a few rose maroon spots in centre of flower. Stems 0.75m.

VII (-/-) 'White Heron'

cv. (Parentage unknown) R: F. Jury, pre-1998. Fls large, pure white.

IX (c/d) 'White Orchid'

See *speciosum* 'White Orchid'.

IX (b-c/a) 'White Satin'

See *longiflorum* 'White Satin'.

VII (b/b) 'White Surprise'

cv. (Parentage unknown) R (pre-1985), N & REG (2002): Sun Valley Bulb Farms. Inside shiny white (155D), with a white (155D) throat and pink markings; outside white (155D), outer petals with greenish midveins; nectaries green; spots and papillae present; pollen orange-brown; stigma pale grey-green. Petals with strongly ruffled margins and slightly recurved tips. Stems light green/greyish, with strongly scented fls.

I (a/-) Wilconking Group

grp (*wilsonii* var. *flavum* × 'Connecticut King') R (pre-1982) & N: L. Woodriff. Fls yellow-orange, flushed old gold, tips yellow; with maroon spots on ½ of each petal; nectaries white; pollen dark orange. Fls 140mm wide; petals 85 × 40mm, ruffled, tips slightly recurved. Lvs alternate, 95 × 22mm, dark green. Stems 0.8m, dark green, with c10 fls. July.

VIII (a-b/-) 'Wild Treasure'

cv. (*longiflorum* 'Lorina' × 'Muscader') H (1993), G (1995), N (2000), I (2002) & REG (2002): De Jong Beheer B.V. Inside white (155C), base of midvein yellow-green, shading apically into vivid purplish red (57C); outside white (155B); nectaries green; pollen red-brown; stigma grey; spots and papillae absent. Petals 162mm long, slightly ruffled. Lvs alternate, 230 × 30mm, more impressed than in 'Sea Treasure'. Stems 1.05m, green, with c6 fls. Late. (A *longiflorum*/oriental hybrid.)

I (a/b-d) 'Yellowlita'

cv. ('Aelita' × Yellow Blaze Group) H (1991) & G (1994): G. Steinbrück N, I & REG: 2002. Inside vivid yellow (9A), shading to yellowish white (158D) at tips; outside yellowish white (158D), with pale green midribs; many, large, elliptic, deep purplish red (61A) spots over ¾ of each petal; nectaries pale yellow-green (157C) hoary; pollen moderate red (181A), anthers on long, thin, very pale greenish yellow filaments; stigma greyish brown (166A) on a long, dark greenish yellow style. Fls 130mm wide; petals 70 × 37mm, not ruffled, tips strongly recurved. Lvs alternate, 75 × 18mm. Stems 1.0m, dark green, with c9 fls. Mid July. (Standard: colour print provided by registrant (WSY 0032273). Etymology: name derived from parts of the name of both parents.)

I (a-b/-) Yellow Tiger Group

grp (Parentage unknown) H (pre-1963) & G: N.A. Lamb I: F.M. Abbey. Fls in varying shades of yellow and orange. July. (Published in *Gardenside Gossip* 28: 8 (1963); unacceptable name (Art. 17.13).)

I (a/-) 'Yolanda'

cv. (Parentage unknown) R (pre-1985) & N: Oregon Bulb Farms. Bright yellow, with a few dark spots on basal ¼ of each petal. Stems 1.0-1.2m. (Unacceptable name (Art. 17.12).)

I (a/b-c) 'Zauberflöte'

cv. (Un-named seedling × mixed pollen of 'Mandarin' and 'Snowcap') H (1989), G (1992), N (1999) & I (2001) & REG: G. Steinbrück REG: 2002. Inside pale yellow (158A), tips yellowish white (158D); outside similarly coloured with soft green midribs which are edged with pale pink; numerous, small, deep reddish purple (77A) spots

on basal $\frac{1}{2}$ of each petal; nectaries soft green; pollen strong orange (169B); stigma moderate orange (173C). Fls 150mm wide; petals 82×37 mm, not ruffled, tips very slightly recurved. Lvs alternate, 93×18 mm. Stems 0.7m, dark green, with c8 fls. Late June. (Standard: colour print provided by registrant (WSY 0032274).)

I (c/-) 'Zeus'

cv. (*lancifolium* hybrid) L: Blackthorne Gardens, 1982. Fls golden yellow, with dark spots. Petals recurved. Stems 1.2-1.5m. July. (Unacceptable name (Art. 17.2).)

I (a/b) 'Ziedoklis'

cv. (Parentage unknown) H (1990), G (1993), N (1996) & I (1996): J. Vasarietis REG: 2001. Inside vivid orange (28A) with a pink overlay, throat vivid orange (28A); outside vivid orange (28B); black spots of moderate size on basal $\frac{1}{2}$ of each petal; nectaries white; pollen brown; stigma greyish. Fls 130mm wide; petals 70×40 mm, not ruffled, tips slightly recurved. Lvs alternate, 100×12 mm. Stems 1.05m, nearly black at base, greenish brown above, with c19 fls in a pyramidal inflorescence. Mid July. (First published in *Lilium Balticum* Lily Register 1996. Standard: colour print provided by registrant (WSY 0030819). Etymology: Ziedoklis is Latvian for the word credence.)

I (b/c) 'Zimtstern'

cv. ('Winnetou' \times 'Goldprinzessin') H (1992) & G (1995): G. Steinbrück N, I & REG: 2002. Inside vivid reddish orange (32A), with a suffusion of pale orange-yellow (24A) (especially along centre in basal $\frac{1}{2}$); outside vivid reddish orange (32A); numerous, small, round, dark red (187B) spots on basal $\frac{1}{2}$ of each petal; nectaries light yellowish pink (159A); pollen dark red (187A); stigma moderate red (179A). Fls 140mm wide; petals 70×40 mm, not ruffled, not recurved. Lvs alternate, 60×20 mm. Stems 0.7m, dark green, with c11 fls. Mid June. (Standard: colour print provided by registrant (WSY 0032241).)

CORRECTIONS TO THE REGISTER

VII (b/b) 'Dali'

cf. *The International Lily Register, Nineteenth Supplement*, p. 14 (2001). Published in error as Div. VIII; it is Div. VII.

I (a/-) 'Dublin'

cf. *The International Lily Register, Nineteenth Supplement*, p. 16 (2001). This entry is for the same plant here listed as 'Darwin'. The latter name has been used as part of a grant of Plant Breeders' Rights (LEL 1759, The Netherlands) and the use of this later epithet is sanctioned.

VIII (a/b) 'Eurostar'

cf. *The International Lily Register, Twentieth Supplement*, p. 18 (2002). The Standard for this cultivar cannot be the illustration indicated, but must be the documentation held by the Plant Breeders' Rights authority in The Netherlands where rights were granted in 1999 (LEL 1481).

I (a/b) 'Ina'

cf. *The International Lily Register, Nineteenth Supplement*, p. 24 (2001). The petal measurements should be 75×40 mm.

VIII (a/b) 'Palermo'

cf. *The International Lily Register, Nineteenth Supplement*, p. 36 (2001). This is a *longiflorum*/asiatic hybrid.

VIII (a-b/b) 'Serrada'

cf. *The International Lily Register, Nineteenth Supplement*, p. 45 (2001). This is a *longiflorum*/asiatic hybrid.

V (b/a) 'Worldwhite'

cf. *The International Lily Register, Nineteenth Supplement*, p. 56 (2001). This cultivar is probably best placed as a Div. IX *longiflorum* cultivar.

LIST OF RAISERS, REGISTRANTS AND INTRODUCERS

(† = deceased)

†ABBEY, F.M.	Formerly of North Ferrisburg, Vermont, USA
ALEXANDER, A.E.	Christchurch, New Zealand
ASANO, Y.	Hokkaido University, Sapporo, Japan
BALDWIN, M.K.	Ellinbank Nurseries, Ellinbank, Victoria, Australia
BALL, J.	Regina, Saskatchewan, Canada
†BARRY, H.E.	Formerly of Papatoetoe, Auckland, New Zealand
BAZETT, R.H.	Kelowna, British Columbia, Canada
B & D LILIES	Port Townsend, Washington, USA
†BEMIS, R.W.	Formerly of Blackthorne Gardens, Holbrook, Massachusetts, USA
BERG, M.A. van den	't Zand, The Netherlands
BEUTNAGEL, O.	Braunschweig, Germany
BISCHOFF TULLEKEN LELIES B.V.	Wieringerwerf, The Netherlands
†BLACKTHORNE GARDENS	Formerly of Holbrook, Massachusetts, USA
BÖHM, H.	Dresden, Germany
BOON, Fa. Jaap	Hem, The Netherlands
BORBELETA GARDENS	Faribault, Minnesota 55021, USA
BOS FLOWERBULBS, Van den	Honselersdijk, The Netherlands
BOTTEMA, G.	Berkhout, The Netherlands
BOTTEMA & Zn, Mts S.E.	Berkhout, The Netherlands
BOWEN, Mrs V.E.	Rockville, Maryland, USA
BRAUN, K.	Rousínov, Czech Republic
BURNETT, A.J.	Vermilion, Alberta, Canada
†CEBECO LILIES Inc.	Formerly of Aurora, Oregon, USA
CHANDLER, G.M.	Victoria, Australia
COLUMBIA-PLATTE LILIES	Boring, Oregon 97009, USA
COÖP. KWEKERSVERENIGING "THE ORIGINALS" u.a.	Sint Maartensvlotbrug, The Netherlands
CPRO	Wageningen, The Netherlands
DE JONG LELIES HOLLAND B.V.	Andijk, The Netherlands
DENKEWITZ, L.	Hamburg, Germany
DIERSSEN, H.	Lemgo, Germany
†DOERR, E.H.	Formerly of New Ulm, Minnesota, USA
DOUWE WIERSTRA B.V.	Andijk, The Netherlands
DUDEK, J.	Hlučín, Czech Republic
EWALD, F. & G.	Leifherde/Gifhorn, Germany
FAIRDALE NURSERY	Wilsonville, Oregon, USA
FEDOROV, A.M.	Volzhok Mary El Republic, Russia
†FELDMAIER, C.	Formerly of Pfarrkirchen, Germany
FELLNER, F.	Vermilion, Alberta, Canada
FOSTER, M.C.	Sevenoaks, Kent, England
FOX, E.E.	Millet, Alberta, Canada
FOX LILY RANCH	Millet, Alberta, Canada
FUJISHIMA, S.	Hokkaido, Japan
GEERING, G. & A.	Ashburton, New Zealand
GEISER, P.	Münchenstein, Switzerland

GEORGI, H.
 HARRISON, R.E.
 HAY, D.J.
 HAYLER, B.G.

HOFF & Zn, Fa. P.
 HOFFGAARDE B.V.
 HONEYWOOD LILIES
 HOOGLAND, P.M.
 IMANSE B.V., H.P.
 IMANSE LELIEVEREDELING B.V.
 †JURY, F.

JURY, M.
 KAISER, J.-M.
 KAPIAS, J.
 KIREEVA, M.F.
 KLEIN, J.
 KÖPKE, G.
 KOVÁŘ, J.
 KROMHOUT, D.
 & HOGEWONING, C.

LAAN Bros, N. & B.
 †LAMB, N.A.
 LIGTHART, W.
 LILY NOOK, The
 †LIMERICK, Lord
 MAK B.V., Gebr.
 MAK-LEEK, Inc.
 MAK LELIEVEREDELING B.V.
 MARKLILY C.V.
 †MARKS, J.
 MARTIN, T.R.
 MIČULKA, B.
 MIROSHNICZENKO, Mrs N.A.
 MIYOSHI & CO. Ltd
 MÜLLER, P.
 NELSON, D.
 NORTH, C.

OREGON BULB FARMS Inc.
 †OREKHOV, V.P.
 PAAUW & Zn, W.
 PARKER-JERVIS, †J. & E.

PÄTZOLD, G.
 †PAYNE, F.H.

PEACOCK B.V.
 PETRUSKE, J.
 PIASSEK, F.
 †PYNAERT van GEERT, E.

Birkenwerder, Germany
 Palmerston North, New Zealand
 Saskatoon, Saskatchewan, Canada
 Formerly of Golden Ray Gardens,
 Kallista, Victoria, Australia
 See Hoffgaarde B.V.
 Steenberg, The Netherlands
 Parkside, Saskatchewan, Canada
 Dirksborn, The Netherlands
 Voorhout, The Netherlands
 Lisserbroek, The Netherlands
 Formerly of Waitara, Taranaki,
 New Zealand
 Urenui, New Zealand
 Germany
 Radostowice, Poland
 Michurinsk, Russia
 Obersalbach, Germany
 Neuss, Germany
 Volyně, Czech Republic
 Rijnsburg, The Netherlands

Bovenkarspel, The Netherlands
 Formerly of Mystic, Connecticut, USA
 't Zand, The Netherlands
 Neepawa, Canada
 Formerly of West Hoathly, Sussex, England
 't Zand, The Netherlands
 Anna Paulowna, The Netherlands
 The Netherlands
 Lisse, The Netherlands
 Formerly of Hořice, Czech Republic
 Binbrook, Ontario, Canada
 Velehrad, Czech Republic
 Zhitomir, Ukraine
 Tokyo, Japan
 Dossenheim, Germany
 Saskatchewan, Canada
 Formerly of Scottish Horticultural
 Research Institute, Invergowrie,
 Scotland
 Aurora, Oregon, USA
 Formerly of Jēkabpils, Latvia
 Rijnsburg, The Netherlands
 Longworth, Abingdon, Oxfordshire,
 England
 Wurzen, Germany
 Formerly of Piedmont Gardens,
 Waterbury, Connecticut, USA
 Rijnsburg, The Netherlands
 Berlin, Germany
 Immekath, Germany
 Formerly of Ghent, Belgium

†RECH, A. RIVERSIDE GARDENS ROHRMEIER, M. ROLLINSON, M.E. ROOYAKKERS B.V., H.J.A. SANDE B.V. SCHAEFER, Mrs D.B. SCHARF, A. SCHENK, P.	Formerly of Heusweiler, Germany Riverside Estates, Saskatchewan, Canada Address unknown Timaru, New Zealand Breezand, The Netherlands 't Zand, The Netherlands Waukee, Iowa, USA Canada Bischoff Tulleken Lelies B.V., Wieringerwerf, The Netherlands Berkhout, The Netherlands Berkhout, The Netherlands Formerly of Port Townsend, Washington, USA Edmonton, Alberta, Canada Ogre, Latvia Berlin, Germany Formerly of Strahm Bulb Farm, Harbor, Oregon, USA Harbor, Oregon, USA Formerly of Erlangen-Kriegenbrunnen, Germany Arcata, California, USA Kita-uonuma-gun, Niigata Prefecture, Japan Plymouth Meeting, Pennsylvania, USA Kyoto, Japan Hillegom, The Netherlands
SCHOUTEN, J.J. S. en S., V.o.f. †SHAVER, J.M.	Edmonton, Alberta, Canada Ogre, Latvia Berlin, Germany Formerly of Strahm Bulb Farm, Harbor, Oregon, USA Harbor, Oregon, USA Formerly of Erlangen-Kriegenbrunnen, Germany Arcata, California, USA Kita-uonuma-gun, Niigata Prefecture, Japan Plymouth Meeting, Pennsylvania, USA Kyoto, Japan Hillegom, The Netherlands
SIMONET, R. ŠKIMELE, D.V. STEINBRÜCK, G. †STRAHM, H.J.	Edmonton, Alberta, Canada Ogre, Latvia Berlin, Germany Formerly of Strahm Bulb Farm, Harbor, Oregon, USA Harbor, Oregon, USA Formerly of Erlangen-Kriegenbrunnen, Germany Arcata, California, USA Kita-uonuma-gun, Niigata Prefecture, Japan Plymouth Meeting, Pennsylvania, USA Kyoto, Japan Hillegom, The Netherlands
STRAHM BULB FARM †STRASSER, V.	Harbor, Oregon, USA Formerly of Erlangen-Kriegenbrunnen, Germany Arcata, California, USA Kita-uonuma-gun, Niigata Prefecture, Japan Plymouth Meeting, Pennsylvania, USA Kyoto, Japan Hillegom, The Netherlands
SUN VALLEY BULB FARMS SUZUKI, W.	Arcata, California, USA Kita-uonuma-gun, Niigata Prefecture, Japan Plymouth Meeting, Pennsylvania, USA Kyoto, Japan Hillegom, The Netherlands
SWOSZOWSKI-TRAN, K. TAKII & CO. Ltd TESTCENTRUM VOOR SIERGEWASSEN THÜRMER, G. TIL-HARTMAN B.V., van UNGERMAN, A.N. VAN DER SALM (HOLLAND) VASARIETIS, J. VEG, Zierpflanzen VELDE, A.J. van den VENARD, J. Mrs VLETTER & DEN HAAN BEHEER B.V. VLIER, M. van VOLKOV, V.A. †WADEKAMPER, J. †WEINREICH, A. WERELD, J.W.A. van der WINKLER, A. †WITTNEBEN, C. †WOODRIF, L.	Weinböhl, Germany Hillegom, The Netherlands Kishinev, Moldavia Den Helder, The Netherlands Vecumnieki, Bauskas raj., Latvia Erfurt, Germany c/o H.P. Imanse B.V. Waukee, Iowa, USA Rijnsburg, The Netherlands 't Zand, The Netherlands Dmitrov, Moscow, Russia Formerly of Parkdale, Oregon, USA Formerly of Wolmierstedt, Germany Breezand, The Netherlands Kleinsendelbach, Germany Formerly of Luchow, Germany Formerly of Mckinleyville, California, USA Breezand, The Netherlands Breezand, The Netherlands Formerly of Riga, Latvia Bensheim, Germany Riga, Latvia
WORLD BREEDING WORLD FLOWER B.V. †ZELTAN, K. ZIEPKE, S. ZORGÉVIC, A.	Breezand, The Netherlands Breezand, The Netherlands Formerly of Riga, Latvia Bensheim, Germany Riga, Latvia