

## THE INTERNATIONAL LILY REGISTER & CHECKLIST TWENTY-FOURTH SUPPLEMENT

The following list contains details of lilies whose epithets were registered between 1 October 2004 and 30 September 2005. In addition there are details of a few lilies whose epithets have not been registered: these form part of the Checklist function.

Raisers and introducers are urged to ensure that all their plants have been registered and are reminded that registration should take place before a plant is released. Registration forms can be obtained from the International Lily Registrar (address below), or downloaded from the RHS web site ([www.rhs.org.uk](http://www.rhs.org.uk)).

The International Cultivar Registration Authority (ICRA) is indebted to the regional representatives for the effort they make to obtain and pass on information about lily cultivars. They are as follows:

Regional representative: Australia

Neil Jordan, 27 Scarfe Street, Camdale, Tasmania 7320, Australia.

Regional representative: Czech Republic

Ing. Břetislav Mičulka, 687 07 Velehrad 225, Czech Republic.

Regional representative: The Netherlands

Saskia Bodegom, Koninklijke Algemeene Vereeniging voor Bloembollencultuur (Royal General Bulbgrowers' Association), Postbus 175, 2180 AD Hillegom, The Netherlands.

Regional representative: New Zealand

Mrs Merle I. Lepper, P O Box 1394, Christchurch, New Zealand.

Regional representative: North America

Mrs Kathy Andersen, 2565 Charlestown Road, Phoenixville, PA 19460, USA.

This Supplement continues the use of a dual subdivision for every entry. Thus, both flower aspect (up-, out- or down-facing) and flower form (trumpet, bowl, flat or recurved) are indicated for each entry when this information is available. The existing lettering is retained, with flower aspect given first, separated from the flower form by a slash mark (/). An entry such as I(b/c) 'Aloft' therefore indicates that this is an Asiatic lily with out-facing, flat flowers. It is not a requirement that dual subdivisions should be used at all times. The way in which the information is presented means that, where it is necessary for a particular purpose, one (or both) can be ignored. Their inclusion, however, together with the division number, helps to present in an abbreviated form a quick 'picture' of the flower concerned. For a full account of any plant it is necessary to refer to the description.

The following changes have been introduced into the lily descriptions. Petals are now referred to as tepals (a more botanically correct term). Alternate leaves are now referred to as scattered: the latter term is more generally accepted in lily literature and more accurately describes the leaf arrangement. The longitudinal central vein that is present in each tepal is now given the following terminology: the vein on the upper surface (inside) of a tepal is a midvein, whereas the vein on the lower surface (outside) is a midrib.

Victoria Matthews, International Lily Registrar, 13 Chatsworth Close, Market Deeping, Peterborough PE6 8AZ, United Kingdom. Tel: +44 (0)1778 380 256. E-mail: [victoriamatthews@rhs.org.uk](mailto:victoriamatthews@rhs.org.uk)

### **VII(a/b-c) 'Action'**

cv. Parentage: Seedling 90-007 × seedling 90-058-1

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside vivid purplish red (57C/D), towards tip and margins almost white, midveins yellow at base changing upwards to deep red (53A); outside vivid purplish red (57D); spots dark red (187A), on basal half of tepals; papillae present; nectaries green changing into yellow; filaments pale green; pollen moderate reddish orange (171A); stigma dark red (187A). Fls 180 mm wide; tepals 110-115 × 44-62 mm, margins ruffled, tips recurved. Lvs scattered, 135-175 × 34-50 mm, dark green. Stems 0.95-1.4 m, green, with 3-10 fls. Midseason.

(Nomenclatural standard: colour print supplied by registrant (WSY 0096389).)

### **VIII(a-b/b) 'Albany'**

cv. Parentage: Seedling E819-1 × seedling 93,40-033-1

R: World Breeding B.V., N: World Breeding B.V. (2001)

Inside light greenish yellow (5D), each tepal with a deep red (53A) flame from nectary, midveins yellow at base; outside pale greenish yellow (2D), flushed with red at base; spots absent; papillae present; nectaries yellow-green; pollen strong reddish orange (169A); stigma orange with purple spots. Fls 190 mm wide; tepals 140 × 27-64 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 13-25 × 49-65 mm, dark green. Stems 0.9-1.2 m, dark green, with 2-9 fls. Late.

(An Orienpet (OT) lily. Registration was refused under ICNCP, 2004: Art. 19.25, because the name is too similar to 'Albanie', however it must be accepted because Dutch PBR were granted in 2004 (ICNCP, 2004: Art. 24.4). Nomenclatural standard: colour print supplied by registrant (WSY 0096409).)

### **VIII(a-b/b) 'Altari'**

cv. Parentage: Seedling 819-18 × seedling 90-033-1

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside dark red to deep red (187C/53A), pale yellow-green (155A) towards tepal-margins; outside yellowish white (155D) with some purple-red; spots dark red to deep red (187C/53A); papillae present; nectaries yellow-green; pollen orange-brown; stigma purple, spotted. Tepal-margins slightly ruffled, tips recurved. Lvs scattered, dark green. Stems medium to tall, with c.10 fls. Late.

(An Orienpet (OT) lily.)

### **VIII(a-b/b) 'Amarna'**

cv. Parentage: Seedling PH 94-01 × seedling RH 96-03

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals brilliant greenish yellow (6C), pale yellow-green (4D) towards margins and tips, margins light greenish yellow (5D), throat yellowish white (155D), outer tepals brilliant greenish yellow (6C) but less yellow; outside with inner tepals pale yellow-green (155A), margins light greenish yellow (5D), midribs strong yellow-green (144C) at base changing to dark yellowish pink (181D) towards top, outer tepals pale yellow-green (155A), with dark yellowish pink (181D) along midribs which are strong yellow-green (143C) at base and top; spots absent; papillae brilliant greenish yellow (6C); nectaries strong yellow-green (143C); pollen dark red (183B); stigma deep purple (79C). Fls 235 mm wide, fragrant; tepals 170 × 51-81 mm, margins ruffled, tips recurved. Lvs scattered 181 × 30 mm. Stems 1-1.3 m, green with darker markings, and c.3 fls.

(An Orienpet (OT) lily.)

### **VIII(a-b/b) 'Ambiente'**

cv. Parentage: Seedling PG 95-57 × seedling RH 95-01

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light yellow (10C), brilliant greenish yellow (6B) on the narrow margins and along midveins, outer tepals light greenish yellow (8C), brilliant yellow (8A) on the narrow margins and along midveins, throat pale yellow-green (4D); outside with inner tepals pale yellow (8D), light greenish yellow (5D) towards margins, midribs moderate yellow-green (147D), at base dark pink (182C) changing to light greenish yellow (5D), outer tepals pale greenish yellow (2D), dark pink (182C) along midribs, at base moderate yellow-green (147D) and dark pink (182C), strong yellow-green (143C) at top; spots absent; papillae brilliant greenish yellow (6B); nectaries strong yellow-green (143B); pollen moderate orange (167B); stigma dark red (187A). Fls 250 mm wide, fragrant; tepals 155 × 58-82 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 195 × 28 mm. Stems 1.55 m, green with darker margins, and with c.4 fls. (An Oripenet (OT) lily.)

### **I(b-c/c) 'Anna Theresia'**

cv. Parentage unknown

R: J. Wadekamper, N: Lily Company B.V. (2004), REG: Lily Company B.V. (2004)

Inside vivid yellow (12A), each tepal with a dark red (183A) flame, throat brilliant greenish yellow (2B); outside vivid yellow (12A), tepal-margins vivid yellow (13A); spots dark red (183A); papillae absent; nectaries strong yellow-green (143B); pollen moderate reddish orange (171A); stigma dark red (59A). Fls 120-155 mm wide; tepals 90 × 30-40 mm, margins smooth to slightly ruffled, tips recurved; peduncles with felty white hairs especially at base. Lvs scattered, 137 × 9 mm. Stems 0.8-1.15 m, green with darker markings, with c.14 fls.

### **VII(c/b) 'Antonia'**

cv. Parentage unknown

R: J.W.A. van der Wereld ?, N: Lily Company B.V. (2004), REG: Lily Company B.V. (2004)

Inside dark purplish pink (186C), inner tepals with light greenish yellow (4B) above nectaries; outside with outer tepals dark to moderate purplish pink (186C/D), midribs strong yellow-green (144B) at base, changing from moderate purplish red (185C) to brilliant yellow-green (154C), at top some strong yellow-green (144A), inner tepals similar in colour but lacking green top; spots absent; papillae pale yellow-green (155A), pale yellow-green (4D) and moderate purplish pink (186D); nectaries strong yellow-green (144B); pollen strong brown (172A); stigma dark red/dark purple (187A/79A). Fls 150-250 mm wide, fragrant; tepals 144-150 × 39-46 mm, margins strongly ruffled, tips recurved. Lvs scattered, 165 × 30 mm, dark green with lighter veins. Stems 0.7-1 m, light green, with c.9 fls. (The young fls have more whitish inside.)

### **VII(a/b) 'Arabian Pearl'**

cv. Parentage: 'Medusa' × 'Solai'

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Beheer B.V. (2004)

Inside deep purplish pink (66C), moderate purplish pink (66D) towards yellowish white (155D) margins, moderate purplish pink (66D) along midveins (with age the colour becomes more light purple (75A) with (74B) midveins), throat yellowish white (155D); outside with outer tepals vivid purplish red (67B), deep purplish pink (66C) towards narrow yellowish white (155D) margins, midveins strong yellow-green (143C) at base and top, the middle part brilliant yellow-green (150A) in basal half and in apical

half moderate red (179A) with strong yellow-green (143C), inner tepals very pale purple (69D) at base changing into strong purplish pink (68B), towards margins moderate purplish pink (65A), from halfway up margins narrow, yellowish white (155D), midribs strong yellow-green (143C) at base, changing into brilliant yellow-green (150B), towards top strong purplish red (60D); spots few, strong purplish red (60D); nectaries strong yellow-green (144B) with narrow light greenish yellow (1C) edge; pollen strong orange (168A); stigma light purple (77C). Fls 210 mm wide, fragrant; tepals 130 × 50-65 mm, margins slightly ruffled, tips recurved. Lvs scattered, 165 × 30 mm. Stems 1.1-1.2 m, green with darker flush, and with 4-7 fls.

#### **VIII(a/b) 'Arctic Diamond'**

cv. Parentage: Seedling LA 93096-1 × 'Sancerre' (tetraploid form?)

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Beheer B.V. (2004)

Inside yellowish white (155D), throat light yellow-green (138D); outside yellowish white (155D), with little light purplish pink (65B) stripes from base to about halfway of each tepal along midrib, midrib brilliant yellow-green (142B) at base and top, inner tepals less light purplish pink (65B); spots few, dark red (59A); papillae yellowish white (155D); nectaries strong yellow-green (143B); pollen dark reddish orange (175B); stigma light greenish yellow (3D). Fls 160 mm wide; tepals 103 × 55 mm, margins slightly ruffled, tips recurved. Lvs scattered, 140 × 20 mm. Stems 0.9-1.15 m, green with darker markings, and with 2-7 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **VII(a/b) 'Armonia'**

cv. Parentage: Seedling 92-22 × seedling RF 94-01

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside deep purplish red (61A), tepal-edges dark purple (79A), along midveins strong purplish red (60B), throat yellowish white (155D); outside with outer tepals deep purplish red/strong purplish red (61A/60B), edges dark purple (79A), midribs light yellow-green (145B) at base, yellowish white (155D) up to half way, changing to light yellow-green (145D) with deep purplish red (61A), at top strong yellow-green (143B), inner tepals strong purplish red (60C), deep purplish red (61A) towards margins and with dark purple (79A) edges, midribs similar to outer tepals but lacking green top, spots absent; papillae dark red (59A), nectaries strong yellow-green (145A); pollen strong orange (25B); stigma light greenish grey (190D) with dark purple (79B). Fls 250 mm wide, fragrant; tepals 145 × 52-76 mm, margins slightly ruffled, tips recurved. Lvs scattered, 160 × 40 mm. Stems 0.9-1.4 m, red-brown with green, with 2-5 fls.

#### **I(a/c) 'Barča'**

cv. Parentage: 'Kolovrat' × 'Amarone'

R: J. Rezek (2001), N: J. Rezek (2005), I: J. Rezek (2005), REG: J. Rezek (2005)

Inside with deep yellow tepal-tips, centre darker; outside pale yellow, brownish at base; spots few, brown-violet, oblong, on a fifth of each tepal; papillae in throat; nectaries white-granulated; pollen chocolate-brown; stigma dark brown. Fls 160 mm wide; tepals 92 × 48 mm, margins ruffled, tips not recurved; pedicels 85 mm, brown-spotted. Lvs scattered, 115 × 18 mm, glossy dark green. Stems 0.85 m, thick, dotted with violet-brown, with dark green bulbils. Early July. Probably triploid.

#### **I(a-b/b-c) 'Barunka'**

cv. Parentage: ('Rusalka' × unknown) × 'Blizzard'

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside creamy with a yellowish centre, brownish at nectaries; outside pale pink; nectaries

white-granulated; spots small, brownish pink, fairly numerous, over three-quarters of each tepal; pollen chocolate-brown; stigma dark brown. Fls 125 mm wide; tepals 75 × 33 mm, slightly recurved, the outer tepals more so; pedicels 65 mm, shiny brown. Lvs scattered, 85 × 14 mm, dull mid green. Stems 0.9 m, thick, finely brown-dotted. Late June.

#### **VIII(a/b) 'Basic Diamond'**

cv. Parentage: Seedling 3473.vi-2008.rs zl × seedling LA 93064-14T

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside with inner tepals moderate purplish pink (68C) with a small pale yellow-green (157C) flame, midveins light yellow-green (144D) colouring towards top, outer tepals very pale purple (69B) with larger pale yellow-green (157C) flame, light yellow-green (144D) along midveins, strong purplish pink (68B) from half way up and surrounding veins, throat light yellow-green (144D); outside with outer tepals deep purplish pink (70C), midribs strong reddish purple (70B) at top, inner tepals very pale purple (73D), strong purplish pink (73B) along midribs; spots and papillae absent; nectaries light yellow-green (144B); pollen moderate reddish brown (175A); stigma brilliant yellow-green (142B). Fls 170 mm wide; tepals 110 × 37-52 mm, margins smooth, tips recurved. Lvs scattered, 155 × 19 mm. Stems 0.85-1.15 m, green with darker marking especially at base, and with 3-8 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **VI(c/d) 'Beautiful Victoria'**

cv. Parentage: 'Brno' × 'Caramba'

R: A. Mego (1994), N: A. Mego (2005), I: A. Mego (2005), REG: A. Mego (2005)

Inside white, tepals with apricot centre, throat gold-orange; outside white, midribs green; spots absent; nectaries deep green with yellow edges; pollen brown-red; stigma green. Fls 100 mm wide; tepals 100 × 30 mm, strongly recurved, margins wavy; pedicels 180 mm, brown-dotted, with secondary buds. Lvs scattered, 220 × 30 mm, dull mid green. Stems 1.5 m, brown-dotted, infl. long. Early July.

#### **I(a/b) BENFICA**

Trade designation of 'Zanlzfica'.

#### **VII(a/b) 'Benin'**

cv. Parentage: Seedling 89-59 × seedling 93-14

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside yellowish white (155D), flushed with yellowish white (155B) along midveins; outside with inner tepals yellowish white (155D), midribs strong yellow-green (143C) changing into pale yellow (11D) flush, pale greenish yellow (2D) at top, outer tepals flushed with purple-red at base; spots absent; papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen moderate reddish orange (171A); stigma light purple (81C). Fls 225 mm wide, fragrant; tepals 155 × 48-83 mm, margins strongly ruffled, tips recurved. Lvs scattered, 155 × 40 mm. Stems 1-1.2 m, green, with c.3 fls.

#### **I(b-c/b-c) 'Beskyd'**

cv. Parentage: (('Hornback's Gold' × 'Rhabus') × 'Apricot Supreme') × white-flowered tetraploid

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside pale salmon-apricot, tepal-tips and throat paler; outside pale salmon, midribs green-yellow, brownish at base; spots fairly numerous, large, oblong, dark pinkish brown, over a third of each tepal; nectaries white-granulated and ciliate; pollen orange-brown; stigma dark brownish crimson. Fls 130 mm wide; tepals 95 × 47 mm, thick,

somewhat recurved; pedicels 100 mm long, thick, dotted. Lvs scattered, 145 × 23 mm, shiny dark green. Stems 0.85 m, very thick, densely spotted dark brown, with a medium-length infl. Early July. Polyploid.

#### **VIII(a/b) 'Birgi'**

cv. Parentage: Seedling BM 95-02 × seedling RS 93-16

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2000), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside strong purplish red (63A), tepal-edges flushed with (59B), throat light yellow-green (144D); outside strong purplish red (68B), changing into strong purplish red (59D) towards tepal-margins; spots very few, dark red (59A); papillae absent; nectaries strong yellow-green (144A); pollen strong orange-yellow (163B); stigma brilliant greenish yellow (3B). Fls 250 mm wide; tepals 150 × 50-73 mm, margins smooth, tips recurved. Lvs scattered, 155 × 28 mm. Stems 1.15-1.40 m, green with darker markings at base, and with 4-5 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **VIII(a/b) 'Birthday'**

cv. Parentage: Seedling 92-136 × seedling 89-50-101-1

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside brilliant to light yellow (10A/C), lighter towards tepal-tops, midveins yellow at base; outside pale greenish yellow (10D), flushed with red; spots and papillae absent; nectaries green; pollen orange-brown; stigma purple. Fls 200 mm wide; tepal-margins very slightly ruffled, tips recurved. Lvs scattered, dark green. Stems 0.9-1.1 m, green with darker spots and stripes, with 2-10 fls. Late.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096390).)

#### **VIII(b-c/b-c) Blomberg Group**

grp Parentage: Yellow-flowered Trumpet × ('Black Beauty' × (*auratum* × *henryi*))

R: H. Dierssen (1998), N: H. Dierssen, I: H. Dierssen, REG: H. Dierssen (2004)

Inside very pale beige-apricot, tepal-centres deeper apricot, tepal-tips and throat whitish; outside with light pink streaks, and greenish midribs on outer tepals; spots brown and yellow, the latter raised, in lines on two-fifths of each tepal; nectaries green; pollen brown; stigma grey. Fls 135 mm wide; tepals 115 × 40 mm, moderately recurved, margins wavy; pedicels 105 mm, densely brown-dotted. Lvs scattered, 90 × 17 mm, dull dark green.

Stems 1.25 m, brown-dotted. Mid July.

(Orienpet (OT) lilies.)

#### **VIII(a/b) 'Boa Vista'**

cv. Parentage: Yellow-flowered seedling × (*longiflorum* × 'Aristo')

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside vivid yellow (9B), throat pale greenish yellow (1D); outside with outer tepals brilliant yellow (13C), inner tepals light greenish yellow (8B); spots very few, greyish red (178A); papillae vivid yellow (9B); nectaries moderate olive-green (137B); pollen deep orange-yellow (163A); stigma greyish red (178A). Buds hairy. Fls 190 mm; tepals 120 × 44-62 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 170 × 15 mm. Stems 1.22 m, green with darker markings, and with c.5 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **VIII(a-b/b) 'Boogie Woogie'**

cv. Parentage: Seedling 132-1 × seedling 88-17-1

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside pale yellow-green (4D), lighter toward tepal-tips, tepal-margins narrow, red-pink; outside pale yellow-green (4D); spots absent; papillae sparse; nectaries light green; pollen vivid orange (28B); stigma yellow-green. Fls 190 mm wide; tepals 125-130 × 34-54 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 171-240 × 38-65 mm, dark green. Stems 1-1.25 m, green with darker spots and stripes, and with 2-9 fls. Late. (An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096392).)

#### **VIII(a/b) 'Bourbon Brilliant'**

cv. Parentage: 'Dame Blanche' (tetraploid form?) × seedling LO 93003.2BT  
R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Beheer B.V. (2004)

Inside yellowish white (155D), greenish white (157D) along midveins, throat light yellow-green (142D); outside yellowish white (155D), midribs light yellow-green (144D) at base and top, greenish white (157D) in middle part; spots absent; papillae yellowish white (155D); nectaries light yellow-green (142D); pollen strong reddish-orange (31A); stigma light yellow-green (138D). Fls 270 mm wide, slightly fragrant; tepals 161 × 53-68 mm, margins smooth, tips recurved. Lvs scattered, 190 × 44 mm. Stems 0.9-0.95 m, green with darker markings at base, and with c.3 fls. (A *longiflorum*/Asiatic (LA) hybrid.)

#### **VIII(b/b) 'Bradano'**

cv. Parentage: Seedling PG 95-57 × seedling RH 95-01  
R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light greenish yellow (8C), brilliant yellow (8A) along midveins, outer tepals pale yellow (8D), brilliant yellow (11A) along midveins, tepal-margins light greenish yellow (8B), throat pale yellow (8D); outside with inner tepals pale yellow (8D), tepal-margins light greenish yellow (8C), outer tepals with strong purplish red-flushed (60C) midribs; spots absent; papillae brilliant yellow (8A); nectaries strong yellow-green (144A); pollen orange-brown; stigma dark purple (79A). Fls 210 mm wide, slightly fragrant; tepals 150 × 55-70 mm, margins smooth, tips strongly recurved. Lvs scattered, 170-220 × 30-5- mm, dark green. Stems 1.25-1.5 m, green with darker markings, with c.4 fls. (An Orienpet (OT) lily.)

#### **VII(b/b) 'Branko'**

cv. Parentage: Seedling 91-28 × seedling RM 95-40  
R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals pale purplish pink (65C), outer tepals very light purple (75C), margins very pale purple (69D), moderate purplish pink (65A) along midveins, throat yellowish white (155D); outside with inner tepals yellowish white (155D) with pale purplish pink (65C), midribs light yellow-green (154D), outer tepals strong purplish red (63B), light purplish pink (65B) towards margins, midribs light yellow-green (154D); spots absent; papillae few, deep purplish pink (70C); nectaries strong yellow-green (144C); pollen dark reddish orange (172B); stigma moderate yellow-green (139D). Fls 200 mm wide; tepals 130 × 50-70 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 130 × 45 mm, dark green. Stems 0.9-1.2 m, green, darker at base, with 3-6 fls.

#### **VII(a-b/b) 'Brenta'**

cv. Parentage: Seedling RF 94-05 × seedling RM 97-30  
R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter &

Den Haan Beheer B.V. (2004)

Inside with inner tepals dark red (59A/187B), margin in basal third yellowish white (155D), in apical two-thirds yellowish white with moderate purplish red (59C) veins, outer tepals (59B), narrow margin yellowish white (155D), dark red (59A) along midveins, throat yellowish white (155D); outside with inner tepals deep red (60B), margin in basal part veined yellowish white (155D), in apical part narrow, yellowish white (155D), midribs at base with a small moderate red (183C) dot, changing into deep red (60C) with some strong yellow-green (144C) dots; spots absent; papillae deep reddish purple (77A); nectaries light green with a narrow brilliant greenish yellow (1B) edge; pollen moderate orange (172D); stigma brilliant violet (92A). Fls 180 mm wide; tepals 110 × 38-54 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 165 × 35 mm. Stems 1-1.35 m, brownish with green, with c.3 fls.

#### **VIII(a/b) 'Bright Brilliant'**

cv. Parentage: 'Sea Treasure' (tetraploid form?) × 'Dame Blanche' (tetraploid form?)

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside yellowish white (155D), with a pale yellow (8D) stripe along midveins (more prominent on outer tepals); outside with outer tepals yellowish white (155D), with a pale yellow (8D) stripe along midribs changing into strong yellow-green (144C) towards top, inner tepals with pale yellow (8D) midribs that have some strong yellow-green (144C) at base; spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144B); pollen strong orange (25A); stigma light yellow-green (145C). Fls 295 mm wide, fragrant; tepals 165 × 60-80 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 190 × 45 mm. Stems 0.8-1.1 m, green with darker markings, and with 1-5 fls.

(A *longiflorum*/Oriental (LO) hybrid.)

#### **VIII(a/b) 'Bright Diamond'**

cv. Parentage: 'Esra' × seedling LA 93096-IT

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside with inner tepals yellowish white (155D), outer tepals yellowish white (155D), with a strong purplish red (59D) flush next to the light yellow-green (144D) throat; outside yellowish white (155D), midribs strong yellow-green (143C) at base and top, middle part flushed with strong purplish red (59D); spots and papillae absent; nectaries strong yellow green (143A); pollen dark reddish orange (175C); stigma light greenish yellow (3C). Fls 235 mm wide, slightly fragrant; tepals 130 × 50-65 mm, margins smooth, tips slightly recurved. Lvs scattered, 195 × 27 mm. Stems 1.1-1.45 m, green with darker markings, and with 2-8 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **I(a/b-c) 'Brūnactīņa'**

cv. Parentage: 'Rotaļa' × seedling 238-S-1

R: Ā. Zorgevics (1985), G: Ā. Zorgevics (1990), N: Ā. Zorgevics (2005), I: University of Latvia Botanical Garden (2005), REG: Ā. Zorgevics (2005)

Inside strong orange (26B), throat brilliant yellow-green (149B); outside cinnamon, spots brownish; nectaries light purple; pollen light brownish; stigma light purple. Fls 120 mm wide; tepals 70 × 20 mm, margins smooth, tips not recurved. Lvs scattered, 60 × 15 mm, glossy. Stems 1 m, brownish green, with c.9 fls. Bulbs white. Early July.

#### **VII(a/b) 'Carbonia'**

cv. Parentage: Seedling 88-99 × seedling 86-282

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter &

Den Haan Beheer B.V. (2004)

Inside with inner tepals strong purplish red (63B), strong purplish red (63A) along midveins, margins light purple (75A), outer tepals strong purplish red (63B), margins strong purplish red (61B), midribs vivid purplish red (61C), throat yellowish white (155B); outside with inner tepals yellowish white (155B) in basal third, becoming strong purplish red (64C), and finally strong purplish red (64B) at top, margins strong reddish purple (70B), midribs light green at base, yellowish white (155D) towards top, outer tepals strong purplish red (63B), towards margins strong purplish red (63A), yellowish white (155B) at base, next to midribs deep purplish pink (64D), midribs strong yellow-green (144C) at base and top, middle part yellowish white (155D); spots absent; papillae deep purplish red (59B); nectaries strong yellow-green (144A) with brilliant yellow (12B) edge; pollen strong orange (25A); stigma moderate yellow-green (138C). Fls 235 mm wide; tepals 130 × 55-75 mm, margins ruffled, tips recurved. Lvs scattered, 160-190 × 45-60 mm, dark green. Stems 0.9 m, light green with some darker markings, and with c.5 fls.

(The fls become a little paler in colour as they age.)

### VII(a/b) 'Carmina'

cv. Parentage: Red-flowered seedling × ('Rosalie' × 'Star Gazer')

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2002), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside with inner tepals strong purplish red (63A), moderate purplish red (66D) towards margins, base yellowish white (155D), strong red (53B) along midveins, outer tepals strong to vivid purplish red (67A/B), deep purplish red (68A) towards margins, base yellowish white (155D), strong red (46A) along midveins, throat yellowish white (155D); outside with inner tepals strong purplish red (63A), dark purplish pink (186C) towards margins, base light yellow-green (150D) changing into yellowish white (155D), strong purplish red (63A) along midribs, outer tepals moderate purplish red (64B), moderate purplish pink (66D) towards margins, base yellowish white (155D), midribs brilliant greenish yellow (151D) at base, strong purplish red (63A) towards top; spots absent; papillae deep red (60A); nectaries strong yellow-green (144C) with narrow light greenish yellow (1C) edge; pollen deep orange-yellow (163A); stigma greenish white (192D). Fls 225 mm wide, fragrant; tepals 135 × 45-60 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 140 × 35 mm. Stems 0.8-1 m, green, darker at base, with 6-8 fls.

### VIII(b/b-c) 'Carpaccio'

cv. Parentage: Seedling PG 96-59 × seedling RH 96-04

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light greenish yellow (5D), pale yellow-green (155A) along midveins towards top, outer tepals pale yellow-green (4D), light greenish yellow (8B) on the narrow margins and along midveins, throat pale yellow-green (155A); outside with inner tepals light greenish yellow (5D), pale yellow-green (155A) along midribs, strong yellow-green (144A) at base, outer tepals pale yellow-green (4D/155A), margins light greenish yellow (4C), at base with some strong greenish yellow (151A), along basal two-thirds of each midrib flushed with vivid purplish red (61C) (which fades with age), along apical third brilliant yellow-green (154B), apical fifth with light olive/strong yellow-green (152A/144A); spots absent; papillae brilliant yellow (8A); nectaries strong yellow-green (144A/B); pollen deep orange-yellow (163A); stigma dark purple (79A). Fls 200 mm wide; tepals 120-130 × 45-70 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 150-160 × 20-30 mm, dark green. Stems 1.2 m, dark green, with c.8 fls.

(An Orienpet (OT) lily.)

### VII(a/b-c) 'Cash'

cv. Parentage: 'New Age' × (unnamed seedling × 'Joy')

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside light purplish pink (73C) in apical two-thirds of each tepal, strong purplish pink (73B) towards margins, above each nectary is a brilliant yellow (12B) spot, at this level the area towards margins is yellowish white (155D), throat yellowish white (155D), outside with inner tepals light purplish pink (73C), towards margins strong purplish pink (73B), yellowish white (155D) at base, midribs brilliant yellow (14C) at base, yellowish white (155D) towards top, strong purplish pink (73B) from two-thirds upwards, outer tepals strong purplish pink (73B), yellowish white (155D) at base, midribs light yellow-green (144C) at base and top, middle part yellowish white (155D); spots absent; papillae yellowish white (155D) and brilliant yellow (12B); nectaries light yellow-green (144C); pollen moderate red/moderate reddish orange (180A/178C); stigma moderate purplish pink (186D). Fls 255 mm, fragrant; tepals 135 × 46-67 mm, margins slightly ruffled, tips recurved. Lvs scattered, 138 × 47 mm. Stems 1-1.15 m, reddish brown with green, with 6-8 fls.

### VIII(a/b) 'Cecil'

cv. Parentage unknown

R: Testcentrum voor Siergewassen B.V., N: Marklily C.V. (2001), REG: Marklily C.V. (2004)

Inside with inner tepals pale yellow-green (4D), light greenish yellow (5C) on margins and along midveins, margins and basal half of each tepal with dark red (187A/187B) spots, with outer tepals similar but lacking the light greenish yellow (5C) along midveins (young fls are light yellow (10B), brilliant yellow (8A) on margins and along midveins), throat pale yellow-green (4D) with dark red (187B) spot; outside greenish white (157D), flushed with pale yellow-green (4D) at base and along midveins, tepal-margins vivid yellow (9A) with dark red (187B) spots, tip strong yellow-green (144A), (young fls are pale yellow (8D), tepal-margins vivid yellow (9A) with spots; spots (see above); papillae absent; nectaries strong yellow-green (144B); pollen light olive-brown/light yellowish brown (199B/C); stigma strong greenish yellow (151C). Buds hairy. Fls 215 mm wide; tepals 137-144 × 45-65 mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs scattered, 180 × 16 mm, dark green. Stems 1.1-1.5 m, green, with 3-5 fls.

(*A. longiflorum*/Asiatic (LA) hybrid.)

### I(a/c) 'Cervia'

cv. Parentage: Seedling RS 91-23 × seedling AM 95-07

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light purplish pink (63D) with strong purplish red (63B) stripes, each nectary with a strong purplish red (63B) spot at top end, outer tepals more blue-pink than 63B, throat brilliant yellow-green (142B), yellowish white (155D) towards margin; outside light purplish pink (63D), very pale purple (69B) along midribs; spots absent; papillae strong purplish red (63B); nectaries brilliant yellow-green (142B); pollen moderate reddish brown (175A); stigma light yellow-green (142C). Fls 165 mm wide; tepals 85 × 45-55 mm, margins smooth, tips recurved. Lvs scattered, 95 × 15 mm, light green. Stems 0.95-1.2 m, green with darker markings, and with c.6 fls.

### VIII(a/b) 'Champagne Diamond'

cv. Parentage: Seedling LA 93007-24 × 'Sancerre' (tetraploid form?)

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside pale yellowish pink (27C), light yellowish pink (27B) towards tepal-margins, tips light orange (29B), throat strong yellow-green (144C); outside pale yellowish pink (27C/D), tepal-margins light yellowish pink (29C), along midribs with a moderate red/moderate purplish pink (182A/186A) stripe fading towards top into moderate purplish pink (186B) on a light ground; spots and papillae absent; nectaries pale yellowish pink (27D); pollen dark red/strong brown (187A/172A); stigma moderate reddish orange (173B). Fls 205 mm wide, fragrant; tepals 112-117 × 49-64 mm, margins smooth, tips recurved. Lvs scattered, 153 × 20 mm. Stems 1.1-1.15 m, brown-red, green towards top, with 2-9 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

### VIII(b/b) 'Charente'

cv. Parentage: Seedling PG 95-57 × seedling RH 97-07

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside pale greenish yellow (9D), brilliant yellow (9C) along midveins, tepal-margins light yellow (11B), throat yellowish white (155D); outside with outer tepals pale yellow-green (4D), midribs light yellow-green (145D) at base, area next to midribs vivid purplish red (57C) (absent in younger fls), inner tepals pale yellow-green (4D), midribs light yellow-green (145D) at base, margin pale greenish yellow (9D); spots absent; papillae vivid yellow (9B); nectaries strong yellow-green (144C); pollen strong brown (172A); stigma light purple (77C). Fls 210 mm wide, fragrant; tepals 140 × 50-80 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 175 × 41 mm, light and mid green. Stems 1.05-1.1 m, green with darker markings, and with c.5 fls. (An Orienpet (OT) lily.)

### I(b/b-c) 'Cheney Josette'

cv. Parentage: 'Catherine Laburi' × 'Sutherland Flamethrower'

R: D.J. Hay (2001), G: D.J. Hay (2003), N: D.J. Hay (2004), REG: D.J. Hay (2004)

Inside pink with gold overtones; outside pink; spots small, at base of tepals, surrounding nectaries; nectaries white; pollen orange; stigma pink. Fls 126 mm wide; tepals 63 × 30 mm, margins smooth, tips slightly recurved. Lvs scattered, 68 × 12 mm. Stems 0.45-0.5 m, dark brown, with c.6 fls. Midseason.

(Nomenclatural standard: colour print supplied by registrant (WSY 0096412).)

### I(a/b) 'Conamara'

cv. Parentage: Seedling AM 95-12 × seedling AW 94-09

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2000), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals with a large yellowish white (155B) spot changing towards margins and top into light purplish pink (65B), at margins and top moderate purplish pink (65A), outer tepals more moderate purplish pink (65A) at top, throat yellowish white (155D) changing into strong purplish pink (67D) towards edge; outside with inner tepals moderate purplish pink (65A) in basal half and along margins, apical half pale purplish pink (65C), midribs strong yellow green (144C) at top, outer tepals light purplish pink (65B) with narrow moderate purplish pink (65A) margins, light yellow-green (144D) along midribs, strong yellow-green (143C) towards top; spots few, on papillae, deep purplish red (59B); papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen vivid reddish orange (30B); stigma deep yellowish pink (43C). Fls 175 mm wide; tepals 95 × 36-54 mm, margins smooth, tips slightly recurved; pedicels hairy at base. Lvs scattered, 115 × 17 mm. Stems 0.85-1.4 m, green with darker markings, and with c.8 fls.

### **I(a/b-c) 'Conan'**

cv. Parentage: 'Kolovrat' × 'Amarone'

R: J. Rezek (2001), N: J. Rezek (2005), I: J. Rezek (2005), REG: J. Rezek (2005)

Inside dark purple-red, centre of tepals blackish; outside wine-red with green tepal-tips; spots absent; nectaries white-granulated; pollen dark purple; stigma dark purple. Fls 190 mm wide; tepals 115 × 49 mm, scarcely recurved; pedicels 100 mm, dark green.

Lvs scattered, 120 × 18 mm, glossy dark green. Stems 1 m, very thick, dark green. Early July. Probably triploid.

### **VIII(a/b) 'Condor Diamond'**

cv. Parentage: ((Seedling 1032-88-RSWI × 'Michella') × 'Esra') × seedling LA 93019-2T

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside brilliant yellow (7A), throat light greenish yellow (1C); outside light greenish yellow (4C), brilliant greenish yellow (6A) towards tepal-margins, light yellow-green (142C) at base; spots few, dark red (59A); papillae absent; nectaries strong yellow-green (143C); pollen strong brown (172A); stigma strong purplish red (60B) on greenish ground. Fls 185 mm wide; tepals 125 × 42-58 mm, margins smooth, tips slightly recurved. Lvs scattered, 143 × 18 mm. Stems 1.2-1.35 m, green with darker markings, and with 2-8 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

### **VIII(a-b/b-c) 'Contrast'**

cv. Parentage: Seedling 819-1 × seedling 90-033-1

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside pale yellow-green (155A), each tepal with a vivid red (57A) flame from nectary upwards; outside pale yellow-green (155A) with some purple red; spots and papillae present; nectaries yellow-green; filaments yellow-green; pollen red-brown; style yellow-green; stigma purple, spotted. Fls 170 mm wide; tepal-margins slightly ruffled, tips strongly recurved. Lvs scattered, dark green. Stems 0.9-1.1 m, green with darker spots and stripes, and with 2-7 fls. Very late.

(Nomenclatural standard: colour print supplied by registrant (WSY 0096391).)

### **VII(a/b) 'Copaya'**

cv. Parentage: Seedling 88-07-2 × seedling 89-225-3

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside vivid purplish red/deep purplish pink (67B/C), midveins vivid yellow (12A) at base changing into purple-red; outside deep to strong purplish pink (68A/B); spots dark purple-red on basal half of each tepal; papillae present; nectaries green; pollen strong reddish orange (31A); stigma grey-green. Fls 190-230 mm wide; tepals 130-135 × 45-63 mm, margins ruffled, tips slightly recurved. Lvs scattered, 110-253 × 30-38 mm, dark green. Stems 0.85-1.1 m, green with darker spots and stripes, and with 2-9 fls. Midseason.

(Nomenclatural standard: colour print supplied by registrant (WSY 0096394).)

### **VII(a/b) 'Coral Pearl'**

cv. Parentage: 'Melbourne' × 'Opus One'

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside moderate purplish red (186C), yellowish white (155D) above nectaries and on narrow tepal-margins, throat yellowish white (155D); outside with outer tepals moderate purplish red/moderate purplish pink (186C/D), yellowish white (155D) at base and on narrow margins, midribs strong yellow-green (143C) at base changing into moderate orange (170D); spots few, strong purplish red (61B); papillae yellowish

white (155D) and strong purplish red (61B); nectaries strong yellow-green (143C); pollen deep orange-yellow (163A); stigma light greenish grey (188C). Fls 240 mm wide, slightly fragrant; tepals 145 × 47-70 mm, margins ruffled, tips recurved. Lvs scattered, 170 × 28 mm. Stems 1.1-1.3 m, green, with 4-7 fls.

#### **VII(a/b) 'Corda'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside moderate purplish red (186B), tepal-margins narrow, strong reddish purple (70B), deep pink (48A) along midveins, throat greenish white (155C); outside with outer tepals moderate purplish red/deep purplish pink (186B/70C), greenish white (155C) at base, midveins strong yellow-green (144C) at base, moderate pink (35D) towards top, inner tepals dark purplish pink (186C), greenish white (155C) at base, strong reddish purple (70B) towards margins, midribs strong yellow-green (144C) at base, moderate pink (35D) towards top; spots absent; papillae dark red/deep purplish red (59A/B); nectaries strong yellow-green (144B), brilliant yellow (10A) towards midveins; pollen vivid yellowish pink (28A); stigma greyish yellow-green (148D). Fls 200 mm wide, strongly fragrant; tepals 125 × 60 mm, margins smooth to ruffled, tips slightly recurved. Lvs scattered, 170 × 38 mm, dark green with lighter veins. Stems 1-1.5 m, light green with darker markings, and with c.8 fls. (Colour of fls becomes paler with age.)

#### **II(c/d) 'Corneil's Yellow'**

cv. Parentage unknown

R: E.A. Corneil (pre 1975), G: E.A. Corneil, N: L.M. Barber (2005), REG: L.M. Barber (2005)

Inside vivid yellow (14B), throat brilliant yellow (14C); outside vivid yellow (14B), midribs strong yellow-green (144A); spots moderate red (179A), evenly distributed over whole tepal surface; nectaries strong yellow-green (144A); pollen deep orange-yellow (163A); stigma brilliant yellow (14C). Fls 51 mm wide, slightly scented; tepals 37 × 16 mm, margins smooth, tips recurved. Lvs whorled, 160 × 45 mm. Stems 1.3 m, strong yellow-green (144A), with c.11 fls in a pyramidal infl. June. (Misspelling: 'Cornell Yellow'. Named in memory of the raiser.)

#### **VII(b-c/b) 'Corte'**

cv. Parentage: Seedling RM 94-41 × seedling RM 97-08

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals strong purplish red (60B/C), moderate purplish red (70A) towards margins, outer tepals strong purplish red (60C/71B), throat greenish white (155C); outside with inner tepals strong reddish purple (70B), moderate to strong purplish red (64A/B) towards tip, midribs strong yellow-green (144C) at base changing into greenish white (157D), outer tepals deep to strong purplish red (71A/B), base strong yellow-green (144C), margins greenish white (155C) at base, pale yellow-green (155A) along midribs, strong yellow-green (144A) at top; spots absent; papillae dark red (187C); nectaries strong yellow-green (144A) with brilliant yellow (10A) edge; pollen strong brown (172A); stigma light purple (77C). Fls 210 mm wide; tepals 135-140 × 52-65 mm, margins slightly ruffled, tips recurved. Lvs scattered, 145 × 34 mm, dark green. Stems 0.85-1.2 m, light green with darker markings, with 3-4 fls.

#### **VIII(a/b) 'Cosenza'**

cv. Parentage: Seedling BW 94-03 × seedling CW 98-2

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (1999), REG: Vletter &

Den Haan Beheer B.V. (2004)

Inside yellowish white (155D) flushed with light pink, throat yellowish white (155D); outside yellowish white (155D), midribs strong yellow-green (143B) at base with moderate red (179B) flush; spots and papillae absent; nectaries strong yellow-green (143B); pollen light brown; stigma light greenish yellow (1C). Fls 165 mm wide, slightly fragrant; tepals 110 × 60 mm, margins smooth, tips recurved; pedicels hairy at base. Lvs scattered, 135 × 28 mm. Stems 1.1-1.3 m, green with some darker markings at top, and with c.6 fls.

### **I(a/b) 'Costa'**

cv. Parentage unknown

N: Sande B.V. (2001), REG: Brandenburg-Elshof B.V. (2005)

Inside light greenish yellow (4C) to pale yellow-green (4D); outside pale yellow-green (4D); spots and papillae absent; nectaries green; pollen brown; stigma yellow-green. Tepal-margins slightly ruffled, tips slightly recurved. Lvs scattered, dark green. Stems 1.1-1.25 m, green with darker markings, and with 3-6 fls. Midseason to late.

### **I(a-b/b-c) 'Crystal Baby'**

cv. Parentage: Seedling A-534-86 × seedling 90-83-042-1

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside pale yellowish pink (158D), lighter towards tepal-tips; outside pale yellowish pink (158D); spots and papillae absent; nectaries green; pollen brown; stigma purple. Tepals rather short and narrow, margins slightly ruffled, tips slightly recurved. Lvs dark green. Stems short, green with darker spots and stripes. Very early.

### **VI(b-c/d) 'Cyprián'**

cv. Parentage: 'Holubice' × unknown

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside pale apricot-beige, tepal-tips and margins paler; outside pale beige, midribs green-yellow, green at base; spots rusty, sporadic; papillae in lines; nectaries deep green; pollen brown; stigma green. Fls 150 mm wide; tepals 115 × 43 mm, very recurved, margins roughly waved, sometimes twisted; pedicels 125 mm long, with red-brown dots. Lvs 125 × 14 mm, dull dark green, dense. Stems 1.45 m, slightly dotted with red-brown. Mid July.

### **VIII(b/b) 'Daladier'**

cv. Parentage: Seedling PG 95-32 × 'Aubade'

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light yellow (10C), midveins brilliant yellow (8A), apical two-thirds of each tepal with area next to midveins pale yellow-green (155A), outer tepals light yellow (10C), brilliant yellow (8A) along midveins, throat pale greenish yellow (10D); outside with inner tepals light yellow/pale greenish yellow (10C/D), midribs light greenish yellow (4C) with adjacent area pale yellow-green (155A), outer tepals pale yellow-green (4D), margins pale greenish yellow (10D), midribs strong yellow-green (144B) in apical half, light yellow/pale greenish yellow (10C/D) at base with weak moderate purplish red (59C) flush; spots absent; papillae many, brilliant yellow (8A); nectaries strong yellow-green (144A); pollen strong orange (25A); stigma very pale purple (97D). Fls 200 mm wide, fragrant; tepals 110-130 × 45-70 mm, margins slightly ruffled, tips recurved. Lvs scattered, 130 × 20-30 mm. Stems 1-1.5 m, brown-red, green towards top, with 3-4 fls. (An Orienpet (OT) lily. The younger fls are more yellow.)

### **I(a-b/c) 'Dark Moon'**

cv. Parentage: Simonet Asiatic unnamed seedling × mixed pollen from Fellner Asiatic seedlings

R: J.J. Brown (1991), G: J.J. Brown (1992), N: J.J. Brown (2005), I: Manitoba Regional Lily Society (2005), REG: J.J. Brown (2005)

Inside dark red (187A), tepal-tips dark red (187B); outside similar but with slightly less lustre; spots darker than dark red (187A), raised; nectaries dark red (187A) with a silver glow; pollen dark brown; stigma dark crimson-brown. Fls 150 mm wide; tepals 90 × 40 mm, margins smooth, tips slightly recurved; pedicels 110-180 mm long. Lvs scattered, thick, 150 × 15 mm, dark green. Stems 1.5 m, brownish green, with 17 or more fls. Mid July.

(Published in *Manitoba Regional Lily Soc. Newsletter* 22(3): 8 (2005). Fls do not fade with age, but the density of the colour is affected by sunlight. Nomenclatural standard: colour print supplied by registrant (WSY 0096425).)

### **VII(a/b) 'Dassa'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals yellowish white (155D), each tepal with a light greenish yellow (7D) flame in basal half, outer tepals similar but with a vivid purplish red (57D) streak at tip (which disappears with age); outside with inner tepals yellowish white (155D), outer tepals yellowish white (155D), midribs strong yellow-green (144C) at base changing into light yellow-green (150D), at top strong yellow-green (143B), area next to midribs flushed with vivid purplish red (57D); spots absent; papillae many, brilliant yellow (7A); nectaries strong yellow-green (144C); pollen strong orange (25A); stigma light greenish (190C). Fls 235 mm wide, fragrant; tepals 125 × 57-75 mm, margins ruffled, tips recurved. Lvs scattered, 180 × 69 mm. Stems 1-1.1 m, green with darker markings, dark at base, with c.3 fls.

### **VII(a/b) 'Delos'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals dark/moderate purplish pink (186C/D), midveins light greenish yellow (3C) changing into moderate purplish pink (66D) towards top, throat greenish white (157D), outer tepals dark/moderate purplish pink (186C/D), area next to midveins deep purplish pink (70C), midveins light greenish yellow (3C) changing to moderate purplish pink (65A) towards top; outside with inner tepals light purple/very light purple (75B/C), narrow margin light purple (75A), base yellowish white (155D), midribs strong yellow-green (144C), outer tepals moderate purplish pink (186D), base yellowish white (155D), area next to midribs light yellowish pink (36B), midribs light yellow-green (145B); spots absent; papillae brilliant greenish yellow (3A) and vivid red (57A); nectaries strong yellow-green (144C); pollen dark reddish orange (175B); stigma greenish white (192D). Fls 214 mm wide, fragrant; tepals 135 × 54-75 mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs scattered, 150 × 59 mm. Stems 0.9-1.2 m, green with darker markings, with c.3 fls.

### **VII(a/b) 'Demeter'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside vivid purplish red (57C), towards margins strong purplish red (64C), along nectaries with a little light orange-yellow (16C), along midveins vivid purplish red

(57B), throat yellowish white (155D); outside with inner tepals vivid purplish red (57D), towards margins vivid purplish red (67B), yellowish white (155D) at base, midribs brilliant yellow-green (142B) at base, up to half their length deep purplish pink (58D), outer tepals strong purplish red (63A), yellowish white (155D) at base, midribs brilliant yellow-green (142B) at base, up to half their length deep purplish pink (58D); spots absent; papillae light orange-yellow (16C) and strong purplish red (60B); nectaries strong yellow-green (144C); pollen deep orange-yellow (163A); stigma very pale green (192C). Fls 230 mm wide, slightly fragrant; tepals 140 × 50-75 mm, margins slightly ruffled, tips recurved. Lvs scattered, 165 × 37 mm. Stems 1.1-1.2 m, lower third brownish, upper two-thirds green with brownish markings, and with c.7 fls.

#### VIII(b-c/b-d) 'Detmold'

cv. Parentage: Yellow-flowered Trumpet × ('Black Beauty' × (*auratum* × *henryi*))  
R: H. Dierssen (1997), N: H. Dierssen, I: H. Dierssen, REG: H. Dierssen (2004)  
Inside dark crimson, tepal-edges and tips cream, throat yellow; outside whitish with pink shining through, midribs greenish at base; spots dark brown-crimson, numerous, medium-sized to large, on two-thirds of each tepal; nectaries yellow-green; pollen brown; stigma pale green. Fls 160 mm wide; tepals 115 × 40 mm, reflexed, margins wavy; pedicels 115 mm, thick, strongly brown-dotted, with secondary buds. Lvs scattered, 155 × 36 mm, dull mid green. Stems 1.15 m, thick, strongly brown-dotted. Mid July.  
(An Orienpet (OT) lily.)

#### VIII(a/b) 'Dipo Lolo'

cv. Parentage: Pink-flowered seedling × (*longiflorum* × 'Aristo')  
R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V., REG: Bischoff Tulleken Lelies B.V. (2004)  
Inside pale purplish pink (65D), each tepal with moderate/light purplish pink (65A/B) flush along midvein in basal half, changing towards top into stripe along midvein, throat strong yellow-green (144C) with greenish white (157D) edges; outside with all midribs hairy, inner tepals pale purplish pink (65C), light yellow-green (144D) along midribs, outer tepals pale purplish pink (65D), margins strong purplish red/pink (63B/C), with strong yellow-green (144C) stripe/flush along midveins, at top a little dark purple (79A); spots dark purple (79A); papillae few; nectaries yellowish white (158D); pollen strong brown (172A); stigma dark greenish yellow (152D) with little strong purplish red (59D) stripes. Fls 150-195 mm wide; tepals 122 × 47-65 mm, margins smooth, tips slightly recurved. Lvs scattered, 180 × 26 mm, dark green. Stems 0.7-0.8 m, green, with c.3 fls.  
(A *longiflorum*/Asiatic (LA) hybrid.)

#### VIII(a-b/b) 'Donato'

cv. Parentage unknown  
R: Testcentrum voor Siergewassen B.V., N: Marklily C.V. (2001), REG: Marklily C.V. (2004)  
Inside with inner tepals moderate purplish red (186B), margins strong purple (77B), yellowish white (155D) around nectaries, along midveins strong purplish red (63A), at base on yellowish white (155D) ground with some strong orange-yellow (163B), outer tepals similar but margins moderate purplish red (186A), throat yellowish white (155D); outside with inner tepals moderate purplish red /dark purplish pink (186B/C), margins strong purple (77B), yellowish white (155D) around nectaries, midribs strong yellow-green (144A) at base, changing up to a sixth into light yellow-green (144B), apical five-sixths strong red (53B), outer tepals moderate purplish red (186B) in basal half, moderate purplish red (186A) in apical half, margins deep purplish red (59B), flushed with strong orange-yellow (163B) along midribs which are moderate olive-green (146A) at top; spots absent; papillae strong purplish red (60C); nectaries strong yellow-green

(144A); pollen dark reddish orange (175B); stigma dark red (187A/B). Fls 250 mm wide; tepals 160 × 40-70 mm, margins ruffled, tips recurved. Lvs scattered, 170-230 × 35-45 mm, dark green. Stems 1.4-2.05 m, green with darker markings, and with 5-8 fls. (An Orienpet (OT) lily.)

#### **I(a/b) 'Don Camillo'**

cv. Parentage: Seedling AM 96-28 × seedling AM 96-40

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals strong purplish pink (63C) with light purplish pink (73C) stripes, margins strong purplish pink (73B), outer tepals strong purplish pink (63C), margins strong purplish pink (73B), throat yellowish white (155B); outside with inner tepals strong purplish pink (63C), deep purplish pink (64D) towards margins, midribs light purplish pink (63D) surrounded by very pale purple (73D), outer tepals light purplish pink (62C), moderate purplish pink (62B) towards margins, strong purplish pink (62A) at margins, along midribs light yellow-green (150D), at top strong yellow-green (144C); spots absent; papillae yellowish white (155B); nectaries strong yellow-green (143C); pollen vivid yellowish pink (28A); stigma strong purplish red (59D). Fls 160 mm wide; tepals 102 × 40-53 mm, margins smooth, tips very slightly recurved. Lvs scattered, 135 × 25 mm. Stems 1.2-1.4 m, brownish red, with c.9 fls.

#### **I(a/c) 'Dot Com'**

This is the correct name for 'Dot.com' (published in *ILR Twentieth Supplement*). This

cv. was registered as 'Dot.com' but it was granted Dutch PBR in 2004 under the name 'Dot Com' and this latter name must be accepted (ICNCP, 2004: Art. 24.4).

#### **VIII(a-b/b-c) 'Dreamcatcher'**

cv. Parentage unknown

R: Sande B.V., N: Fa Q.J. Vink & Zn (2003), REG: Fa Q.J. Vink & Zn (2005)

Inside strong purplish red/deep purplish pink (54A/B); outside deep purplish pink/strong pink (54B/C); spots few; papillae present; nectaries light green; pollen red-brown; stigma orange-pink. Tepals short, margins slightly ruffled, tips slightly recurved. Lvs scattered. Stems small to mid height, green with darker markings. Early to midseason.

#### **VIII(a-b/b) 'Easly'**

cv. Parentage: Seedling E819-2 × seedling 88.22-1

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside vivid yellow (9B), lighter towards tepal-tips; outside brilliant yellow/pale greenish yellow (9C/D), greenish at base; spots brown-red; papillae present; nectaries green; filaments pale green; pollen (28B); stigma green with purple spots. Fls 200 mm wide; tepals 120-135 × 34-54 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 83-220 × 38-85 mm, dark green. Stems 0.8-1.25 m, green, with 2-9 fls. Early to midseason.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096393).)

#### **VIII(a/c) 'Elegant Crown'**

cv. Parentage: 'Sorbonne' × 'Goldina'

R: De Jong Lelies Holland B.V. (1996), G: De Jong Lelies Holland B.V. (1998), N: De Jong Lelies Holland B.V. (2003), I: De Jong Lelies Holland B.V. (2002), REG: De Jong Lelies Holland B.V. (2003)

Inside deep purplish pink (70C), tepal-margins and tips brilliant greenish yellow (6C),

along midveins strong purplish red (60D), throat pale yellow-green (4D); outside deep purplish pink (70C), each tepal yellowish white (155D) at base, margin brilliant yellow (13C), midribs brilliant yellow-green (142B) at top; spots strong purplish red (60D); papillae brilliant greenish yellow (6C); nectaries strong yellow-green (141D); pollen strong reddish orange (169A); stigma light purplish pink (63D). Fls 195 mm wide; tepals 115 × 58 mm, margins smooth, tips recurved. Lvs scattered, 107 × 21 mm. Stems 0.9-1.2 m, green with dark markings, and with c.5 fls. June.

(This cv. was omitted from the *ILR Twenty-third Supplement* and is published here.)

#### **VIII(b-c/d) 'Emír'**

cv. Parentage: 'Longida' × ('Gran Paradiso' × 'Puchanta')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside orange, tepal-tips and throat paler; outside pale orange; spots brown, oblong, moderately numerous, on half of each tepal; nectaries white-granulated; pollen orange-brown; stigma brown. Fls 140 mm wide; tepals 95 × 43 mm, very recurved, margins slightly wavy; pedicels 60 mm long, very thick, brown-dotted, with tertiary buds. Lvs scattered, 145 × 17 mm, mid green. Stems 1.05 m, thick, dark green with brown dots. Early July. Tetraploid.

#### **VII(a/b) 'Endios'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals deep/strong purplish pink (68A/B), margins strong reddish purple (70B), pale greenish yellow (2D) along midvein in basal two-thirds of each tepal, strong purplish red (63B) towards top, outer tepals strong purplish red (63B), margins (64C), pale greenish yellow (2D) along midvein for up to basal two-thirds of each tepal, throat yellowish white (155B); outside with inner tepals deep/strong purplish pink (68A/B), basal third of each tepal yellowish white (155D), margins moderate purplish red (64A), midribs strong purplish red (64B) on moderate yellow (162A) ground, strong yellow-green (144C) at top; outer tepals moderate/strong purplish red (63A/B), margins strong purplish red (64B), area next to midribs in basal sixth yellowish white (155D), midribs strong yellow-green (144B) at base changing into moderate purplish red (63A) on moderate yellow (162A) ground, strong yellow-green (144A) at top; spots absent; papillae brilliant greenish yellow (5A), strong purplish red (63B) and pale greenish yellow (2D); nectaries strong yellow-green (144A); pollen strong brown (172A); stigma moderate yellow-green (138B). Fls 240 mm wide; tepals 130 × 60-80 mm, margins ruffled, tips recurved. Lvs scattered, 160-210 × 40-65 mm, dark green. Stems 1.1-1.2 m, green with darker markings, and with c.4 fls. (Fl. colour becomes paler with age.)

#### **VII(a/b-c) 'Ensor'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer, B.V. (2004)

Inside with inner tepals dark/moderate purplish pink (186C/D), midveins deep purplish pink (68A), outer tepals very light purple (75C), dark purplish pink (186C) towards margins, light purple (75A) along midveins, throat yellowish white (155D); outside with inner tepals dark/moderate purplish pink (186C/D), very pale purple (69A) towards margins, yellowish white (155D) at base, midribs light yellow-green (144D) at base, changing upwards to moderate purplish red (186B), outer tepals dark/moderate purplish pink (186C/D), with a moderate purplish red (186B) stripe along midribs, base of midribs strong yellow-green (144B) changing into yellowish white (155D), dark greenish yellow (152D) at top; spots absent; papillae pale purplish pink

(65D); nectaries strong yellow-green (144C); pollen brownish orange (171B); stigma yellowish white (196D). Fls 250 mm wide, fragrant; tepals 155 × 60-85 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 190 × 57 mm. Stems 1.1-1.25 m, green with darker markings, with c.3 fls.

### **I(a/b-c) 'Equinox'**

cv. Parentage: Seedling AM 96-01 × seedling RS 93-18

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside strong purplish red (63A), throat strong yellow-green (144B); outside strong purplish red (63A), tepal-margins moderate purplish red (58A), midribs strong yellow-green (143C) at base and top, outer tepals paler; spots and papillae absent; nectaries moderate yellow-green (137C); pollen deep orange-yellow (163A); stigma yellowish white (155D), strong purplish red (60B) at edge. Fls 180 mm wide; tepals 105 × 40-50 mm, margins slightly ruffled, tips very slightly recurved. Lvs scattered, 135 × 23 mm. Stems 1.05-1.5 m, green with darker markings, and with c.6 fls.

### **VII(a/b-c) 'Escape'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals dark/moderate purplish pink (186C/D), brilliant yellow (10A) for up to two-thirds along midveins, bordered with pale yellowish green/greenish white (155A/C); outer tepals dark/moderate purplish pink (186C/D), brilliant/light yellow(10A/B) for up to two-thirds along midveins, pale yellowish green/greenish white (155A/C) at top, throat pale yellow-green (157C); outside with inner tepals very pale purple/moderate purplish pink (69A/186D), margins somewhat darker, base strong yellow-green (144B) and pale yellowish green (155A), outer tepals dark/moderate purplish pink (186C/D), margins greenish white/pale yellow-green (157D/155A), base with some strong yellow-green (144B), midribs pale yellowish green (155A), at top strong yellow-green (144B) with some strong yellow-green (144A); spots absent; papillae brilliant yellow (10A); nectaries strong yellow-green (144B); pollen moderate reddish brown (175A); stigma reddish grey (201C). Fls 210 mm wide; tepals 122-125 × 50-65 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 170 × 32 mm, green with lighter-coloured veins. Stems 1.1-1.2 m, green with darker margins, and with c.9 fls. (The fl. colour is not always solid.)

### **VII(b/b) 'Evelina'**

cv. Parentage unknown

R: Trior Lelies B.V., N: Trior Lelies B.V. (1994), REG: Lily Company B.V. (2004)

Inside with inner tepals yellowish white (155D) in basal half, towards margins and tips strong purplish pink (68B), brilliant greenish yellow (3B) along midveins towards top, outer tepals similar but with yellowish white (155D) margins and more purplish pink, throat yellowish white (155D); outside with inner tepals pale yellow-green (155A) in basal half, towards margins and tips dark purplish pink (186C), midribs strong yellow-green (144C) at base, dark purplish pink (186C) towards top, inner tepals dark purplish pink (186C), towards margins yellowish white (155D), midribs strong yellow-green (144C) at base with some dark purplish pink (186C), pale yellow-green (155A) towards top with a small strong yellow-green (144C) patch; spots absent; papillae brilliant greenish yellow (3B); nectaries strong yellow-green (143C) with narrow brilliant greenish yellow (3B) edge; pollen strong orange (170A); stigma dark purple (79A). Fls 150-240 mm wide, fragrant; tepals 140 × 42-65 mm, margins ruffled, tips recurved. Lvs scattered 160 × 28 mm. Stems 0.6-1 m, green, with c.9 fls. (Younger fls are a little more purplish pink.)

### VI(c/d) 'Ewunia'

cv. Parentage: 'Henrietina' × 'Brighton'

R: R. Widerski, N: R. Widerski, I: R. Widerski (2004), REG: R. Widerski (2004)

Inside pale orange; outside with pale green midribs; dense brown dashes on half of each tepal; nectaries green; pollen brown; stigma green. Fls 85 mm wide; tepals 100 × 40 mm, strongly recurved, slightly twisted; pedicels 10 mm long, brown and green, with tertiary buds. Lvs scattered, 210 × 40 mm, glossy mid green. Stems 1.3 m, thick, brown-dotted, with a long infl. Mid July.

### I(a/-) 'Fair'

This is the correct name for the cv. that was originally named, introduced and registered as 'Flair', and published in *ILR Fourth Supplement*. However, in 1986 Dutch PBR were granted under the name 'Fair', so this name must be accepted (ICNCP, 2004: Art. 24.4).

### VIII(b/b) 'Faros'

cv. Parentage: Seedling PH 94-68 × 'Aubade'

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals pale greenish yellow (10D) each with a large brilliant yellow (10A) spot, midveins brilliant yellow (10A) in basal half, apical half light yellow (10B), outer tepals pale greenish yellow (10D) with a light yellow (10B) flame, margins light yellow (10C), throat pale greenish yellow/yellowish white (10D/155D); outside with inner tepals pale yellow (11D), margins pale yellow (11C), midribs light yellow (11B), light green at base, flushed with moderate purplish red (184C) at top, outer tepals yellowish white/pale yellow (155D/11D), along midribs with a moderate purplish red (184C) streak, margins and midribs pale yellow (11C), base of midribs light yellow-green (145B), top strong yellow-green (144B); spots absent; papillae vivid yellow (13A); nectaries strong yellow-green (144A) with brilliant yellow (10A) margins; pollen greyish brown (166A); stigma dark purple (79A). Fls 250 mm wide; tepals 155-163 × 45-70 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 205 × 35 mm. Stems 0.8-1.25 m, light green with dark green stripes, and with c.4 fls. (An Orienpet (OT) lily.)

### VII(a-b/b) 'First Arrival'

cv. Parentage: Seedling 89-50-10 × seedling 90-058-4

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside and outside greenish white (155C); spots absent; papillae greenish white (155C); nectaries green; pollen yellow-brown; stigma purple. Fls 220 mm wide; tepals 145-152 × 50-58 mm, margins ruffled, tips recurved. Lvs scattered, 79-205 × 32-42 mm, dark green. Stems 0.8-1.25 m, green, with 3-11 fls. Midseason. (Nomenclatural standard: colour print supplied by registrant (WSY 0096396).)

### VII(a/-) 'First Love'

Published in *ILR Twenty-second Supplement* as an unacceptable name (ICNCP, 1995: Art 27.1 - name already used for a Cultivar Group). However, it has been found that this cv. was granted Dutch PBR in 1995 and therefore the name, although not registered, must now be accepted (ICNCP, 2004: Art. 24.4).

### I(a-b/b-c) 'Flevo Gold'

cv. Parentage unknown

REG: De Vries Veredelings B.V. (2005)

Inside vivid yellow (9A), midveins yellow-green at base; outside vivid yellow (9A/B);

spots absent; papillae few; nectaries green; pollen orange; stigma purple-red. Tepals short, margins slightly ruffled, tips slightly recurved. Stems 0.4 m, green with darker markings, with 2-7 fls. Early.

#### **VII(a/b) 'Florian'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside dark/moderate purplish pink (186C/D), with some yellowish white (155D) along and above nectaries and along tepal-margins; outside with inner tepals very pale purple (69A), margins narrow, yellowish white (155D), midribs light yellow-green (144D) at base and top, in between brilliant yellow-green (150C), yellowish white (155D) along midribs in basal half, outer tepals moderate purplish pink (186D), margins narrow, yellowish white (155D), midribs strong yellow-green (144B) at base and top, in between light yellow-green (145B); spots absent; papillae yellowish white (155D) and strong purplish pink (58C); nectaries strong yellow-green (144C); pollen vivid yellowish pink (28A); stigma pale green (189C). Fls 250 mm wide, slightly fragrant; tepals 130 × 50-71 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 179 × 44 mm. Stems 1.1-1.3 m, green with darker markings at base, with c.3 fls.

#### **VIII(a/b) 'Freya'**

cv. Parentage: Seedling BG 95-07 × seedling AG 98-21

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside brilliant yellow (7A), tepal-margins vivid yellow (9A), throat strong yellow-green (144B) with light yellow-green (150D) margin, becoming yellowish white (155D) with age; outside light yellow-green (2C), brilliant greenish yellow (4A) towards tepal-margins, margins brilliant yellow (7A), base with light yellow-green (144D) flush (older fls are yellowish white (155D) inside, with brilliant yellow (7A) tepal-margins and along midveins flushed with pale greenish yellow (2D)); outside yellowish white (155D), with brilliant yellow (7A) tepal-margins, and flushed with pale yellow-green (4D) along midribs); spots and papillae absent; nectaries pale greenish yellow (1D) (in older fls yellowish white (155D) with strong yellow-green (144B) edge); pollen moderate orange (167A/B); stigma deep greenish yellow (153A). Fls 220 mm wide; tepals 119 × 20 mm, margins smooth, tips recurved. Lvs scattered, 172 × 40 mm, dark green. Stems 1-1.25 m, dark green with darker markings, and with 3-6 fls.  
(A *longiflorum*/Asiatic (LA) hybrid.)

#### **VII(a-b/b) 'Gasparone'**

cv. Parentage unknown

REG: Hoogland B.V. (2005)

Inside greenish white (155C), basal part of each tepal along midvein brilliant yellow/light greenish yellow (8A/B); outside greenish white (155C), spots and papillae yellow; nectaries yellow-green; pollen orange-brown; stigma purple. Tepals medium to long, margins ruffled, tips strongly recurved. Lvs scattered. Stems 1.25 m, green, with 3-9 fls. Midseason to late.

#### **VII(a/b) 'Gerez'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside moderate purplish red/strong reddish purple (70A/B), basal half of each tepal with a large greenish white (155C) spot, throat greenish white (155C); outside with inner tepals light purple (75B), strong reddish purple (70B) towards margins, greenish

white (155C) at base, midribs pale yellow-green (4D), with purple flush towards top, outer tepals strong reddish purple (70B), moderate purplish red (70A) towards margins, greenish white (155C) at base, midribs pale yellow-green (4D), at base light yellow-green (145C) and at top light yellow-green (145B); spots absent; papillae at base few, greenish white (155C), others deep purplish red (71A); nectaries strong yellow-green (144A); pollen strong brown (172A); stigma dark purple/light purple (79B/77D). Fls 210 mm wide; tepals 150-160 × 57-77 mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs scattered, 152 × 37 mm, dark green. Stems 0.9-1.2 m, light green, with c.4 fls.

#### **VII(b/b-c) 'Giacondo'**

cv. Parentage: Seedling 93-14 × seedling RW 97-05

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals yellowish white (155D), with deep purplish pink (58D) streaks towards margins and tips, outer tepals similar but streaks stronger-coloured; outside with inner tepals yellowish white (155D), with deep purplish pink (58D) streaks, strong purplish red (59D) along midribs (less so with age), base and top of midribs strong yellow-green (144C) with light yellow-green (150D) in middle part, outer tepals similar but streaks more prominent, spots absent; papillae yellowish white (155D) and deep purplish pink (58D); nectaries moderate yellow-green (143D); filaments pale green; pollen strong brown (172A); stigma very pale green (192C). Fls 250 mm wide, fragrant; tepals 140 × 55-70 mm, margins gently ruffled, tips recurved. Lvs scattered, 160 × 38 mm, dark green. Stems 1.1-1.2 m, brownish with green, with 5-8 fls.

#### **VII(a-b/b) 'Gisborn'**

cv. Parentage: Seedling 89-14 × seedling RW 94-16

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside yellowish white (155D); outside with inner tepals yellowish white (155D), midribs strong yellow-green (144C) at base, changing towards top from light yellow-green (150D/154D), outer tepals yellowish white (155D), with midribs strong yellow-green (144C) at base changing towards top to light yellow-green/brilliant yellow-green (150D/149A), strong yellow-green (143B) at top, up to a fifth along midribs moderate red (185B); spots absent; papillae many, yellowish white (155D); nectaries strong yellow-green (144A); pollen strong brown (172A); stigma dark red (59A). Fls 260 mm wide; tepals 155 × 50-80 mm, margins ruffled, tips recurved. Lvs scattered, 160-210 × 40-60 mm, dark green. Stems 0.7-1.1 m, light green, with 2-5 fls.

#### **I(a/-) 'Golden Joy'**

cv. Parentage unknown

REG: Fa G.A. Verdegaaal & Zn Export B.V. (2005)

Inside vivid yellow (12A)/13A), at base strong orange (25A/B); outside light orange-yellow (25D), vivid yellow (9A) towards tepal-margins; spots few; papillae present; nectaries yellow-green; pollen brown; stigma dark purple. Tepals short, margins slightly ruffled, tips slightly recurved. Stems short. Early to midseason.

#### **VI(b-c/b) 'Gradient'**

cv. Parentage: (Palmer seedling 56221 × 'Crème de la Crème') × ((Palmer seedling 582812 × 'Gold Eagle') × 'Mimosa Star')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside white, centre and throat with deep yellow rays; outside with creamy midribs that are greenish at base; spots small, numerous, cinnamon, in lines over half of each tepal; pollen brown-apricot; stigma green. Fls 125 mm wide; tepals 115 × 30 mm, margin

slightly waved, tips recurved; pedicels 50 mm long, dotted with red-brown. Lvs 130 × 18 mm, dull pale green. Stems 1.15 m, rigid, freely dotted with red-brown. Mid July.

#### **I(a/-) 'Grand Cru'**

This is the correct name for 'Gran Cru' (published in *ILR Tenth Supplement*). This cv. was registered as 'Gran Cru', but it was granted Dutch PBR in 1988 under the name 'Grand Cru' and this latter name must be accepted (ICNCP, 2004: Art. 24.4).

#### **VII(a/b-c) 'Grasa'**

cv. Parentage: Seedling 93-80 × seedling RM 97-11

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside strong reddish purple (72B), midveins flushed with vivid reddish orange (32A) in basal half, top yellowish white (155D), throat yellowish white (155D); outside with inner tepals light purple (75A), midribs light yellow-green (145D) at base, changing into yellowish white (155D), outer tepals strong reddish purple (72B), at tips yellowish white (155D) at edge, midribs light yellow-green (144D) at base, changing into pale yellow-green (155A), light yellow-green (145D) at top; spots absent; papillae deep red (60A); nectaries strong yellow-green (144B) with light greenish yellow (1C) edge; pollen strong orange (168B); stigma yellowish grey (194C). Fls 200 mm wide, slightly fragrant; tepals 114 × 45-65 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered 130 × 30 mm. Stems 0.85-1.1 m, brownish with a little green, and with c.4 fls. (Young fls are more reddish purple.)

#### **VI(c/b-d) 'Gratulant'**

cv. Parentage: 'Favoritka' × unknown

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside creamy, centre and throat with golden yellow rays; outside with yellow shining through and with pale yellow midribs; spots dusty, very numerous, cinnamon, over half of each tepal; pollen apricot-brown; stigma green. Fls 135 mm wide, bowl- and turban-shaped; tepals 110 × 33 mm, very recurved, margins roughly waved, pedicels 105 mm long, thick, slightly dotted with red-brown. Lvs 145 × 19 mm, dull dark green. Stems 1.2 m, moderately thick, brown-dotted. Mid July.

#### **VI(b-c/b,d) 'Gurmán'**

cv. Parentage: ('Andrea' × 'Sommertraum') × ('Henrietina' × 'Mimosa Star')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside pale orange-apricot, tepal-tips paler; outside pale beige, midribs green; spots absent; nectaries olive-green; pollen pale brown; stigma violet-grey. Fls 135 mm wide, bowl- and turban-shaped; tepals 105 × 42 mm, moderately recurved, rigid; pedicels 75 mm long, dark brown-dotted. Lvs 110 × 9 mm, dull dark green. Stems 1.1 m, rigid, slightly black-dotted. Late July.

#### **I(c/d) 'Gwyneth Evans'**

cv. Parentage: Yellow-flowered, up-facing, speckled lily × unknown

R: I.R. Evans (1991), G: I.R. Evans (1993), N: I.R. Evans (2004), REG: I.R. Evans (2004)

Inside and outside light orange-yellow (19A); spots mid brown, moderately heavy, uniformly distributed; nectaries yellow-orange; pollen bright orange; stigma pale yellow. Fls 120-140 mm wide; tepals 60-70 × 30-40 mm, margins smooth, tips strongly recurved. Leaves scattered, 120-140 × 20-30 mm, dark green. Stems 1-2 m, dark green, with 10-50 fls.

### **I(b/c-d) 'Hadrián'**

cv. Parentage unknown

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside golden yellow, tepal-tips yellow, throat pale yellow-green; outside yellow with pale yellow in middle, midribs pale green-yellow; brushmarks mahogany-brown, spots large, mahogany-brown, over a third of each tepal; nectaries green, white-granulated; pollen ochre-brown; stigma brown. Fls 123 mm wide, out-facing to slightly pendent; tepals 85 × 35 mm, margins smooth, tips somewhat recurved; pedicels c.50 mm, brown-dotted. Lvs scattered, dull pale green, 100 × 12 mm. Stems 0.95 m, green, hairy above, with a mid-sized infl. Late June.

### **VIII(a/b) HALLEY**

Trade designation of 'Zanlahalley'.

### **I(a/b) 'Harry Wharton'**

cv. Parentage: 'Merryspring' × 'Sutherland Peach Melody'

R: D.J. Hay (1996), G: D.J. Hay (1999), N: D.J. Hay (2004), REG: D.J. Hay (2005)

Inside cream in basal part of tepals, pink in apical part, throat pink; outside pink, spots absent; nectaries pink; filaments reddish; pollen rust-orange; stigma dark pink. Fls 120 mm wide; tepals 74 × 38 mm, margins scarcely ruffled, tips very slightly recurved. Lvs scattered, 63 × 12 mm. Stems 0.5 m, brown, with c.5 fls. Mid July.

(Named after the raiser's great uncle. Nomenclatural standard: colour print supplied by registrant (WSY 0096411).)

### **VIII(b/b-c) 'Heartbreaker'**

cv. Parentage: Seedling PG 95-57 × seedling RH 97-07

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light greenish yellow (6D), light greenish yellow (8C) towards margins, brilliant greenish yellow (6B) along midveins, outer tepals with a streak of (53B) along midveins, this streak sometimes present on inner tepals. throat pale yellow (8D); outside with inner tepals pale yellow-green (4D) with a narrow light greenish yellow (8C) margin, midribs light yellow-green (144D) at base, outer tepals pale yellow-green (4D), midribs light yellow-green (144D) at base, light yellow-green (150D) with light yellow-green (143B) at top, with a strong red (53B) streak along midribs (stronger in older fls); spots absent; papillae brilliant yellow (8A); nectaries light yellow-green (144C); pollen dark reddish orange (175B); stigma deep reddish purple (77A). Fls 275 mm wide, slightly fragrant; tepals 150 × 55-86 mm, margins smooth to slightly ruffled, tips strongly recurved. Lvs scattered, 165 × 40 mm. Stems 0.9-1.1 m, green with darker markings, and with c.3 fls. (An Orienpet (OT) lily.)

### **I(a-b/b) 'Hillcrest Cynthia'**

cv. Parentage unknown

R: D. Putland (1990), G: Hillcrest Harmony Flowers (1992), N: Hillcrest Harmony Flowers (1995), I: Hillcrest Harmony Flowers (2005), REG: D. Putland of Hillcrest Harmony Flowers (2005)

Inside strong orange (25B) changing to pale orange-yellow (24D) in apical half of tepals; outside light orange-yellow (25D), midribs of inner tepals overlaid with green, midribs of outer tepals brilliant orange (29C); spots elongated, dark red (59A) on basal third of each tepal; nectaries green; pollen vivid reddish orange (32A); stigma strong red (51A). Fls 172 mm wide; tepals 90 × 45 mm, margins smooth, tips slightly recurved; pedicels with secondary buds. Lvs scattered, 125-170 × 13-26 mm, glossy. Stems 1.1-1.2 m, green with brown streaks, with some bulbils, and with c.25 fls. July.

(Named after the raiser's daughter.)

### **VI(a-b/c) 'Hold'**

cv. Parentage: Cream-flowered trumpet lily × yellow-green-flowered trumpet lily  
R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)  
Inside pale yellow, tepal-tips whitish, throat yellow; outside yellow, midribs brownish; spots absent; nectaries dark green; pollen ochre-apricot; stigma green. Fls 135 mm wide, shortly trumpet-shaped; tepals 130 × 55 mm, margins ruffled, tips recurved; pedicels 55 mm long, thick, freely brown-dotted. Lvs scattered, 90 × 14 mm, dark green, slightly shiny. Stems 1.15 m, greyish, slightly brown-dotted. Early July.

### **VIII(a-b/b) 'Honey Gold'**

cv. Parentage: Seedling E 854 × seedling OR 2  
R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)  
Inside brilliant yellow (9C), lighter towards tepal-tips, midveins yellow at base; outside light yellow (10C), darker towards tepal-margins, midribs green; spots and papillae absent; nectaries yellow-green; pollen brown-red; stigma green. Fls 190 mm wide; tepals medium-sized, margins ruffled, tips recurved. Lvs scattered. Stems 1-1.4 m, dark green, with 2-9 fls. Late.  
(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096395).)

### **VIII(a-b/b) 'Honeymoon'**

cv. Parentage: Seedling 164-5 × seedling 90-058-8  
R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)  
Inside light/pale yellow (11B/C), lighter towards tepal-margins and tips, brilliant yellow (13C) in basal half of midveins; outside pale yellow (11C); spots absent; papillae present; nectaries green; pollen brown-red; stigma purple. Fls 200 mm wide; tepals 155-162 × 42-67 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 185-205 × 29-40 mm, dark green. Stems 1-1.2 m, green, with 2-12 fls. Late.  
(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096407).)

### **VII(a/b-c) 'Honfleur'**

cv. Parentage unknown  
R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter & Den Haan Beheer B.V. (2004)  
Inside with inner tepals strong/vivid purplish red (68A/B), light purple (75A) towards margins, midveins vivid purplish red (57B), outer tepals strong/vivid purplish red (68A/B), basal half very pale purple (73D), midveins vivid purplish red (57B), throat yellowish white (155D); outside with inner tepals strong reddish purple/deep purplish pink (70B/C), yellowish white (155D) at base, midribs strong yellow-green (144A) at base, changing to deep purplish pink (54B), towards top pale greenish yellow (2D) with light yellow-green (144D) at top, outer tepals light purple (75B), margins light purple (75A), midribs strong yellow-green (144A) at base changing into yellowish white (155D), towards top strong yellow-green (144A); spots absent; papillae strong purplish pink (63C) and vivid red (57A); nectaries strong yellow-green (144C); pollen deep orange-yellow (163A); stigma very light purple (85C). Fls 270 mm wide, slightly fragrant; tepals 155 × 61-86 mm, margins ruffled, tips recurved. Lvs scattered, 175 × 58 mm. Stems 1.15-1.2 m, green with darker markings, and with c.3 fls.

### **I(a/c) 'Ice Berry'**

cv. Parentage: White-flowered seedling × 'Vivaldi'  
R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2000), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside light purplish pink (63D), each tepal with a yellowish white (155D) flame and a narrow strong purplish pink (63C) margin; outside pale purplish pink (65C), tepal-margins narrow, strong purplish pink (67D); spots few, dark red (59A); papillae light purplish pink (63D); nectaries strong yellow-green (144B); pollen strong brown (172A); stigma moderate orange (170C). Fls 195 mm wide; tepals 105 × 33-52 mm, margins slightly ruffled, tips recurved. Lvs scattered, 130 × 16 mm. Stems 1.01 m, green with darker markings, and with c.6 fls.

#### **VIII(a/b) 'Ice Diamond'**

cv. Parentage: (Seedling 3217.rs × 'Esra') × seedling LA 93055.5T

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside yellowish white (155D), throat light yellow-green (145D); outside yellowish white (155B), midribs with some moderate yellow-green (143D) at base and top; spots and papillae absent; nectaries strong yellow-green (143C); pollen brownish orange (164A); stigma greenish white (157D). Fls 175-190 mm wide; tepals 132 × 45-59 mm, margins smooth, tips slightly recurved. Lvs scattered, 150 × 15 mm. Stems 1.15 m, green with some darker markings at base, and with 2-10 fls.

(*A longiflorum*/Asiatic (LA) hybrid.)

#### **VIII(a-b/b) 'Icepearl'**

cv. Parentage: 'Monte Rosa' × unknown

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside pale yellow-green (4D), becoming whiter with age; outside pale yellow-green (4D), midribs green; spots few; papillae absent; nectaries yellow-green; pollen brown; stigma light yellow-green. Tepals short to medium-sized, margins scarcely ruffled, tips slightly recurved. Lvs scattered, dark green. Stems 1-1.4 m, green with darker markings, and with 2-8 fls. Midseason.

(*A longiflorum*/Asiatic (LA) hybrid. Nomenclatural standard: colour print supplied by registrant (WSY 0096410).)

#### **I(a-b/b-c) 'Idylle'**

cv. Parentage: Seedling AW 94-09 × seedling W 92-01

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside greenish white (155C), throat greenish white (155C) with small strong/light yellow-green (144C/145D) stripes; outside with inner tepals greenish white (155C), along midribs strong yellow-green/moderate red (144C/181A), outer tepals light/strong yellow-green (145C/144B), margins greenish white (155C), striped or flushed with moderate red (181A/B) along midribs; spots and papillae absent; pollen moderate brown (165A); stigma vivid red (45B). Fls 170 mm wide; tepals 90-97 × 40-54 mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs scattered, 129 × 25 mm, dark green. Stems 1.05-1.3 m, dark green with darker markings, and with c.7 fls.

#### **I(a-b/b-c) 'Jaap Aap'**

cv. Parentage: Seedling AG 94-01 × seedling AG 95-12

R: Gebr. Vletter & Den Haan, REG: Vletter & Den Haan Beheer B.V. (2004)

Inside vivid yellow (14A), lighter towards base; outside vivid yellow (14A); spots absent; papillae few, small; nectaries green; pollen reddish brown; stigma yellow with small red spots. Tepals short, tips slightly recurved. Lvs scattered. Stems 1.2 m, green. Early to midseason.

### **VII(b/b) 'Jacobi'**

cv. Parentage: Seedling 93-13 × seedling RM 97-22

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside strong purplish pink (63B), tepal-margins moderate purplish red (70A), brilliant yellow/light greenish yellow (8A/B) around basal third of nectary, strong purplish pink (63A) along midveins, throat yellowish white (155D); outside with inner tepals deep purplish pink (70C), margins strong reddish purple (70B), yellowish white (155D) along midribs in basal third, midribs strong yellow-green (144A) at base, changing into brilliant yellow-green (150C), brilliant yellow-green (150A) at top, outer tepals strong purplish pink (63B/C), margins strong purplish red (63A), base yellowish white (155D), light purplish pink (63D) along midribs in basal half, midribs strong yellow-green (144A) at base, changing into brilliant yellow-green (150C), brilliant yellow-green (150A) at top; spots and papillae strong purplish pink (60B); nectaries strong yellow-green (144A); pollen strong brown (172C); stigma dark red (187A). Fls 190 mm wide; tepals 160 × 70-100 mm, margins ruffled, tips recurved. Lvs scattered, 140-180 × 35-45 mm, dark green. Stems 1.1-1.2 m, green with darker markings, and with c.3 fls.

### **VI(c/b-d) 'Jakobín'**

cv. Parentage: (Silver Sunburst Group × 'Gold Eagle') × (((*henryi* × unknown) × 'Gold Eagle') × unknown)

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside white, centre and throat yellow-green; outside with yellow midribs, green and brownish at base; small cinnamon spots on two-thirds of each tepal; nectaries yellow-green; pollen apricot-brown; stigma green. Fls 145 mm wide, bowl- and turban-shaped; tepals 110 × 40 mm, margins slightly wavy, tips recurved. Lvs 100 × 13 mm, yellowish green, sparse. Stems 1.15 m, brown-dotted. Mid July.

### **I(a-b/b) 'Jarmilka'**

cv. Parentage: ('Pinegos Fire' × 'Tetra Nutmegger') × white-flowered tetraploid

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside creamy tinged with beige, tepal-tips white; outside with greenish and brownish midribs; spots numerous, dark crimson-brown, large, oval, on half of each tepal; nectaries green, white-granulated; pollen chocolate-brown; stigma beige. Fls 165 mm wide; tepals thick, 100 × 47 mm, slightly recurved, the outer more so; pedicels 70 mm, thick, dotted with dark brown. Lvs scattered, 135 × 21 mm, dull dark green, dense. Stems 0.75 m, dotted with dark brown. Early July. Tetraploid.

### **I(a/b) 'Just Jazz'**

cv. Parentage unknown

R: R.H. Bazett (1992), G: R.H. Bazett (1995), N: R.H. Bazett (2002), I: Valley K Greenhouses (2005), REG: R.H. Bazett (2005)

Inside mid pink, apical half of tepals lighter, throat red with yellow undertone; outside similar, midribs grey-green; spots and papillae absent; nectaries yellow edged with dark grey; pollen red-brown; style pinkish; stigma pale green. Fls 170 mm wide; tepals 95 × 45 mm, margins slightly ruffled, tips not recurved. Lvs scattered, 150 × 17 mm. Stems 1.25 m, dark green, with c.16 fls. June to July. Tetraploid.  
(Nomenclatural standard: colour print supplied by registrant (WSY 0096398).)

### **I(b/b-c) 'Kampa'**

cv. Parentage: ('Dawn Star' × 'Parfait') × ('Heloma' × ('Sioux' × 'Wattle Bird'))

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside orange, with small brownish flame over nectaries; outside with green midribs;

spots dark pink-brown, medium-sized, numerous, on three-fifths of each tepal; nectaries finely white-granulated; pollen orange-brown; stigma crimson-brown. Fls 120 mm wide; tepals 75 × 40 mm, thick and rigid, moderately recurved; pedicels 80 mm long, thick, brown-dotted. Lvs scattered, 120 × 11 mm, dull dark green. Stems 0.9 m, finely brown-dotted. Late June.

#### **VII(a/b) 'Kanton'**

cv. Parentage: Seedling RM 94-20 × seedling RM 97-02

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside vivid purplish red (57D), tepal-margins and tips yellowish white (155B), vivid purplish red (57C) along midveins, midveins strong purplish red (58B) in basal half, apical half strong purplish red (63A), throat yellowish white (155B); outside with outer tepals strong purplish red (63B), towards margins strong purplish pink (63C), margins yellowish white (155B), midribs strong yellow-green (144C) at base, changing into light yellow-green (154D) with strong purplish red (64B), brilliant/strong yellow-green (149A/143B) towards top, inner tepals paler; spots few, deep red (60A); papillae deep red (60A); nectaries strong yellow-green (144C); pollen strong brown (172C); stigma light purple (75A). Fls 230 mm wide; tepals 130 × 45-55 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 120-130 × 30-40 mm, dark green. Stems 1-1.2 m, light green with some darker marking, and with c.5 fls.

#### **VI(b-c/a) 'Karkulka'**

cv. Parentage: (Unknown × 'Schokolade') × (unknown × 'Toni')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside creamy, tepal-tips slightly pink-veined, throat with lemon-yellow rays; outside pale yellow, tepal-margins pink-veined, midribs green-yellow with pink veining; spots absent; nectaries dark green; pollen apricot; stigma whitish. Fls 130 mm wide and 80 mm long, shortly trumpet-shaped; tepals 130 × 56 mm, very recurved with twisted tips, margins slightly wavy; pedicels 65 mm, green. Lvs scattered, 130 × 9 mm, dull dark green. Stems 1.15 m, green. Early July.

#### **VIII(a/b) 'Karnak'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan., N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside strong purplish pink (68B), strong purplish red (63A) along margins and midveins, throat brilliant yellow-green (142B), yellowish white (155D) towards edge; outside strong purplish pink (63C), tepal-margins strong purplish red (63B); spots few, strong purplish red (63A); papillae strong purplish red (63A); nectaries strong yellow-green (143A); pollen strong brown (172C); stigma light greenish yellow (3C). Fls 225 mm wide; tepals 135 × 48-70 mm, margins smooth, tips recurved. Lvs scattered, 165 × 23 mm. Stems 1.1-1.5 m, green with darker markings, and with c.4 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **I(a-b/b-c) 'Kastelán'**

cv. Parentage: (('Sioux' × 'Wattle Bird') × ('Rusalka' × unknown)) × (unknown × ('Bélásek' × 'Mánes'))

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside very pale golden yellow, tepal-tips whitish, centre of tepals stained with crimson-brown, throat pale yellow; outside tinged with old rose-pink (especially on outer tepals); spots dark brown, oval, medium-sized, on basal half of each tepal (more numerous on inner tepals); nectaries white- and yellow-granulated; pollen brown; stigma crimson-brown. Fls 135 mm wide; tepals 80 × 37 mm, slightly curved; pedicels 70 mm long,

dark brown-dotted. Lvs scattered, 125 × 14 mm, dull light green. Stems 0.85 m, thick, dark brown-dotted.

#### **VI(b/b) 'Katowice'**

cv. Parentage: Unknown × ('Kremona' × 'Melone')

R: R. Widerski, N: R. Widerski, I: R. Widerski (2004), REG: R. Widerski (2004)

Inside lemon-yellow; outside with pale green midribs; spots absent; pollen yellow-brown; stigma green-yellow. Fls 140 mm wide; tepals 130 × 55 mm, outer tepals recurved, inner less so, twisted, tough; pedicels 120 mm long, thick, green. Lvs scattered, 110 × 10 mm, dull mid green. Stems 1.5 m, thick, green, with medium-sized infl. Mid July.

#### **I(a/b) 'Keiko'**

cv. Parentage: (unknown × ('Mohykán' × 'Juno')) × ('Odysseus' × 'Mánes')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside pale apricot, tepal-tips paler, each tepal with a brown-crimson flame over nectary; outside with pinkish brown midribs; spots absent; nectaries white-granulated; pollen brown; stigma pale brownish old rose-pink. Fls 123 mm wide; tepals 73 × 40 mm, scarcely curved; pedicels 75 mm long, with secondary buds. Lvs scattered, 130 × 18 mm, dark green, slightly shiny. Stems 1 m, thick, green. Late June.

#### **VIII(a/b) 'Kentucky'**

cv. Parentage: ('Tango' × unknown) × ('Sterling Star' × 'Connecticut King')

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2001), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside light orange-yellow (24C), throat pale yellow-green (4D); outside with outer tepals pale orange-yellow (23D), margins light orange (26C) with dark red (187B) spots, along midrib strong yellow-green (143C) changing upwards into light yellow-green (145C), inner tepals with broader margins; spots dark red (187B) on basal third of each tepal; papillae dark red (187B); nectaries moderate yellow-green (139C); pollen deep orange-yellow (163A); stigma vivid yellowish pink (30C). Fls 200-230 mm wide, fragrant; tepals 133 × 37-55 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 190 × 19 mm. Stems 1 m, green with darker markings at base, and with c.9 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **I(a/b) 'Kissime'**

cv. Parentage unknown

R: R.H. Bazett (1998), G: R.H. Bazett (2000), N: R.H. Bazett (2005), I: Valley K

Greenhouses (2005), REG: R.H. Bazett (2005)

Inside ivory-white, throat ivory-white to grey-green; outside with inner tepals ivory-white, outer tepals ivory white, midribs grey-green; spots and papillae absent; nectaries grey-green; pollen dark orange; stigma light brown. Fls 160 mm wide; tepals 90 × 45 mm, margins smooth, tips slightly recurved. Leaves scattered, 130 × 15 mm, dark green. Stems 1.05 m, dark green, with c.12 fls. June to July.

#### **I(a/b) 'Korsikán'**

cv. Parentage: ('Mirage' × 'Impact') × ('Target' × 'Windahloo')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside pale brownish red-orange, each tepal with pale yellow margin and pale tip; outside pale orange; spots red-brown, medium-large, on half of each tepal, brushmarks large, red-mahogany, with longer lines; feathered papillae in throat; nectaries white-granulated; pollen ochre-brown; stigma brownish pink. Fls 125 mm wide; tepals 85 × 40 mm, margins incised, tips slightly curved; pedicels 65 mm long, green. Lvs scattered, 130 × 13 mm, dull dark green. Stems 0.85 m, thick, green. Early June.

### **VI(b-c/b) 'Kvarteto'**

cv. Parentage unknown

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside white, outer tepals pink-veined, centre of fl. with narrow pink rays, throat lemon-yellow; outside pink; spots absent; nectaries green; pollen red-brown; stigma brown. Fls 130 mm wide; tepals thick, 105 × 42 mm, somewhat recurved; pedicels 75 mm, brown-dotted. Lvs scattered, 100 × 11 mm, dull deep green, dense. Stems 1.15 m, dotted red-brown. Early July.

### **I(a/b) 'Lady Willemse'**

cv. Parentage unknown

R: R. Griesbach, N: Lily Company B.V. (2004), REG: Lily Company B.V. (2004)

Inside vivid reddish orange (30A), each tepal with a large, dark greyish purple/dark red (202A/187A) flame; outside strong reddish orange (31A), tepal-margins vivid reddish orange (32A); spots few, dark greyish purple/dark red (202A/187A); papillae vivid reddish orange (30A); nectaries strong yellow-green (143A); pollen dark reddish orange (173A); stigma strong reddish orange (31A). Fls 120 mm wide; tepals 55 × 16-25 mm, margins smooth, tips slightly recurved. Lvs scattered, 90 × 20 mm, dark green. Stems 0.65-0.7 m, green with darker markings, and with c.5 fls.

### **VII(a/b-c) 'Lake Carey'**

cv. Parentage: Red-flowered seedling × ('Rosalie' × 'Star Gazer')

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V., REG: Bischoff Tulleken Lelies B.V. (2004)

Inside deep/strong purplish red (61A/67A), strong purplish red (60D) towards tepal-margins, apical half of tepals with a narrow yellowish white (155D) margin, strong red (53B) along midveins towards top, throat yellowish white (155D); outside with outer tepals strong purplish red (64C), apical half with moderate purplish pink (65A) margin, base yellowish white (155D), midribs strong yellow-green (144C) at base, changing into yellowish white (155D) with moderate purplish pink (65A), at top with some moderate yellow-green (146D), inner tepals similar but with narrow yellowish white (155D) margins; spots absent; papillae few, (59B); nectaries strong yellow-green (144C) with narrow brilliant greenish yellow (1B) edge; pollen moderate reddish orange (171A); stigma dark purple (79A). Fls 215-250 mm wide, fragrant; tepals 123 × 50-65 mm, margins slightly ruffled, tips recurved. Lvs scattered, 115 × 40 mm. Stems 0.8-1.2 m, green, darker at base, with c.4 fls.

### **VII(a/b) 'Lake Huron'**

cv. Parentage: (Unnamed seedling × 'Hit Parade') × ('Allegra' × 'Star Gazer')

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside moderate purplish pink (186D), basal half of each tepal with light greenish yellow (3C) along midveins, pale yellow (19C) towards top, throat yellowish white (155D); outside moderate purplish pink (186D), dark purplish pink (186C) towards tepal-margins, base yellowish white (155D), midribs strong yellow-green (144C) at base, changing into light yellow-green (145D), strong yellow (153D) towards top; spots dark purplish pink (186C); papillae light greenish yellow (3C) and yellowish white (155D); nectaries strong yellow-green (144C); pollen moderate reddish orange (178C); stigma pale green (189C). Fls 255 mm wide, fragrant; tepals 150 × 45-63 mm, margins smooth to ruffled, tips recurved. Lvs scattered, 140 × 35 mm. Stems 0.9-1.25 m, green with darker markings, and with c.6 fls.

### VII(b/b) 'Lake Volta'

cv. Parentage: 'Aubade' × unknown

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside yellowish white (155D), each tepal with a light greenish yellow (6D) flame, brilliant greenish yellow (6C) along basal half of midveins; outside yellowish white (155D), midribs strong yellow-green (144C) at base, above base with a streak of strong purplish red (63A), brilliant greenish yellow (4A) towards top; spots absent; papillae brilliant greenish yellow (6A); nectaries strong yellow-green (144A); pollen dark reddish orange (172B); stigma dark red (187A). Fls 240 mm wide; tepals 160 × 50-70 mm, margins slightly ruffled, tips recurved. Lvs scattered, 140-170 × 40-50 mm, dark green. Stems 0.8-1.15m, light green with some darker markings, and with c.6 fls.

### I(a/b) 'Lapkritis'

cv. Parentage: 'Rotāja' × seedling 420-S-88

R: Ā. Zorgevics (1985), G: Ā. Zorgevics (1989), N: Ā. Zorgevics (2005), I: University of Latvia Botanical Garden (2005), REG: Ā. Zorgevics (2005)

Inside vivid reddish orange (33A), throat light red; outside vivid reddish orange (33A); spots and papillae absent; nectaries orange; pollen dark brown; stigma pale. Fls 120 mm wide; tepals 80 × 40 mm, margins very slightly ruffled, tips slightly recurved. Lvs scattered, 70 × 10 mm. Stems 1.1m, green, with c.9 fls. Early July.

### I(a/-) LAVA

Trade designation of 'Brandaris'.

(In *ILR Twenty-second Supplement* LAVA was erroneously given as a trade designation of 'Apollo'.)

### VIII(a-b/b-c) 'Lavon'

cv. Parentage: Seedling 819-1 × seedling 90-033-1

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside light greenish yellow (5D), each tepals with a deep pink (50B) flame from top of nectary, midveins purple-red at base; outside light greenish yellow (5D) with some dark pink-red; spots and papillae present; nectaries green, yellow-green at top; pollen brown-red; stigma green with purple star. Fls 180 mm wide; tepals 137-144 × 40-60 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 160-195 × 30-40 mm, dark green. Stems 0.9-1.1 m, green with darker spots and stripes, and with 1-9 fls. Very late.

(An Orienpet (OT) lily.)

### VIII(a-b/b) 'Leda'

cv. Parentage: Seedling E 164-1 × seedling 90-045-8

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside pale yellow (20D), lighter towards tepal-tips, basal half of each tepal brilliant yellow (14C) in centre, midveins brilliant yellow (14C) at base; outside pale orange-yellow (19D); spots absent; papillae present; nectaries green; filaments pale green; pollen dark yellow-orange; style green; stigma purple. Fls 200 mm wide; tepals 155-160 × 36-63 mm, margins slightly to strongly ruffled, tips strongly recurved. Lvs scattered, 139-224 × 24-39 mm, dark green. Stems 1-1.2 m, green, with 1-9 fls. Late.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096397).)

### VIII(b-c/b-d) Lemgo Group

grp Parentage: Yellow-flowered Trumpet × ('Black Beauty' × (*auratum* × *henryi*))

R: H. Dierssen (1997), N: H. Dierssen, I: H. Dierssen, REG: H. Dierssen (2004)

Inside deep crimson, tepal-margins and tips pale beige-apricot; outside beige, midribs green with mahogany speckles; spots dash-like, raised, dark crimson, on a third of each tepal; nectaries deep yellow-green; pollen chocolate-brown; stigma violet-red. Fls 155 mm wide; tepals 140 × 42 mm, strongly reflexed, margins ruffled; pedicels 120 mm, strongly spotted. Lvs scattered, 85 × 18 mm, slightly glossy dark green. Stems 1.15 m, moderately thick, strongly brown-speckled. Mid July.  
(Orienpet (OT) lilies.)

### VII(a/b) 'Lepanto'

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside light purplish pink/very pale purple (73C/D), tepal-margins light purple (75A), throat greenish white (155C); outside with inner tepals very pale purple/light purplish pink (69C/73C), margins vivid reddish purple (74A), midribs strong yellow-green (144B) at base, changing into deep purplish pink (73A), outer tepals dark/moderate purplish pink (186C/D), dark purplish pink (186C) along midribs, midribs strong yellow-green (144B) at base and top, in between moderate orange (170C); spots absent; papillae white and very pale purple (73D); nectaries strong yellow-green (144B); pollen strong reddish orange (169A); stigma moderate/very pale purple/(79D/84D). Fls 275 mm wide; tepals 165-172 × 54-76 mm, margins ruffled, tips recurved. Lvs scattered 205 × 36 mm, dark green. Stems 1.2 m, light green with darker markings, and with c.4 fls.

### VII(a/b) 'Lido'

cv. Parentage: Seedling 93-40 × seedling RW 97-01

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside moderate/light purplish pink (62B/C), strong purplish pink (62A) towards tepal-tips, throat yellowish white (155B); outside light/pale purplish pink (65B/C), midribs strong yellow-green (144B) at base, light yellow-green (154D) towards top; spots absent; papillae strong purplish red (63B) and yellowish white (155B); nectaries strong yellow-green (144B); pollen dark reddish orange (175C); stigma brilliant/light yellow-green (142B/C). Fls 250 mm wide; tepals 150 × 55-80 mm, margins ruffled, tips recurved. Lvs scattered, 130-155 × 30-40 mm, dark green. Stems 0.8-1.2 m, light green with darker markings, and with c.3 fls.

### II(c/d) 'Lois Hole'

cv. Parentage unknown

R: (1994), G: M. Joslin (2001), N: M. Joslin (2004), I: Estate Perennials (2004), REG: M. Joslin (2004)

Inside vivid yellow (9A); outside white; spots and papillae absent; nectaries yellow; pollen orange; stigma yellow. Fls 85 mm wide; tepals 35 × 10 mm, margins smooth, tips slightly recurved. Lvs whorled, 40 × 9 mm, mid green. Stems 1-1.2 m, mid green, with 13-15 fls. Early July.

(Named for Lois Hole, Lieutenant Governor of Alberta, Canada.)

### VII(a/b) 'Lutine'

cv. Parentage: Seedling 92-12 × seedling RW 96-05

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals moderate purplish red (70A), strong reddish purple (72B)

towards margins, midveins strong purplish red (71B), yellowish white (155D) at top, outer tepals strong purplish red (71B), moderate purplish red (71A) towards margins, midveins strong purplish red (64B), yellowish white (155D) at top, throat yellowish white (155D); outer tepals with inner tepals strong reddish purple (70B), moderate purplish red (71A) towards margins, base with some yellowish white (155D), midribs strong yellow-green (144C) at base, moderate purplish pink (70D) towards top, yellowish white (155D) at top, outer tepals moderate purplish red (70A), moderate purplish red (71A) towards margins, base with some yellowish white (155D), midribs strong yellow-green (143C) at base and top, in between light yellow-green (154D) with moderate purplish red (70A); spots absent; papillae strong purplish red (60B); nectaries moderate yellow-green (139D), yellowish white (155D) along edges; pollen moderate reddish brown (166B); stigma moderate brown (200D). Fls 260 mm wide, fragrant; tepals 170 × 55-80 mm, margins ruffled, tips recurved. Lvs scattered, 200 × 53 mm. Stems 1.2-1.25 m, green with some darker markings, and with c.3 fls.

### **I(a-b/b) 'Lužánky'**

cv. Parentage: (('Quinella' × unknown) × ('Juno' × 'Honey Creme')) × ('Polárka' × unknown)

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside pale pink, with gold shining though around nectaries, throat crimson-pink; outside pink with beige midribs; spots present; nectaries pink-granulated; pollen ochre; stigma greenish. Fls 130 mm wide; tepals 73 × 36 mm, margins slightly ruffled, tips slightly curved; pedicels 70 mm long, dotted. Lvs scattered, 75 × 14 mm, dull mid green. Stems 0.85 m, thick, freely finely brown-dotted. Mid June.

### **VI(b/a) 'Maestro'**

cv. Parentage: 'Spītniece' × seedling L-11-35

R: Ā. Zorgevics (1987), G: Ā. Zorgevics (1993), N: Ā. Zorgevics (2005), I: University of Latvia Botanical Garden (2005)

Inside brilliant yellow (9C), throat brilliant yellow-green (149B); outside with brownish shading; spots and papillae absent; nectaries whitish; pollen dark brown; stigma grey. Fls 220 mm wide, strongly scented; tepals 150 × 30 (outer) and 60 (inner) mm, margins very slightly ruffled, tips recurved. Lvs scattered, 150 × 15 mm. Stems 1.7 m, green with brown, and with c.9 fls. Bulbs purple. Late July to early Aug.

(Unacceptable name, ICNCP, 2004: Art. 27.1 - name already used for another lily, so registration refused.)

### **VII(a/b) 'Magic Pearl'**

cv. Parentage: 'Medusa' × 'Solaia'

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside strong purplish red (63B), tepal-margins yellowish white (155B), moderate purplish red (63A) along midveins, throat yellowish white (155B); outside with outer tepals strong purplish red (60D), margins yellowish white (155B), midribs strong yellow-green (143B) at base, up to a half light yellow-green (150D), strong yellow-green (142A) at top, inner tepals deep/strong purplish pink (68A/B), margins yellowish white (155B), midribs strong yellow-green (142A) at base, towards top light yellow-green (150D), changing into brilliant yellow-green (150A) with some deep purplish pink (68A); spots strong purplish red (61B); papillae deep purplish red (61A) and at base yellowish white (155B); nectaries strong yellow-green (144A); pollen dark reddish orange (175C); stigma dark red (187A). Fls 240 mm wide, slightly fragrant; tepals 140 × 40-70 mm, margins slightly ruffled, tips recurved. Lvs scattered, 130-160 × 18-28 mm, dark green. Stems 0.8-1.3 m, green with some darker markings, and with 5-7 fls.

### VIII(a-b/b) 'Malibu'

cv. Parentage: Seedling E 905-5 × seedling OR 2

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)  
Inside brilliant yellow (12B/13C), lighter towards tepal-tips; outside light yellow (11B); spots and papillae absent; nectaries green; pollen orange-brown; stigma green. Fls 180 mm wide; tepals 130-135 × 35-56 mm, margins slightly ruffled, tips recurved. Lvs scattered, 195-230 × 45-84 mm, dark green. Stems 1.45-1.65 m, green, with 2-12 fls. Late.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096400).)

### VII(b/b) 'Mantegna'

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside yellowish white (155D); outside with outer tepals yellowish white (155D), midribs light yellow-green (145B) at base, light yellow-green (150D) at top with some light yellow-green (144D), along midribs flushed with strong purplish red (58C) - which disappears with age, inner tepals similar but midribs light yellow-green (145D) at base changing into pale yellow-green (155A); spots absent; papillae yellowish white (155D); nectaries light yellow-green (144D); pollen strong orange (24A); stigma light greenish grey (190C). Fls 185 mm wide, fragrant; tepals 121 × 45-78 mm, margins smooth to ruffled, tips recurved. Lvs scattered, 140 × 35 mm. Stems 0.85-0.9 m, green with darker markings, and with c.4 fls.

### I(a-b/b) 'Marcelka'

cv. Parentage: 'Mafían' × white-flowered tetraploid

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside pink with salmon tinge, throat whitish, area around nectaries pale apricot; outside pink, midribs brown; nectaries white-granulated. Fls 155 mm wide; tepals thick, 95 × 47 mm, somewhat recurved.; pedicels 75 mm, very thick, heavily dotted with dark brown. Lvs scattered, 135 × 20 mm, glossy mid green. Stems 1.15 m, dark brown, very thick. Triploid.

### I(a-b/b) 'Maruška'

cv. Parentage: (((('Tetra Enchantment' × 'Tetra Tabasco') × 'Tetra Connecticut Yankee') × 'Avignon') × ('Gran Paradiso' × 'Tetra Aristo'))

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside bright orange-red, throat with yellow shining through; outside orange with crimson mottling, midribs brownish; spots and papillae present; nectaries white-granulated; pollen chocolate-brown; stigma dark crimson-brown. Fls 130 mm wide; tepals 85 × 47 mm, scarcely curved; pedicels 65 mm long, abundantly dotted dark brown, with secondary buds. Lvs scattered, 120 × 18 mm, dull mid green. Stems 0.85 m, very thick, dark brown. Late June. Tetraploid.

### VII(a-b/b-c) 'Maryland'

cv. Parentage: Seedling 92-96 × seedling 92-36

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals greenish white (155C), midveins light/pale yellow (10C/11C) in basal half, moderate purplish pink (186D) at top, outer tepals greenish white (155C), midveins with light yellow (10B/C) along basal half, apical half stronger moderate purplish pink (186D) than on inner tepals; outside with inner tepals greenish white (155C), light yellow-green (145B) at base, greenish white (157D) along midribs,

light greenish yellow (4C) towards top, outer tepals yellowish white (155B), light yellow-green (145B) at base, light yellow-green (150D) along midribs, strong yellow-green (144C) with strong yellow-green (144A) at top, next to midribs dark purplish pink (186C); spots absent; papillae greenish white/pale greenish yellow (155C/10D); nectaries stronger yellow-green (144B); pollen moderate reddish brown (175A); stigma dark purple (79A). Fls 225 mm wide, fragrant; tepals 115-120 × 42-62 mm, margins ruffled, tips recurved. Lvs scattered, 135 × 30 mm, dark green with lighter veins. Stems 0.9-1.2 m, light green, darker in lower part, with c.9 fls.

### **II(c/d) 'Mary Liss'**

cv. Parentage: Backhouse Group × unknown

G: M. Liss (pre 1990), N: D. Dale (2002), I: Estate Perennials (2005), REG: M. Joslin (2005)

Inside vivid yellow (14A), overlaid with purplish pink, tepals with narrow yellow margins, tips with tiny dark red (59A) speckles, throat light yellow (15D) with dark red (59A) lines; outside white with pale purplish pink (65C) highlights; spots large, dark maroon, surrounded by yellow, mainly in central part of each tepal; nectaries light yellow (14D); pollen dark reddish orange (172B); stigma pale cream. Fls 80 mm wide, slightly scented; tepals 45 × 14 mm, margins smooth. Lvs whorled, 200 × 40 mm. Stems 1.75 m, green mottled with black, with 25-35 fls. Early July.

(Originated as a seedling in the garden of the late Mary Liss. Nomenclatural standard: colour print supplied by registrant (WSY 0096428).)

### **VII(a/b) 'Molfetta'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside deep purplish red (59B/187D), throat greenish white (155C); outside with outer tepals deep purplish red (59C/187D), with some strong/light yellow-green (144C/145D) at base, light yellow-green (150D) along midribs, strong yellow-green (144A) at top, inner tepals moderate/strong purplish red (64A/B), at base on greenish white (155C) ground, midribs strong yellow-green (144C) at base, light yellow-green/moderate to strong purplish red (150D/64A/B) at top; spots absent; papillae greenish white (155C) and moderate/strong purplish red (59C/D); nectaries strong yellow-green (144B); pollen strong orange (25A); stigma moderate purple (79D). Fls 230 mm wide; tepals 133-145 × 34-74 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 159 × 34 mm. Stems 0.95-1.1 m, light green, with c.3 fls.

(The fl. colour darkens with age.)

### **I(a/b) 'Moon Jewel'**

cv. Parentage: ('Michelle' (tetraploid form?) × 'Nove Cento') × 'Stella' (tetraploid form?)

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside vivid yellow (9A), throat pale greenish yellow (1D); outside light greenish yellow (7D), brilliant yellow (7A) towards tepal-margins, midribs strong yellow-green (143C) at base and top; spots absent; papillae vivid yellow (9A); nectaries strong yellow-green (143B); pollen dark reddish orange (175B); stigma brilliant yellow (12B) with narrow deep purplish red (59B) edge. Fls 170 mm wide; tepals 105 × 41-60 mm, margins smooth, tips recurved. Lvs scattered, 160 × 18 mm. Stems 1-1.1 m, brownish with green, and with 3-8 fls.

### **VI(c/a) 'My Shadow'**

cv. Parentage: Pink-edged yellow-flowered trumpet lily × deep pink-edged yellow-flowered trumpet lily

R: B. Arthur (1997), G: B. Arthur (2000), N: B. Arthur (2000), REG: B. Arthur (2005)

Inside strong red (46A), with vivid greenish yellow (2A) along centre of each tepal to 35 mm from tip, throat vivid greenish yellow (2A); outside moderate red (182A); spots and papillae absent; nectaries green; pollen strong orange (168A); stigma red. Fls 190 mm wide, slightly scented; tepals 145 × 50 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 195 × 70 mm. Stems pale green (189B), with c.22 fls. Dec. to Jan. (Australia).

(Nomenclatural standard: colour print supplied by registrant (WSY 0096426).)

### **VIII(a/b) 'Mystic Brilliant'**

cv. Parentage: 'Dame Blanche' (tetraploid form?) × seedling LO 93003.2BT

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside yellowish white (155D), with a vivid purplish red (61C) flush along midveins that fades with age; outside with outer tepals yellowish white (155D), midribs pale yellow-green (155A), light yellow-green (145B) at base with a vivid purplish red (61C) stripe, inner tepals similar but lacking purplish red stripe; spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen moderate reddish orange (171A); stigma moderate yellow-green (147D). Fls 235 mm wide, slightly fragrant; tepals 150 × 55-80 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 190 × 35 mm. Stems 0.9-1.1 m, green with darker markings, and with c.3 fls. (A *longiflorum*/Asiatic (LA) hybrid.)

### **VII(a/b) 'Mystique Pearl'**

cv. Parentage: 'Abruzzo' × 'Dynasty'

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V., REG: De Jong Lelies Holland B.V. (2004)

Inside strong/light purplish pink (63D/C), light greenish yellow (3D) above nectaries, tepal-margins narrow, yellowish white (155D), throat yellowish white (155D); outside light purplish pink (62C), midveins strong yellow-green (144C) at base, changing into yellowish white (155D), strong purplish pink (62A) towards top, strong yellow-green (144C) at top; spots strong purplish red (63A); papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen dark reddish orange (172B); stigma pale yellow-green (193B). Fls 175 mm wide, fragrant; tepals 110 × 47-63 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 145 × 26 mm. Stems 0.6-0.9 m, brownish in lower third, green with darker markings in upper two-thirds, and with 3-5 fls.

### **I(a-b/b) 'Nepra'**

cv. Parentage: 'White Princess' × 'Yellow Bowl'

N: J. Dudek, I: J. Dudek (2004), REG: J. Dudek (2004)

Inside creamy white, throat greenish; outside greenish, midribs brownish; spots purple, large, moderately numerous, in a third of each tepal; pollen brown-orange; stigma brown-purple. Fls 170 mm wide; tepals 85 × 32 mm, tips slightly recurved; pedicels 80 mm long, brown-dotted, with secondary buds. Lvs scattered, 85 × 13 mm, glossy deep green. Stems 1 m, thick, brown-dotted, with a moderately long infl. Early July.

(Raised from seeds from a hybrid produced by R.A. Prochaska.)

### **I(a/b-c) 'Netty's Pride'**

cv. Parentage unknown

R: J. Wadekamper, N: Lily Company B.V. (2004), REG: Lily Company B.V. (2004)

Inside dark greyish reddish brown/deep reddish purple (200A/77A) fading to dark red (59A), at top with broad yellowish white (155D) stripe on a third of each tepal, above nectaries with some yellowish white/dark red (155D/59A), throat yellowish white (155D); outside with outer tepals yellowish white (155D), midribs strong yellow-green (144B) at base and top and some dark red (59A) spots, inner tepals yellowish white (155D), margins with dark red (59A) spots; papillae absent; nectaries strong yellow-green (144B); pollen deep orange-yellow (163C); stigma dark red (59A). Fls 115 mm wide; tepals 60 × 16-27 mm, margins smooth, tips recurved. Lvs scattered, 92 × 7 mm. Stems 0.7 m, base brownish changing above into green with darker markings, and with c.4 fls.

#### **I(a/b) 'Nikoleta'**

cv. Parentage: 'Tavello' × 'Lidka'

R: J. Rezek (2002), N: J. Rezek (2005), I: J. Rezek (2005), REG: J. Rezek (2005)

Inside salmon-apricot, centre slightly deeper; outside pale apricot, midribs green, felty; spots dark brown-violet, medium-large, numerous, on a fifth of each tepal; nectaries white-granulated; pollen rusty brown; stigma dark violet-brown. Fls 170 mm wide; tepals 95 × 50 mm, margins smooth, tips slightly recurved; pedicels 115 mm, green, slightly hairy. Lvs scattered, glossy dark green. Stems 1 m, thick, green, slightly hairy. Late June.

#### **VIII(a/c) 'Nimrod'**

cv. Parentage: Seedling PG 95-57 × seedling RH 97-07

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light yellow (10B), brilliant yellow (9C) along midveins, outer tepals light yellow (10C), brilliant greenish yellow (6B) along midveins with some yellowish white (155B) streaks, throat pale yellow-green (4D); outside pale yellow-green (4D), margins brilliant greenish yellow (4A), midribs strong yellow-green (143C) at base, changing into light yellow-green (145B), light greenish yellow (3C) with strong yellow-green (143C) at top, inner tepals pale yellow-green (4D), margins brilliant greenish yellow (4A), pale yellow-green (155A) along midribs, midribs strong yellow-green (143C) at base, changing into pale yellow-green (4D); spots absent; papillae many, vivid yellow (9B); nectaries strong yellow-green (143C); pollen deep orange-yellow (163A); stigma light olive-grey (197B). Fls 300 mm wide, fragrant; tepals 160 × 57-98 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 172 × 44 mm. Stems 0.8-1.3 m, green, with 1 fl.  
(An Orienpet (OT) lily.)

#### **I(a/c) 'Nissouri Daydream'**

cv. Parentage: 'Daydream' × 'Snow Lark'

R: E. Horner (1992), G: E. Horner (1994), N: E. Horner (2005), I: Horner Lilies (2005), REG: E. Horner (2005)

Inside light purplish pink (55C), tepal-margins strong purplish pink (55B), throat light yellowish pink (19B) with green deep in the base; outside strong purplish pink (55B), each tepal with a brilliant yellow (20A) central stripe, outer tepals with light brown tip; spots tiny, maroon, in throat; nectaries deep red (55A); pollen brown; stigma rusty brown. Fls 150 mm wide; tepals 75 × 35 mm. margins smooth, tips not recurved. Lvs scattered, 125 × 20 mm, shiny blue-green. Stems 1 m, dark brown, with c.15 fls. July.

#### **VIII(a-b/b) 'Nova Scotia'**

cv. Parentage unknown

R: Testcentrum voor Siergewassen B.V., N: Marklily C.V. (2005), REG: Marklily C.V. (2005)

Inside yellowish white (155D), inner tepals edged with dark red (59A) in basal third; outside with outer tepals yellowish white (155D), midribs strong yellow-green (143C)

at base, light yellow-green (154D) at top, inner tepals similar but lacking light yellow-green at top, and with dark red (59A) edge in basal third; spots and papillae dark red (59A); nectaries strong yellow-green (143B); pollen brownish orange (164A); stigma light greenish yellow (1C). Fls 195 mm wide; tepals 125 × 52-62 mm, margins slightly ruffled, tips recurved. Lvs scattered, 170 × 30 mm. Stems 1.1 m, green with some darker markings, and with c.4 fls. Early to midseason.  
(A *longiflorum*/Asiatic (LA) hybrid.)

### VII(a-b/-) 'Odessa'

cv. Parentage unknown

Inside deep red/strong purplish red (60A/B/C), tepal-tips lighter, margins almost white, midveins yellow at base changing into strong red (46A); outside moderate purplish pink (66D); spots dark purple-red; papillae present; nectaries green; pollen orange-brown; stigma grey-green. Tepals medium-sized, margins ruffled, tips recurved. Lvs scattered. Stems 1.3 m, green.

(In 2004 J.A. & J. Moerman applied for registration of this name, but this was refused because the name duplicates an earlier usage by Bakker (ICNCP, 2004: Art. 27.1). However, Dutch PBR were granted to Moerman's 'Odessa' in 2004 and under ICNCP, 2004: Art. 24.4 the use of 'Odessa' for this cv. must be accepted, although it is not eligible for registration.)

### I(a-b/b) 'Oktavánka'

cv. Parentage: 'Amos' × 'Silver Bunny'

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside deep pink, tepal-tips of outer tepals paler, yellow shining through area around nectaries; outside of tepals with paler centres, midribs brownish; spots present; pollen orange-brown; stigma beige. Fls 120 mm wide; tepals 85 × 43 mm, thick and rigid, slightly recurved; pedicels 85 mm, green. Lvs scattered, 70 × 12 mm, dull pale green, sparse. Stems 0.75 m. Late June.

### I(a-b/b-c) 'Olympiáda'

cv. Parentage: ('Plavovláška' × Endeavor) × ('Plavovláška' × 'Dekora')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside creamy, centre yellowish, throat whitish, area over nectaries pale violet; outside yellowish at tepal-edges, midribs pink; spots fairly numerous, dark pink-brown, large, on three-fifths of each tepal; nectaries green, white-granulated; stigma brown-crimson. Fls 130 mm wide; tepals 87 × 37 mm, slightly recurved; pedicels 80 mm, dark brown. Lvs scattered, 75 × 12 mm, dull mid green. Stems 0.95 m, densely dotted with black-brown. Mid June.

### VII(a/b) OPIUM

Trade designation of 'Zanloropium'.

### I(a/b) 'Orange Electric'

cv. Parentage unknown

R: Laan Lelie, REG: Lily Company B.V. (2004)

Inside with inner tepals greenish white (155C), along midveins strong orange (25A) in basal half, towards tepal-margins and top changing into little stripes, outer tepals more orange in apical half, throat strong orange (25A) with greenish white (155C) edges; outside with outer tepals strong orange/strong orange-yellow (24A/B) on light orange-yellow (24C) ground, margins greenish white (155C), at top with little strong yellow-green (144B) stripe, inner tepals with broader greenish white margins; spots dark red (187A); papillae strong orange (25A); nectaries moderate yellowish pink (179D); pollen strong orange (25A); stigma strong orange (25A). Fls 110-150 mm wide; tepals 87-93

× 29 mm, margins smooth, tips not recurved. Lvs scattered, 87 × 13 mm, dark green. Stems 0.6-0.8 m, green, light green towards top, with c.9 fls in an infl. that is half total stem-length.

#### **VII(a/b) 'Pathos'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside greenish white (155C); outside with outer tepals greenish white (155C), midribs strong yellow-green (144C) at base, changing into pale yellow-green (4D), towards top light greenish yellow (4C) with some strong yellow-green (144B), inner tepals similar but lacking strong yellow-green at top; spots absent; papillae greenish white (155C); nectaries strong yellow-green (144B/C); pollen strong reddish orange/strong brown (169A/172A); stigma pale yellow-green (192A). Fls 240 mm wide; tepals 140 × 40-74 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 185 × 37 mm. Stems 0.95-1.1 m, light green with darker markings, and with c.4 fls.

#### **VII(a/b) 'Pella'**

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside yellowish white (155D), each tepal with a large light greenish yellow (6D) patch along and above nectaries, brilliant greenish yellow (6B) along midveins in basal two-thirds; outside with outer tepals yellowish white (155D), midribs strong yellow-green (143C) at base and top, in between light yellow-green (154D), inner tepals similar but lacking strong yellow-green top; spots absent; papillae brilliant greenish yellow (6B); nectaries strong yellow-green (144C) with brilliant greenish yellow (6B) edge; pollen dark reddish orange (175B); stigma pale yellow-green (193C). Fls 230 mm wide, fragrant; tepals 146 × 50-80 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 180 × 39 mm. Stems 1-1.2 m, green, with c.5 fls.

#### **I(a/b-c) 'Petrov'**

cv. Parentage: 'Novinka' × 'Windahloo'

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside yellow; outside paler yellow, midribs green; spots few, small, mahogany, on a third of each tepal, brushmarks mahogany; nectaries slightly open, green, yellow-granulated; pollen brown; stigma yellow. Fls 130 mm wide; tepals 80 × 33 mm, slightly recurved; pedicels 60 mm long, green. Lvs scattered, 80 × 14 mm, dull mid green. Stems 0.95 m, thick, green. Early July.

#### **I(b-c/c-d) 'Pilot'**

cv. Parentage: 'Plavovláška' × 'Mokka'

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside lemon-yellow, throat paler; outside paler yellow, midribs brownish at base; spots numerous, pink-brown, oval, fairly large, on two-thirds of each tepal; nectaries white-granulated; pollen brown; stigma pinkish brown. Buds slightly hairy. Fls 130 mm wide; tepals 72 × 35 mm, twisted, margins slightly wavy, tips slightly recurved; pedicels 65 mm, green, slightly hairy. Lvs scattered, 105 × 18 mm, dull pale green. Stems 0.75 m, green, slightly hairy. Early June.

#### **VII(a/b) 'Pimento'**

cv. Parentage: Red-flowered dwarf seedling × 'All Right'

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside strong/vivid purplish red (61B/C), tepal-margins yellowish white (155D), midveins strong red (46A), throat yellowish white (155D); outside with outer tepals moderate purplish red (186B), base and margins yellowish white (155D), midribs (144D) at base, changing into yellowish white (155D), towards top strong yellow-green (144C), at top strong yellow-green (144A), inner tepals moderate/dark purplish red (186B/C), base and margins yellowish white (155D), midribs light yellow-green (144D) at base, changing into yellowish white (155D), towards top strong yellow-green (144C); spots absent; papillae deep purplish red (59B); nectaries strong yellow-green (143C) with narrow light greenish yellow (1C) edge; pollen deep orange-yellow (163A); stigma moderate purplish red (186B). Fls 130-195 mm wide, fragrant; tepals 110 × 37-53 mm, margins ruffled, tips recurved. Lvs scattered, 130 × 34 mm, dark green. Stems 0.45-0.6 m, green, with c.4 fls.  
(A pot lily.)

### VIII(a-b/b-c) 'Pink Brilliant'

cv. Parentage: 'Sorbonne' × seedling LO 93003-2AT

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside vivid purplish red (57C), vivid purplish red (57D) towards tepal-margins, margins yellowish white (155B) in apical half, along midveins strong purplish red (63A), throat yellowish white (155B); outside with outer tepals yellowish white (155B), towards margins yellowish white (155B) on a vivid purplish red (67B) ground, margins yellowish white (155B) in apical half, midribs strong yellow-green (143B) at base, brilliant yellow-green (150C) at top, with a small brilliant yellow-green (150A) spot, inner tepals yellowish white (155B) with small vivid purplish red (67B) streaks, midribs strong yellow-green (143B) at base; spots absent; papillae strong purplish red (63A); nectaries strong yellow-green (144B); pollen strong orange-yellow (163B); stigma light yellow-green (145D). Fls 260 mm wide; tepals 150 × 60-70 mm, margins ruffled, tips recurved. Lvs scattered, 170-210 × 30-70 mm, dark green. Stems 0.9-1.1 m, green, with 2-6 fls.

(A *longiflorum*/Oriental (LO) hybrid.)

### I(a/b) 'Planet Jewel'

cv. Parentage: (Seedling 2504-OJ × seedling 3443.rs) × 'Stella' (tetraploid form?)

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside vivid yellow (12A); outside with outer tepals brilliant yellow (13B), with a light yellow-green (144D) stripe along midveins, strong yellow-green (143B) at top, inner tepals brilliant yellow (12B), margins vivid yellow (14B), strong yellow-green (145A) along midribs; spots absent; papillae brilliant yellow (13B); nectaries strong yellow-green (143B); pollen greyish red (178A); stigma brilliant yellow brilliant yellow (13B). Fls 200 mm wide; tepals 105 × 40-55 mm, margins smooth, tips slightly recurved. Lvs scattered, 145 × 20 mm. Stems 0.95-1 m, green with darker markings, and with 2-9 fls.

### VI(b-c/a) 'Plejáda'

cv. Parentage: (Royal Gold Group × Golden Splendor Group) × (('Black Dragon' × unknown) × unknown)

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside velvety apricot-yellow, throat deeper; outside pale apricot, midribs greenish; spots absent; nectaries green; pollen apricot; stigma yellow-green. Fls 130 mm wide; tepals 125 × 50 mm, margins somewhat wavy, tips strongly recurved; pedicels 60 mm, slightly brown-dotted. Lvs scattered, 95 × 11 mm, dull dark green. Stems 1.05 m, green. Early July.

### **VI(b-c/a) 'Polyglot'**

cv. Parentage: White-flowered tetraploid trumpet lily × deep yellow-flowered tetraploid trumpet lily

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside creamy white, throat apricot-yellow with deep yellow rays; outside with yellow shining through, midribs green with a few red-brown spots; spots absent; pollen apricot-brown; stigma greyish. Fls 160 mm wide; tepals 150 × 52 mm, slightly twisted, margins slightly wavy, tips somewhat recurved; pedicels 80 mm, very thick, brown-dotted, with secondary buds. Lvs scattered, 105 × 12 mm, dull dark green. Stems 1.25 m, dotted with red-brown. Early July. Tetraploid.

### **VI(b/a) 'Prakseda'**

cv. Parentage unknown

R: R. Widerski, N: R. Widerski, I: R. Widerski (2004), REG: R. Widerski (2004)

Inside creamy white, throat golden; outside pale green with darker midribs; spots absent; pollen pale brown; stigma green. Fls 140 mm wide; tepals 140 × 60 mm, slightly bent (outer more so), tough; pedicels 110 mm long, thick, green and brown. Lvs scattered, 100 × 10 mm, dull mid green. Stems 1.4 m, thick, brown-dotted, with long infl. Early July.

### **I(a-b/b & d) 'Pralinka'**

cv. Parentage: White-flowered tetraploid × 'Longida'

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside lemon-yellow, centre deep yellow, throat paler; outside with brownish midribs; spots numerous, dark brown-crimson, large, oval, on two-thirds of each tepal; nectaries white-granulated; pollen chocolate-brown; stigma brownish. Fls 155 mm wide; tepals thick, 90 × 47 mm, very strongly recurved, margins slightly wavy; pedicels 90 mm, very thick, freely dotted with dark brown. Lvs scattered, 130 × 25 mm, dull dark green. Stems 0.85 m, very thick, heavily brown-dotted. Early July.

### **VII(a/b) 'Pranese'**

cv. Parentage: Seedling 93-59 × seedling RH 97-09

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside yellowish white (155B), each tepal with a brilliant/light yellow (10A/B) flame, throat yellowish white (155B) with a narrow brilliant yellow (10A) edge; outside pale yellow-green (155A) flushed with pale yellow-green (4D), strong yellow-green (144C) at base, strong yellow-green (144B) at tepal-tips; spots absent; papillae yellowish white (155B) and brilliant yellow (10A); nectaries strong yellow-green (144A/B); pollen strong brown (172A); stigma pale yellow-green/dark purple (196C/79B). Fls 250 mm wide; tepals 145-155 × 47-66 mm, margins slightly ruffled, tips recurved. Lvs scattered, 190 × 47 mm. Stems 1.2 m, light green, with c.6 fls.

### **VI(b/b) 'Prince of Nitra'**

cv. Parentage: 'White Henryi' × ('Damson' × 'Mabel Violet')

R: A. Mego (1993), N: A. Mego (2005), I: A. Mego (2005), REG: A. Mego (2005)

Inside white suffused with pink, tepal-centres and throat golden yellow; outside deep pinkish purple, midribs green; spots absent; nectaries dark green; pollen ochre-yellow; stigma grey-green. Fls 150 mm wide, slightly scented; tepals 130 × 45 mm, firm and waxy, strongly recurved; pedicels 120 mm, thick, brown-dotted. Lvs scattered, 170 × 20 mm, dull mid green. Stems 1.5 m, thick, dark green. Early July.

### VIII(a/b) 'Prinses Amalia'

cv. Parentage: ('Gran Paradiso' × 'Compass') × ('Sterling Star' × 'Connecticut King')  
R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2005)

Inside strong orange (30D), tepal-margins vivid orange (28B), brilliant orange (25C) along basal half of midveins, throat vivid yellowish pink (28A) with light orange (28D) edges; outside with outer tepals light orange (28D), changing into strong orange (30D) towards margins, inner tepals slightly paler; spots strong purplish red (59D); papillae vivid orange (28B); nectaries strong yellow-green (143A); pollen strong reddish orange (169A); stigma strong orange (168A). Fls 190 mm wide; tepals 107 × 40-60 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 135 × 25 mm. Stems 0.95-1.2 m, green with darker markings, and with 9-12 fls.  
(A *longiflorum*/Asiatic (LA) hybrid.)

### I(b/b) 'Prochora'

cv. Parentage unknown

R: J Dudek, N: J. Dudek, I: J. Dudek (2004), REG: J. Dudek (2004)

Inside pink, tepal-centre cream, throat pink; outside pink, midribs green; spots purple, oblong, large, moderately dense, on half of each tepal; stamens rudimentary, brownish, lacking pollen; stigma purple. Fls 130 mm wide; tepals 68 × 30 mm, slightly twisted, margins slightly ruffled, tips recurved. Lvs scattered, 130 × 14 mm, glossy dark green. Stems 1.5 m, thick, green-brown, hairy in upper part. Mid July.

### VIII(a/b) PROMESA

Trade designation of 'Zanlapromes'

### I(a-b/b) 'Prunotto'

cv. Parentage: Seedling AJ 95-21 × seedling AJ 96-7

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside vivid red/vivid reddish orange (45A/34A), tepal-margins vivid reddish orange (30A), throat moderate reddish orange (35A); outside with outer tepals (169A/171A), margins vivid red (45A), strong orange (170A) at base, strong yellow-green (144A) at top, inner tepals vivid to strong reddish orange (34A/B/C), midribs strong brown (172A); spots dark purple (79A); papillae few; nectaries strong yellowish pink (37B); pollen moderate reddish orange/strong brown (171A/172A); stigma dark red/dark purple (187B/79A). Fls 140 mm wide; tepals 100 × 36-56 mm, margins smooth, tips slightly recurved. Lvs scattered, 141 × 19 mm, dark green. Stems 0.9 m, light green, with c.11 fls. Tetraploid?

### VIII(a/b) 'Purple King'

cv. Parentage unknown

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside pale purplish pink (65C), each tepal with a strong purplish red (60B) flame, throat yellowish white (155D); outside with outer tepals moderate purplish red (186A/B), dark purplish red (186C) towards margins, margins very pale purple (73D), midribs strong yellow-green (144C) at base and top, inner tepals very pale purple (69C), moderate purplish red (186A) along midribs, brilliant yellow-green (142B) at base; spots and papillae absent; nectaries strong yellow-green (144B) with narrow brilliant greenish yellow (2B) margins; pollen greyish brown (166A); stigma light greenish grey (188B). Fls 195 mm wide, fragrant; tepals 150 × 44-63 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 140 × 20 mm. Stems 1-1.15 m, dark green with darker markings, and with 2-5 fls.  
(An Orienpet (OT) lily.)

### **VII(a-b/b) 'Rain Dance'**

cv. Parentage unknown

R: Mak Breeding B.V., N: Liliande B.V. (2003), REG: Mak 't Zand B.V. (2005)

Inside yellowish white (155D), young fls with yellow at base of midveins; outside yellowish white (155D); spots absent; papillae present; nectaries green; pollen brownish orange (171B); stigma deep purplish red (61A). Fls 260 mm wide; tepals 160-162 × 62-90 mm, margins strongly ruffled, tips recurved. Lvs scattered, 180-186 × 23-52 mm. Stems 0.9-1.1 m, green with some darker markings, and with 2-9 fls.

### **VII(a/b) RAVENNA**

Trade designation of 'Zanlorvenna'

### **VIII(a-b/b) 'Redding'**

cv. Parentage: Seedling E 905-6 × seedling OR 2

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside brilliant yellow (12B), lighter towards tepal-margins and tips, midveins vivid yellow (12A) at base; outside light greenish yellow (8C); spots absent; papillae present; nectaries yellow-green; pollen orange-red; stigma green. Tepals short to medium-sized, margins ruffled, tips strongly recurved. Lvs scattered, medium-sized, dark green. Stems c.1.3 m, green, with 2-10 fls. Late.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096408).)

### **VIII(c-d/b) 'Red Hot'**

cv. Parentage: Seedling 821-11 × Oriental mixture

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside strong reddish orange (42C), becoming bluer with age, tepal-margins brilliant yellow/light greenish yellow (8A/B), midveins orange-yellow at base changing into red-pink; outside strong/light purplish pink (63C/D), tepal-margins brilliant yellow (8A); spots and papillae present; nectaries green changing into red; pollen strong orange (168A); stigma green. Fls 190 mm wide; tepals 125-130 × 35-57 mm, margins slightly ruffled, tips recurved. Lvs scattered, 71-207 × 18-40 mm, dark green. Stems 1-1.2 m, green, with 3-12 fls.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096405).)

### **VII(a/b) 'Red Lips'**

cv. Parentage: Red-flowered seedling × ('Rosalie' × 'Star Gazer')

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside with inner tepals deep red (60A), outer tepals deep red (60A) with narrow very pale purple (69D) margins, throat yellowish white (155D); outside with outer tepals deep purplish red (61A), yellowish white (155D) at base, midribs strong yellow-green (143C) at base changing into deep purplish red (61A), inner tepals strong purplish red (60C), towards margins moderate purplish red (64A), yellowish white (155D) at base, midribs strong yellow-green (143C) at base changing into strong purplish red (61B); spots absent; papillae dark red (59A); nectaries strong yellow-green (144C) with narrow brilliant yellow-green (154C) edge; pollen deep orange-yellow (163A); stigma deep reddish purple (77A). Fls 240-260 mm wide, fragrant; tepals 145 × 45-55 mm, margins slightly ruffled, tips recurved. Lvs scattered, 135 × 37 mm. Stems 0.85-1.1 m, green with darker markings at base, and with c.6 fls.

### **I(a-b/b) 'Ripro'**

cv. Parentage unknown

N: J. Dudek, I: J. Dudek (2004), REG: J. Dudek (2004)

Inside pink, throat with cream band; outside with creamy midribs; spots purple, small, sparse, over a half of each tepal; stamens rudimentary, pink, lacking pollen; stigma pinkish. Fls 180 mm wide; tepals 60 × 20 mm, margins slightly ruffled, tips slightly recurved; pedicels 70 mm long. Lvs scattered, 90 × 10 mm, dull mid green. Stems 1 m, thick, dotted, with a moderately long infl. Late June.

(Raised from seeds from a hybrid produced by R.A. Prochaska.)

### **VIII(c/b-c) 'Robert Griesbach'**

cv. Parentage unknown

R: R. Griesbach, N: Lily Company B.V. (2004), REG: Lily Company B.V. (2004)

Inside dark red (59A), changing into yellowish white (155D) towards tepal-margins and tips, brilliant yellow (13B) above and next to nectaries, throat pale yellow green (4D); outside with outer tepals yellowish white (155D), up to two-thirds of each tepal flushed with strong purplish red (59D), strong yellow-green (143C) along midribs changing into light yellow-green (144D), strong yellow-green (143C) at top, inner tepals similar but with light yellow (21D) along midribs at base; spots and papillae absent; nectaries strong yellow-green (143C); pollen strong brown (172A); stigma very pale green (189D). Fls 160-200 mm wide; tepals 120 × 33-43 mm, margins smooth, tips strongly recurved. Lvs scattered, 155 × 27 mm. Stems 1-1.45 m, green with darker markings, and with c.4 fls. (An Orienpet (OT) lily.)

### **VIII(a-b/b-c) 'Robina'**

cv. Parentage: Orienpet (OT) seedling × 'Mero Star'

R: Testcentrum voor Siergewassen B.V., N: Marklily C.V. (2001), REG: Marklily C.V. (2004)

Inside moderate purplish red (70A), strong purplish red (61B) along midveins, yellowish white (155D) next to and above nectaries, throat yellowish white (155D); outside with outer tepals moderate purplish red (70A), margins strong purplish red (71B), yellowish white (155D) at base, midribs strong yellow-green (143C) at base with moderate purplish red (70A) streak, strong yellow-green (143C) at top, inner tepals strong reddish purple (70B), margins moderate purplish red (70A), basal third of tepals yellowish white (155D) with purple veins, midribs strong yellow-green (143C) at base changing towards top into moderate purplish red (186B); spots absent; papillae few, moderate purplish red (70A); nectaries strong yellow-green (143C); pollen strong orange (25A); stigma dark red (187C). Fls 255 mm wide, fragrant; tepals 150 × 50-80 mm, margins slightly ruffled, tips recurved. Lvs scattered, 200 × 41 mm. Stems 1.1-1.5 m, green with darker markings, and with 3-5 fls.

(An Orienpet (OT) lily.)

### **VIII(a/b) 'Rodin'**

cv. Parentage: Seedling BG 97-14 × seedling AG 95-10

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2001), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside light greenish yellow (5D), brilliant greenish yellow (5B) along midveins and on tepal-margins, throat yellowish white (155D); outside pale greenish yellow (2D), tepal-margins light greenish yellow (8B), midribs strong yellow-green (141D) at base; spots and papillae absent; nectaries strong yellow-green (143A); pollen moderate brown (165A); stigma strong yellow-green (144C). Fls 160 mm wide, slightly fragrant; tepals 126 × 43-58 mm, margins smooth, tips recurved. Lvs scattered, 142 × 20 mm. Stems 1-1.35 m, green with some darker markings, and with 5-6 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

### VI(c/d) 'Romanita'

cv. Parentage: 'Henrietina' × 'Brighton'

R: R. Widerski, N: R. Widerski, I: R. Widerski (2004), REG: R. Widerski (2004)

Inside deep orange, tepal-tips paler; outside pale orange with green midribs; brown-orange dashes on a half of each tepal; nectaries green; pollen yellow; stigma green.

Fls 120 mm wide; tepals 85 × 14 mm, strongly recurved, slightly twisted and ruffled; pedicels 100 mm long, thick, green and brown, with tertiary buds. Lvs scattered, glossy mid green. Stems 1.4 m, thick, green and brown, with long infl. Mid July.

### VI(b/a-b) 'Romikanda'

cv. Parentage: Unknown × 'Bamberg'

R: R. Widerski, N: R. Widerski, I: R. Widerski (2004), REG: R. Widerski (2004)

Inside pink, throat yellow; outside dark pink, midribs darker; spots absent; pollen yellow; stigma pale brown. Fls 120 mm wide, shortly trumpet-shaped; tepals 140 × 60 mm, slightly curved, tough; pedicels 80 mm long, brown and green. Lvs scattered, 110 × 11 mm. Stems 1.2 m, thick, brown and green, with medium-sized infl. Early July.

### VIII(a-b/b) 'Roparun'

cv. Parentage: 'Paola' × seedling LA 93008-6T

R: De Jong Lelies Beheer B.V., N: Flora Holland/De Jong Lelies Beheer B.V. (2003), REG: De Jong Lelies Holland B.V. (2004)

Inside with inner tepals moderate red (180B/C), with some moderate red/dark yellowish pink (181C/D) along top of midveins; outer tepals similar but with some moderate red (180A) along midveins and towards margins, throat dark greenish yellow (152C) with dark yellowish pink (181D) on edges; outside with outer tepals moderate purplish red to dark yellowish pink (186A/B-181D), margins strong yellowish pink (34D), tips with some strong yellow-green (144C), inner tepals (181C/D), moderate purplish red (186B) along midribs, margins strong yellowish pink (34D); spots few, dark red (187A); papillae absent; nectaries dark yellowish pink (181D); pollen dark reddish orange (173A); stigma deep purplish red (187D). Fls 180 mm wide; tepals 102-110 × 37-49 mm, margins smooth, tips slightly recurved. Lvs scattered, 176 × 16 mm dark green. Stems 0.85-1.2 m, green with darker markings, and with 3-9 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

### VII(a/b) 'Rosy Dimple'

cv. Parentage: Little Rascal Group (white) × Little Rascal Group (pink)

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V. (2003), REG: Bischoff Tulleken Lelies B.V. (2004)

Inside with inner tepals yellowish white (155D), dark purplish pink (186C) along apical two-thirds of midveins, basal third light greenish yellow (6D), outer tepals with basal half yellowish white (155D), apical half deep purplish pink (70C), yellowish white (155D) towards margins, up to a third along midveins (6C), throat yellowish white (155D); outside with outer tepals yellowish white (155D), along apical two-thirds of midribs moderate purplish pink (186D), base and top of midribs strong yellow-green (144C), in between light greenish yellow (1C), inner tepals similar but less purplish pink; spots strong purplish red (61B); papillae yellowish white (155D) and light greenish yellow (6D); nectaries strong yellow-green (144C); pollen strong orange (169C); stigma pale green (189C). Fls 100-140 × 33-44 mm, margins ruffled, tips recurved. Lvs scattered, 110 × 32 mm. Stems 0.4-0.65 m, green, with 13-15 fls.

### VIII(a/b-c) 'Royal Hall'

cv. Parentage: Seedling BG 95-60 × seedling G92-10

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside pale greenish yellow (1D), tepal-margins and midveins brilliant greenish yellow (1B), throat greenish white (157D); outside pale yellow-green (4D), tepal-margins light yellow-green (2C), midribs brilliant yellow-green (142B) at base; spots absent; papillae brilliant greenish yellow (1B); nectaries strong yellow-green (143C); pollen vivid orange-yellow (23A); stigma pale greenish yellow (1D). Fls 170 mm wide; tepals 120 × 50-60 mm, margins slightly ruffled, tips recurved. Lvs scattered, 150 × 25 mm. Stems 1-1.2 m, green, with c.5 fls.

(*A. longiflorum*/Asiatic (LA) hybrid.)

#### **VIII(a/b) 'Royal Show'**

cv. Parentage: Seedling BG 95-23 × seedling AG 95-04

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside vivid yellow (12A), throat light greenish yellow (1C); outside vivid yellow (12A), midribs flushed with strong yellow-green (143B), strong yellow-green (143C) at top; spots and papillae absent; nectaries strong yellow-green (143B); pollen strong brown (172A); stigma brilliant greenish yellow (4A). Buds slightly hairy. Fls 220 mm wide, slightly fragrant; tepals 120 × 60 mm, margins smooth, tips slightly recurved. Lvs scattered, 140 × 25 mm. Stems 1.05-1.1 m, green with darker markings in lower part, and with c.6 fls.

(*A. longiflorum*/Asiatic (LA) hybrid.)

#### **VI(b-c/b) 'Rybnik'**

cv. Parentage: 'Aldona' × 'Traumstein'

R: R. Widerski, N: R. Widerski, I: R. Widerski (2004), REG: R. Widerski (2004)

Inside creamy yellow, tepal tips creamy, throat with orange-yellow rays; spots absent; pollen yellow; stigma bronze-brown. Fls 180 mm wide; tepals 130 × 55 mm, curved, slightly ruffled and twisted, tough; pedicels 120 mm long, green and brown. Lvs scattered, 150 × 13 mm, dull mid green. Stems 1.4 m, thick, bronze-brown, with long infl. Early July.

#### **VIII(a/b) 'San Stefano'**

cv. Parentage: Seedling BG 95-60 × seedling AG 95-04

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside light greenish yellow (7D), brilliant greenish yellow (5A) along tepal-margins and midveins, inner tepals dark red (187A) on edges at base, throat pale greenish yellow (2D); outside light greenish yellow (4C), brilliant greenish yellow (6B) on tepal-margins; midribs (141D) at base and top, inner tepals dark red (187A) on edges at base; spots few, dark red (59A); papillae brilliant greenish yellow (5A); nectaries strong yellow-green (143A); pollen moderate reddish brown (175A); stigma light greenish yellow (1C). Fls 185 mm wide, slightly fragrant; tepals 115 × 53-65 mm, margins smooth, tips slightly recurved. Lvs scattered, 135 × 28 mm. Stems 1.05-1.4 m, green, with 4-6 fls.

(*A. longiflorum*/Asiatic (LA) hybrid.)

#### **VI(b/a) 'Saulkrasti'**

cv. Parentage: 'Zeltite' × seedling 376-S-16

R: Ā. Zorgevics (1987), G: Ā. Zorgevics (1992), N: Ā. Zorgevics (2005), I: University of Latvia Botanical Garden (2005), REG: Ā. Zorgevics (2005)

Inside brilliant yellow (13C), throat brilliant yellow-green (150A); outside greenish with brownish gleam; spots absent; nectaries colourless; pollen light brown; stigma green. Fls 140 mm wide, strongly scented; tepals 130 × 30 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 70 × 10 mm. Stems 1.2 m, green, with c.9 fls. Late July.

### VII(a/b) 'Savio'

cv. Parentage: Seedling 88-131 × seedling 93-13

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (1999), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside deep/strong purplish pink (68A/B), lighter towards base, midveins white at base; outside dark purplish pink (186C); spots and papillae absent; nectaries green; pollen orange-brown; stigma yellow-green. Tepals short to medium-sized, margins slightly to moderately ruffled, tips recurved. Lvs scattered. Stems 1.25 m, green with darker speckles and stripes. Early to midseason.

### VIII(a/b) 'Sea Brilliant'

cv. Parentage: 'Dame Blanche' (tetraploid form?) × seedling LO 93003.2BT

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside with inner tepals yellowish white (155D), strong purplish red (58C) along basal half of midveins, outer tepals more purplish red (58C) (purplish red fades with age); outside yellowish white (155D), pale yellow-green (155A) along midribs, midribs strong yellow-green (144C) at base and top; spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen moderate reddish brown (175A); stigma light yellow-green (145D). Fls 245 mm wide, fragrant; tepals 155 × 60-75 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 200 × 45 mm. Stems 0.7-1.2 m, green with darker markings, and with 1-5 fls.  
(A *longiflorum*/Oriental (LO) hybrid.)

### VIII(a/b) 'Sidorno'

cv. Parentage: Seedling BG 95-61 × seedling AG 96-15

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals light greenish yellow (3D), margins and above throat brilliant greenish yellow (3A), outer tepals more brilliant greenish yellow (3A), throat pale greenish yellow (1D); outside light greenish yellow (4C), tepal-margins brilliant greenish yellow (5B); spots absent; papillae brilliant greenish yellow (3A); nectaries strong yellow-green (143B); pollen dark greyish reddish brown (200A); stigma deep reddish purple (77A). Buds hairy. Fls 155 mm wide; tepals 112 × 46-64 mm, margins smooth; tips recurved; pedicels hairy at base. Lvs scattered, 145 × 17 mm. Stems 0.95-1.3 m, brownish red with green, with c.6 fls.  
(A *longiflorum*/Asiatic (LA) hybrid.)

### VII(a/b) 'Snow Pearl'

cv. Parentage: 'Vedea' × 'Spinoza'

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside yellowish white (155D); outside yellowish white (155D), midribs strong yellow-green (144C) at base and top, in between pale yellow-green (155A); spots absent; papillae yellowish white (155D); nectaries moderate yellow-green (143D); pollen strong/moderate reddish orange (169A/171A); stigma pale yellow-green (193D). Fls 220 mm wide, fragrant; tepals 130 × 42-68 mm, margins slightly ruffled, tips recurved. Lvs scattered, 140 × 37 mm, dark green. Stems 0.9-1.15 m, green with darker markings, and with 3-8 fls.

### VII(a/-) 'Solo'

cv. Parentage unknown

N: A. van der Marel B.V. (2003), REG: Van der Marel Lelie B.V. (2005)

Inside greenish white (155C), tepals brilliant yellow (12B) in basal part; outside

greenish white (155C); spots yellow; papillae present; nectaries yellow-green; pollen orange-brown; stigma purple. Tepals medium-sized, margins strongly ruffled, tips slightly recurved. Lvs scattered. Stems mid-height, green. Midseason.

#### **VIII(a/b) 'Sonetto'**

cv. Parentage: Seedling PG 95-39 × seedling RH 94-13

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside pale yellow-green (4D), tepal-margins brilliant yellow (10A), basal half of midveins brilliant yellow (10A); outside with outer tepals pale yellow-green (4D), basal half with pale yellow (8D) margins, midribs light yellow-green (144D) at base, changing towards top into pale yellow (8D) and with strong purplish red (60C) flush (absent in young fls), inner tepals similar but with less yellow-green at base, pale yellow margins broader and purple flush absent; spots absent; papillae light greenish yellow (8B); nectaries strong yellow-green (144C); pollen strong orange (25A); stigma light purple (77C). Fls 215 mm wide; tepals 130 × 55-78 mm, margins slightly ruffled, tips recurved. Lvs scattered, 190 × 47 mm. Stems 1.15-1.2 m, green, with 2-3 fls. (An Orienpet (OT) lily.)

#### **VIII(a/b) 'Soprano'**

cv. Parentage: Seedling PG 95-39 × seedling RH 95-01

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals pale yellow-green (4D), brilliant greenish yellow (4A) along basal half of midveins, outer tepals similar but with less brilliant greenish yellow, throat (157D); outside with outer tepals pale yellow-green (4D), midribs strong yellow-green (144C) at base, light greenish yellow (4C) with strong yellow-green (144C) at top, with a strong purplish red (60C) flush along basal half of midribs (disappearing with age), inner tepals similar but lacking yellow-green (144C) at top and lacking purplish red flush; spots absent; papillae brilliant yellow (8A); nectaries strong yellow-green (144C) edged with brilliant greenish yellow (4A); pollen greyish brown/moderate reddish brown (166A/175A); stigma dark purple (79A). Fls 250 mm wide, slightly fragrant; tepals 165 × 59-99 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 175 × 40 mm. Stems 1.05-1.3 m, green with darker markings, and with 2-3 fls. (An Orienpet (OT) lily.)

#### **VIII(a/b) 'Space Mountain'**

cv. Parentage: Seedling 650-3 × seedling 88.212-2

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside strong/moderate purplish red (63C/186B), tepals yellow-green in basal part, midveins yellow at base; outside strong purplish red (63C); spots and papillae present; nectaries green; pollen brown; stigma purple with green star. Tepals short to medium-sized, margins slightly ruffled, tips recurved. Lvs scattered, dark green. Stems tall, green, with c.9 fls. Late.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096399).)

#### **VIII(a/a) SPACER**

Trade designation of 'Zanlaspacer'

#### **VI(b-c/d) 'Split'**

cv. Parentage: (('Oneida' × 'Gold Eagle') × 'Gold Eagle') × ('Oneida' × ('Oneida' × orange-flowered Trumpet))

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside white with beige tinge, each tepal with a broad pale orange ray originating in throat; outside tinged with pale pink; dusting of tiny cinnamon lines on half of each tepal; nectaries yellow-green; pollen chocolate-brown; stigma green. Fls 115 mm wide; tepals 95 × 38 mm, margins roughly wavy, tips very recurved; pedicels 85 mm, lightly brown-dotted. Lvs scattered, 105 × 16 mm, dull dark green. Stems 1.45 m, dotted with dark brown. July.

#### **VIII(a/b) 'Stafford'**

cv. Parentage: Seedling BJ 95-17 × seedling AJ 94-15

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside light yellow/light orange-yellow (18A/16B), tepal-margins and tips vivid yellow/light orange-yellow (16A/B), strong orange-yellow (17A) along midveins, throat vivid orange-yellow (23A), margins vivid/light yellow (17C/D); outside with outer tepals light yellow/pale orange-yellow (18B/16D), margins strong orange-yellow (17A), at tip light yellow (18B), at base with moderate red (181A) patch, inner tepals light yellow (18B), light yellow (18A) towards margins, at tip strong orange-yellow (17A); spots and papillae absent; nectaries light orange-yellow (19A); pollen vivid reddish orange (34B); stigma moderate red/greyish red (183C/184A). Fls 242 mm wide; tepals 150-157 × 49-72 mm, margins smooth, tips slightly recurved. Lvs scattered, 190 × 29 mm. Stems 1.1-1.2 m, green with darker markings, and with c.3 fls. (A *longiflorum*/Asiatic (LA) hybrid.)

#### **I(a-b/b) 'Star Jewel'**

cv. Parentage: (Seedling 2504-OJ × ('Prato' × 'Stella' (tetraploid form?)) × ('Michella' (tetraploid form?) × 'Saba Jewel')

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside strong orange-yellow/vivid yellow (17A/B), throat light yellow-green (154D); outside vivid yellow (12A/13A), tepal-margins strong orange-yellow/vivid yellow (17A/B), at top with some strong yellow-green (144A), at base strong yellow-green (144A/B); spots and papillae absent; nectaries vivid greenish yellow (2A) with strong yellow-green (144A) stripes; pollen dark red (183A); stigma deep red (53A/185A). Fls 145 mm wide; tepals 106 × 37-53 mm, margins smooth, tips slightly recurved. Lvs scattered, 124 × 20 mm, dark green. Stems 0.85-1.1 m, brownish with green spots, and with 2-7 fls.

#### **I(a-b/c) 'Stilnox'**

cv. Parentage: 'Español' × white-flowered tetraploid

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside lemon-yellow, tepal-tips and throat paler; outside paler with mahogany midribs; spots dark pink-brown, tiny, on a third of each tepal; nectaries green, white-granulated; pollen chocolate-brown; stigma brown. Fls 145 mm wide; tepals 80 × 44 mm, tips slightly recurved; pedicels 55 mm, thick, heavily brown-dotted. Lvs scattered, 95 × 16 mm, dull mid green. Stems 0.65 m, thick, heavily dotted with dark brown. Mid July. Triploid.

#### **VII(a/b) 'Striker'**

cv. Parentage: Seedling 88-152 × seedling 93-41

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2003), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside vivid purplish red (57D), tepal-margins pale purplish pink/yellowish white (65D/155D), throat yellowish white (155D); outside dark/moderate purplish pink (186C/D), base of tepals yellowish white (155D), midribs light yellow-green (144D) at base, changing into yellowish white (155D), at top moderate yellow-green (146D);

spots moderate purplish red (58A); papillae vivid purplish red (57D); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma deep reddish purple (77A). Fls 230 mm wide, fragrant; tepals 125 × 50-72 mm, margins slightly ruffled, tips recurved. Lvs scattered, 145 × 34 mm. Stems 1-1.1 m, green with darker markings, and with 4-5 fls.

(Colour of young fls is more intense.)

#### **I(a/b) 'Sunday Parfait'**

cv. Parentage unknown

R: J. Wionzek (1998), G: Hillcrest Harmony Flowers (2000), N: D. Nelson (2004), I: Hillcrest Harmony Flowers (2005), REG: D. Putland (2005)

Inside strong orange (24A) on basal half of tepals, strong red (47B) on apical half; outside with outer tepals light orange-yellow (25D) edged with vivid reddish orange (34B), inner tepals with prominent moderate red (179B) midribs; spots dark red (59A) on basal part of tepals; papillae frosty white; nectaries green; pollen brownish orange (171B); stigma strong red (47B). Fls 119 mm wide; tepals 75-79 × 21-25 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 170 × 20 mm. Stems 0.9 m, light green, with 22-25 fls. July.

(Published in *Hillcrest Harmony Flowers Cat.*, 2005. Nomenclatural standard: colour print supplied by registrant (WSY 0096423).)

#### **I(a/b) 'Sun Kiss'**

cv. Parentage unknown

N: Lily Company B.V. (2004), REG: Lily Company B.V. (2004)

Inside vivid yellow (12A), throat pale greenish yellow (1D); outside brilliant yellow (8A), vivid yellow (12A) towards tepal-margins, strong yellow-green (141D) at top; spots and papillae absent; nectaries strong yellow-green (143C); pollen absent; stigma brilliant yellow (7A). Fls 125 mm wide; tepals 75 × 27-32 mm, margins slightly ruffled, tips slightly recurved; pedicels hairy at base. Lvs scattered, 135 × 13 mm, dark green. Stems 0.5-0.8 m, green, with some darker markings in lower half, and with c.18 fls.

#### **I(b/c) 'Sutherland Fascination'**

cv. Parentage: 'Merryspring' × 'Sutherland Peach Melody'

R: D.J. Hay (1996), G: D.J. Hay (1999), N: D.J. Hay (2001), REG: D.J. Hay (2004)

Inside with tepal-tips and base bright pink, central part gold and cream, throat pink; outside pink; spots tiny, at base of tepals; nectaries pink; pollen orange; stigma pink. Fls 130 mm wide; tepals 65 × 28 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 53 × 14 mm. Stems 0.45 m, red-brown in lower part, green above, with c.7 fls. Midseason.

(Nomenclatural standard: colour print supplied by registrant (WSY 0096414).)

#### **VI(b-c/d) 'Světloňoř'**

cv. Parentage: ((*benyri* × 'Gold Eagle') × 'Mimosa Star') × ('Lady Jane' × 'Gold Eagle')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside vivid yellow, tepal-tips paler; outside with brownish midribs; yellow, raised, papillose spots on half of each tepal; nectaries green; pollen apricot; stigma green. Fls 125 mm wide; tepals 95 × 45 mm, thick, moderately curved, margins slightly wavy; pedicels 60 mm, thick, dotted with dark brown. Lvs scattered, 105 × 15 mm, dull dark green. Stems 1.15 m, freely brown-dotted. Mid July.

#### **VIII(a/b) 'Swansea'**

cv. Parentage: Seedling BG 95-60 × seedling AJ 94-15

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside strong orange-yellow (24B), strong orange (24A) towards tepal-margins and along midveins, throat moderate yellow-green (139D); outside light orange-yellow (19A), tepal-margins strong orange (25A), midribs moderate yellow-green (146D) at base and top; spots absent; papillae strong orange (24A); nectaries moderate yellow-green (146B); pollen vivid reddish orange (30B); stigma strong orange (30D). Fls 222 mm wide; tepals 128 × 50-60 mm, margins smooth, tips recurved. Lvs scattered 155 × 27 mm. Stems 1.1-1.4 m, green with darker markings especially at base, and with c.7 fls.  
(*A longiflorum*/Asiatic (LA) hybrid.)

### VII(a/b) 'Sweet Pearl'

cv. Parentage: 'Montrachet' × 'Medusa'

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside with inner tepals moderate purplish pink (186D), yellowish white (155D) along and above nectaries, tepal-margins yellowish white (155D) in apical half, outer tepals stronger moderate purplish pink (186D), throat yellowish white (155D); outside with basal half of tepals yellowish white (155D), apical half moderate purplish pink (186D), midribs strong yellow-green (143C) at base and top, in between light yellow-green (154D); spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144B); pollen vivid yellowish pink (28A); stigma dark red (187C). Fls 182 mm wide, fragrant; tepals 116 × 41-63 mm, margins smooth to ruffled, tips slightly recurved. Lvs 139 × 45 mm. Stems 0.85-1.3 m, green with darker markings at base, and with 4-7 fls. (The purplish pink colour of outer tepals is less prominent than in inner tepals and it fades with age.)

### I(a-b/b) 'Tábor'

cv. Parentage unknown

R: J. Kovář, N: J. Kovář (2002), I: J. Kovář (2002), REG: J. Kovář (2005)

Inside orange, centre of tepals and throat brighter; outside pale apricot-orange; spots violet-brown, large, elongated, on three-quarters of each tepal; nectaries orange, white-granulated; pollen orange-brown; stigma dark violet-brown. Fls 140 mm wide; tepals 85 × 45 mm, scarcely recurved; pedicels 70 mm, profusely dotted with dark violet-brown. Lvs scattered, 95 × 17 mm, slightly glossy deep green. Stems 1 m, very thick, profusely dotted with dark violet-brown, with brownish bulbils. Late June.

### VI(b-c/d) 'Tanečnice'

cv. Parentage: (('Damson' × unknown) × (*henryi* × 'Gold Eagle') × 'Mimosa Star') × (('Honeydew' × 'Gold Eagle') × (Silver Sunburst Group × 'Gold Eagle'))

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside lemon-yellow, tepal-tips paler; outside with green midribs; spots numerous, linear, tiny, brown, in lines on two-thirds of each tepal; nectaries green; pollen small in amount, brown-orange; stigma green. Fls 105 mm wide, compressed turban-shaped; tepals strongly recurved; pedicels 60 mm, brown-dotted, with secondary buds. Lvs scattered, 95 × 30 mm, dull mid green. Stems 1 m, heavily dotted with red-brown. Early July.

### VI(b-c/c-d) 'Tatra's Eagle'

cv. Parentage: *henryi* × ('Caramba' × (('White Henryi' × unknown) × ('Tetra White Henryi' × unknown)))

R: A. Mego (1994), N: A. Mego (2005), I: A. Mego (2005), REG: A. Mego (2005)

Inside pale ochre-yellow, tepal-centres and throat yellow; outside greenish yellow; sparse chocolate-brown dashes on a third of each tepal; papillae in throat; nectaries dark green; pollen brown; stigma green. Fls 160 mm wide; tepals 100 × 30 mm, moderately

recurved, margins wavy; pedicels 170 mm, brown-dotted, with secondary buds. Lvs scattered, 190 × 25 mm, dark green. Stems 1.8 m, densely brown-dotted, with a very long infl. Mid July.

**VII(a/b) ‘Tesoro’**

cv. Parentage: Seedling RF 94-07 × seedling RM 97-14

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside with inner tepals vivid purplish red (67B), margins moderate purplish red (70A) above each nectary a small light greenish yellow (3C) patch, tips yellowish white (155D), outer tepals similar but strong purplish red (67A) towards margins, throat yellowish white (155D); outside with outer tepals moderate/strong purplish red (64A/B), yellowish white (155D) at base, midribs strong yellow-green (144A) at base changing into strong yellow-green (144B), inner tepals moderate purplish red/dark purplish pink (186B/C), moderate purplish red (64A) towards margins, next to midribs yellowish white (155D), midribs strong yellow-green (143C) at base changing into strong yellow-green (144B); spots absent; papillae moderate purplish red (58A); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma pinkish grey (201D) with a little moderate purplish red (70A). Fls 270 mm wide, fragrant; tepals 140 × 57-72 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 210 × 40 mm, dark green. Stems 1-1.1 m, green, with c.5 fls.

**I(b/-) ‘Tetra Ambrosia’**

REG: ICRA (2005)

Tetraploid form of ‘Ambrosia’.

**I(a/-) ‘Tetra Capri’**

REG: ICRA (2005)

Tetraploid form of ‘Capri’.

**VII(b/b) ‘Tetra Dame Blanche’**

REG: ICRA (2005)

Tetraploid form of ‘Dame Blanche’.

**I(a/-) ‘Tetra Gloria’**

REG: ICRA (2005)

Tetraploid form of ‘Gloria’.

**VI(-/a) Tetra Golden Clarion Group**

REG: ICRA (2005)

Tetraploid form of Golden Clarion Group.

**I(a/-) ‘Tetra Goldina’**

REG: ICRA (2005)

Tetraploid form of ‘Goldina’.

**I(a/b) ‘Tetra Gran Paradiso’**

REG: ICRA (2005)

Tetraploid form of ‘Gran Paradiso’.

**I(a/-) ‘Tetra Michella’**

REG: ICRA (2005)

Tetraploid form of ‘Michella’.

**I(a/b) ‘Tetra Monaco’**

REG: ICRA (2005)

Tetraploid form of ‘Monaco’.

**I(a/-) ‘Tetra Napoli’**

REG: ICRA (2005)

Tetraploid form of ‘Napoli’.

**VII(-/d) ‘Tetra Rachel Pappo’**

REG: ICRA (2005)

Tetraploid form of ‘Rachel Pappo’.

**VIII(a-b/b) ‘Tetra Sea Treasure’**

REG: ICRA (2005)

Tetraploid form of ‘Sea Treasure’.

**I(a/-) ‘Tetra Stella’**

REG: ICRA (2005)

Tetraploid form of ‘Stella’.

**I(a/-) ‘Tetra Unique’**

REG: ICRA (2005)

Tetraploid form of ‘Unique’.

**VIII(a/b) ‘Time Diamond’**

cv. Parentage: ‘Goldina’ × seedling LA 93007-4T

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong

Lelies Holland B.V. (2004)

Inside with inner tepals light orange-yellow (22B), strong orange (24A) along midveins, outer tepals brilliant orange-yellow (23B), strong orange (25A) along midveins, throat greyish yellow-green (191A); outside brilliant yellow (20A), tepal-margins strong orange (24A), strong orange (169C) at base; spots and papillae absent; nectaries strong orange (25A); pollen strong reddish orange (169A); stigma vivid reddish orange (30B). Fls 175 mm wide; tepals 110 × 32–45 mm, margins slightly ruffled, tips slightly recurved. Lvs scattered, 120 × 15 mm. Stems 0.9–1.1 m, brownish with green, and with 3–9 fls. (A *longiflorum*/Asiatic (LA) hybrid.)

**VII(a-b/c) TOPWHITE**

Trade designation of ‘Zanlorwhite’

**VII(-/b) TORERO**

Trade designation of ‘Towny’.

**VIII(c/b) ‘Tornessa’**

cv. Parentage: Seedling 89-102 × seedling 89-22-2

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside light greenish yellow (8B), lighter towards tepal-tips, midveins yellow at base, young fls flushed with red at base; outside light greenish yellow (8B), young fls flushed with red at base; spots and papillae absent; nectaries green; pollen vivid yellowish pink (28A); stigma light yellow-green (145B). Fls 180 mm wide; tepals 148–150 × 46–68 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 205–215 × 32–42 mm, dark green. Stems 1–1.2 m, green with darker spots and stripes, and with 1–9 fls. Late. (An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096401).)

### **VI(b-c/b,d) 'Trend'**

cv. Parentage: (*henryi* × 'Gold Eagle') × unknown) × (('Andrea' (of Geiser) × 'Sommertraum') × (*henryi* × 'Gold Eagle'))

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside ochre-apricot, throat greenish; outside with green midribs; spots absent; nectaries green; pollen ochre-apricot; stigma green. Fls 105 mm wide, bowl- and turban-shaped; tepals 95 × 44 mm, strongly recurved, inner tepal-margins gently wavy; pedicels 120 mm, thin, brown-dotted, with secondary buds. Lvs scattered, 145 × 17 mm, dull dark green. Stems 1.45 m, moderately thick, dotted with red-brown. Mid July.

### **I(a-b/b-c) 'Trial'**

cv. Parentage unknown

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2004), REG: B. Mičulka (2004)

Inside pale apricot, centre deeper, red-pink over nectaries, tepal-tips pinkish; outside pale pink with pale apricot tepal-edges, midribs greenish and brownish; spots few, on a quarter of each tepal; nectaries white-granulated; pollen ochre; stigma pale green. Fls 170 mm wide; tepals 70 × 37 mm, margins slightly wavy, tips slightly recurved, the outer more so; pedicels 60 mm, dotted with dark brown. Lvs scattered, 85 × 12 mm, dull dark green. Stems 0.65 m, thick, sparsely dotted. Mid July.

### **VII(a/b) TROFEO**

Trade designation of 'Zanlortrofeo'

### **VII(a/b-c) 'Tropical'**

cv. Parentage: Seedling 88-225-1 × seedling 88-07

R: World Breeding B.V., N: World Breeding B.V. (2001), REG: World Breeding B.V. (2004)

Inside deep red/strong purplish red (60A/B), lighter towards tepal-margins and tips; outside moderate purplish red (66D); spots and papillae present; nectaries green; filaments pale green; pollen brown-red; stigma purple, spotted. Fls 190-230 mm wide; tepals 140 × 39-64 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 132-206 × 47-61 mm, dark green. Stems 1-1.2 m, green with darker spots and stripes, and with 4-12 fls. Midseason.

(Nomenclatural standard: colour print supplied by registrant (WSY 0096404).)

### **I(a-b/b) 'Truvér'**

cv. Parentage: ('Casyra' × unknown) × (('Alexis' × 'Artistic') × 'Mocha')

R: J. Kovář (1999), N: J. Kovář (2005), I: J. Kovář (2005), REG: J. Kovář (2005)

Inside pale apricot with shining centre; outside pinkish at base; brushmarks dark crimson; nectaries greenish, edged with yellow-green; pollen brown-orange; stigma pink-crimson. Fls 115 mm wide; tepals 70 × 30 mm, slightly recurved; pedicels 70 mm, crimson-dotted. Lvs scattered, 70 × 10 mm, slightly glossy mid green. Stems 0.9 m, finely brown-dotted. Early July.

### **I(a-b/b-c) 'Trylek'**

cv. Parentage: 'Sterling Star' × unknown

R: J. Kovář (1998), N: J. Kovář (2005), I: J. Kovář (2005), REG: J. Kovář (2005)

Inside pale beige-apricot, each tepal with a small pinkish flame around nectary, tepal-tips paler, throat olive-green; outside pale beige-apricot, midribs pinkish at base; numerous crimson-brown spots and dashes on half of each tepal; nectaries silver-granulated, with yellowish border; pollen bright brown-orange; stigma crimson. Fls 140 mm wide; tepals 75 × 33 mm, very slightly recurved; pedicels 50 mm, thin, finely pale brown-dotted. Lvs scattered, 65 × 8 mm, dull yellow-green. Stems 0.8 m, slender, finely pale brown-dotted. Early July.

### VIII(c/b) 'Urandi'

cv. Parentage: Seedling E 905 × seedling 89.32.11-1

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside dark purplish pink (186C) with pale yellow-green (4D) along midveins, midveins brilliant yellow (7B) at base; outside dark purplish pink (186C) flushed with pale yellow-green (4D); spots absent; papillae present; nectaries green; pollen brown-red; stigma green. Fls 200 mm wide; tepals 157-168 × 44-67 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 157-265 × 25-47 mm, dark green. Stems 1-1.2 m, green, with 1-9 fls. Late.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096406).)

### I(a/b) 'Val di Fassa'

cv. Parentage unknown

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside strong orange-yellow (17A), throat light yellow-green (145D); outside vivid yellow (14A), tepal-margins strong orange-yellow (17A), midveins moderate olive-green (146A) at base and top, base also with some brownish orange (171B); spots absent; papillae strong orange-yellow (17A); nectaries strong yellow-green (144A); pollen moderate reddish brown (174A); stigma deep purplish red (59B). Fls 165 mm wide; tepals 115 × 45-60 mm, margins smooth, tips recurved. Lvs scattered, 180 × 15 mm. Stems 1.25-1.5 m, brownish with green spots, and with c.6 fls.

### I(b/b-c) 'Večernice'

cv. Parentage: ('Juno' × 'Honey Creme') × ('Parfait' × 'Wattle Bird')

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside old rose-pink with small yellow gleam around nectaries; tepal-tips and throat paler; spots black-brown, oval, numerous, on half of each tepal; papillae present; nectaries white- and pink-granulated; pollen brown, stigma dark purple-brown. Fls 140 mm wide; tepals 90 × 47 mm, slightly curved; pedicels 70 mm, brown-black. Lvs scattered, 85 × 14 mm, glossy dark green. Stems 0.6 m, thick, dark green. Late June.

### VIII(a/b-c) 'Verriers'

cv. Parentage: Seedling BW 96-33 × seedling AW 94-09

R: Gebr. Vletter & Den Haan, N: Gebr. Vletter & Den Haan (2002), REG: Vletter & Den Haan Beheer B.V. (2004)

Inside yellowish white (155D); outside yellowish white (155D), pale greenish yellow (2D) along midveins, at base and top strong yellow-green (144B), outer tepals also with a strong purplish red (63B) streak; spots and papillae absent; nectaries strong yellow-green (143C); pollen moderate brown (165A); stigma light greenish yellow (1C). Fls 200 mm wide; tepals 117 × 53-65 mm, margins smooth, tips recurved. Lvs scattered, 140 × 22 mm. Stems 0.9-1.4 m, green with darker markings, and with 3-4 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

### VIII(a-b/b) 'Visaversa'

cv. Parentage: Seedling 123-1 × seedling 90.056-14

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside moderate red/moderate purplish red (185B/C), midveins yellow-green at base changing into pink-red; outside moderate purplish red (185C); spots absent; papillae present; nectaries green; pollen brown-red; stigma grey-purple. Fls 200 mm wide; tepals 135-140 × 42-60 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 180-195 × 23-39 mm, dark green. Stems 0.9-1.3 m, green with darker spots and stripes, and with 2-9 fls. Midseason to late.

(An Orienpet (OT) lily. Nomenclatural standard: colour print supplied by registrant (WSY 0096402).)

#### **V(b/-) 'White Magic'**

cv. Parentage unknown

R: J.W.A. van der Wereld, N: Liliande B.V. (2003), REG: J.W.A. van der Wereld (2004)

Inside yellowish white (155D), midveins green at base; outside yellowish white (155D); spots and papillae absent; nectaries green; pollen orange-yellow; stigma light yellow. Tepals long, margins very slightly ruffled, tips recurved. Lvs scattered. Stems tall, green with few fls. Midseason to late.

#### **VIII(a-b/b) 'White Tycoon'**

cv. Parentage unknown

R: Testcentrum voor Siergewassen B.V., N: Marklily C.V. (2003), REG: Marklily C.V. (2004)

Inside with outer tepals yellowish white (155D), strong yellow-green (143A) along nectaries, inner tepals with dark purple (79A) picotee, throat strong yellow-green (144B), edges yellowish white (155D); outside with outer tepals yellowish white (155D), strong yellow-green (143C) at base, midribs moderate purplish red (59C) in basal third, with green-yellow flush when young, purple colour fading with age, strong yellow-green (144B) at top, inner tepals with dark purple (79A) picotee; spots and papillae absent; nectaries strong yellow-green (143A), hairy; pollen dark reddish orange (175C); stigma strong yellow-green (144A). Fls 130 mm wide; tepals 120 × 50-60 mm, margins smooth, tips not recurved. Lvs scattered, 115-130 × 20-30 mm, dark green. Stems 0.95-1.2 m, green, with c.3 fls.  
(A *longiflorum*/Asiatic (LA) hybrid.)

#### **VI(b/a-b) 'Wirena'**

cv. Parentage: Pink Perfection Group × 'Pink Pearl'

R: R. Widderski, N: R. Widderski, I: R. Widderski (2004), REG: R. Widderski (2004)

Inside creamy white with pink veining, throat golden; outside red-brown, midribs brown; spots absent; pollen yellow-brown; stigma brown. Fls 150 mm wide; tepals 130 × 60 mm, strongly recurved, tough; pedicels 90 mm, thick, green and brown. Lvs scattered, 130 × 11 mm, glossy mid green. Stems 1.2 m, thick, green and brown, with medium-sized infl. Mid July.

#### **VI(b/a) 'Wiroma'**

cv. Parentage: 'Anna Maria' × 'Fatima'

R: R. Widderski, N: R. Widderski, I: R. Widderski (2004), REG: R. Widderski (2004)

Inside deep orange, tepal-tips paler; outside brown-red-orange, midribs brown-red; spots absent; pollen yellow; stigma grey-bronze. Fls 100 mm wide; tepals 75 × 50 mm, slightly curved, margins gently ruffled; pedicels 80 mm long, thick, green, with secondary buds. Lvs scattered, 90 × 12 mm, dull mid green. Stems 1.1 m, thick, brown-dotted, with medium-sized infl. Early July.

#### **VII(a/b) 'Woodstock'**

cv. Parentage: White/yellow-flowered seedling × 'Ascari'

R: Bischoff Tulleken Lelies B.V., N: Bischoff Tulleken Lelies B.V., REG: Bischoff Tulleken Lelies B.V. (2004)

Inside with inner tepals moderate purplish pink (62B), light purplish pink (62C) towards margins, with some yellowish white (155D) next to nectaries, and brilliant greenish yellow (3A) above nectaries, outer tepals similar but apical half with narrow yellowish white (155D) margin; outside with outer tepals strong purplish pink (63D), pale purplish pink (65D) towards margins, midribs strong yellow-green (143C) at

base and top, in between brilliant yellow-green (150B), next to each midrib a small deep purplish red (59B) patch, inner tepals pale yellow-green (155A) in basal half, changing into light purplish pink (62C) towards margins and tips, midribs deep purplish red (59B) at base, towards top becoming brilliant yellow-green (150B); spots absent; papillae yellowish white (155D) and brilliant greenish yellow (3A); nectaries strong yellow-green (144C), brilliant greenish yellow (3A) at top and edge; pollen dark reddish orange (173A); stigma brilliant violet (92A). Fls 180-250 mm wide, slightly fragrant; tepals 123 × 50-71 mm, margins smooth to slightly ruffled, tips recurved. Lvs scattered, 155 × 40 mm. Stems 0.7-1 m, green with darker markings, and with c.3 fls.

#### **VII(a-b/b) 'Xotica'**

cv. Parentage: Seedling 88.24 × 'Pink Paradise'

R: World Breeding B.V., N: Liliande B.V. (2003), REG: World Breeding B.V. (2004)

Inside moderate purplish pink (62B), each tepal with brilliant/light yellow (10A/B) flame, midveins brilliant yellow (10A) at base; outside light purplish pink/very pale purple (73C/D); spots and papillae yellow; nectaries green; filaments pale green; pollen brown-red; stigma dark purple. Fls 180 mm wide; tepals 160-165 × 55-85 mm, margins slightly ruffled, tips strongly recurved. Lvs scattered, 106-219 × 34-45 mm, dark green. Stems 0.8-1.1 m, green, with 3-11 fls. Midseason to late.

(Nomenclatural standard: colour print supplied by registrant (WSY 0096403).)

#### **VIII(b/b) 'Yellow Diamond'**

cv. Parentage: 'Jonita' × seedling LA 93007 T

R: De Jong Lelies Beheer B.V., N: De Jong Lelies Beheer B.V. (2004), REG: De Jong Lelies Holland B.V. (2004)

Inside vivid yellow (12A), throat light yellow/pale greenish yellow (10C/D) with some strong yellow-green (144C); outside brilliant/light yellow (10A/B), tepal-margins vivid yellow (12A); spots dark red (187A); papillae few, vivid yellow (12A); nectaries strong/light yellow-green (144C/D); pollen strong brown/dark red (172A/187A); stigma moderate purplish red/dark purplish pink (186B/C). Fls 190 mm wide; tepals 110-115 × 39-50 mm, margins smooth, tips slightly recurved. Lvs scattered, 155 × 22 mm. Stems 0.95-1.1 m, light green with darker markings, and with 3-11 fls.

(A *longiflorum*/Asiatic (LA) hybrid.)

#### **VI(b/a) 'Yellow Surf'**

cv. Parentage: 'Anna Maria' × 'Afričanka'

R: A. Mego (1994), N: A. Mego (2005), I: A. Mego (2005), REG: A. Mego (2005)

Inside lemon-yellow; outside with green and brown midribs; spots absent; nectaries green; pollen golden yellow; stigma dark green. Fls 140 mm wide, strongly scented; tepals 160 × 50 mm, strongly recurved, margins smooth; pedicels 100 mm long, brown-dotted, with tertiary buds. Lvs scattered, 150 × 10 mm, dull mid green. Stems 1.5 m, thick, dark green. Late June.

#### **VIII(a/b) 'Zanlahalley'**

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2001), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside light orange-yellow (22C), tepal-margins and tips strong orange (24A), throat light yellow-green (145D); outside light orange-yellow (24C), tepal margins strong orange-yellow (24B), light yellow-green (144D) along midribs; spots dark red (59A); papillae absent; nectaries strong yellow-green (144B); pollen greyish red (178A); stigma brilliant orange (29A). Fls 172 mm wide; tepals 115 × 40-54 mm, margins smooth, tips recurved. Lvs scattered, 120 × 22 mm. Stems 0.8-1.3 m, green, with c.3 fls.

Trade designation: HALLEY.  
(*A. longiflorum*/Asiatic (LA) hybrid.)

### **VIII(a/b) 'Zanlapromes'**

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2003), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside pale yellow-green (4D), with strong yellow-green (144C) flush/stripe along basal half of each midvein, throat strong yellow-green (144C), with greenish white (155C) on edges; outside with outer tepals pale yellow-green/pale yellow (4D/8D), midribs with a strong yellow-green (144C) stripe, strong yellow-green (144B) at top, at base with little moderate red (181B) stripes, inner tepals pale yellow-green (4D), with moderate red (181B) stripe at base; spots and papillae absent; nectaries greenish white (155C); pollen moderate brown (165A); stigma moderate purplish red (185C) on a light yellow-green (144D) ground. Fls 165 mm wide; tepals 130-133 × 46-59 mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs scattered, 140 × 22 mm. Stems 0.9-1.5 m, green with darker markings, and with 4-6 fls.

Trade designation: PROMESA.

(*A. longiflorum*/Asiatic (LA) hybrid. The fls age to pale yellow-green (155A).)

### **VIII(a/a) 'Zanlaspacer'**

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2002), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside moderate pink (51D), tepal-margins with strong reddish orange (31B) flush (which becomes more prominent with age) and dark red (59A) picotee, along midveins deep pink (51B), throat strong purplish red (59D); outside with outer tepals deep purplish pink (70C), flushed with moderate yellowish pink (39C) towards margins, margins strong red (47B), midribs moderate yellow-green (139D) at base and top, inner tepals moderate purplish pink (70D), strong pink (50C) towards margins, deep purplish pink (70C) along midribs, midribs moderate yellow-green (139D) at base; spots many, dark red (59A); papillae absent; nectaries strong yellow-green (144B); pollen vivid reddish orange (32A); stigma strong reddish orange (31B). Fls 140 mm wide; tepals 130 × 41-51 mm, margins smooth, tips slightly recurved. Lvs scattered, 140 × 20 mm. Stems 1.15-1.2 m, green with darker markings, and with c.5 fls.

Trade designation: SPACER.

(*A. longiflorum*/Asiatic (LA) hybrid.)

### **I(a/b) 'Zanlazfica'**

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2003), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside strong orange (25B), vivid yellowish pink (28A) towards tepal-margins and tips; outside vivid orange (28B), tepal-tips moderate yellow-green (143D); spots absent; papillae vivid yellowish pink (28A); nectaries strong yellow-green (143C), pollen moderate red (179A); stigma vivid reddish orange (30B). Fls 160 mm wide; tepals 100 × 40-50 mm, margins smooth, tips recurved. Lvs scattered, 150 × 18 mm. Stems 0.9-1.25 m, brownish with green, and with 7-8 fls.

Trade designation: BENFICA.

### **VII(a/b) 'Zanloropium'**

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2003), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside deep purplish red (61A), strong purplish red/strong reddish purple (72A/B) towards tepal-margins, apical part of each tepal with narrow yellowish white (155D) margin, midveins deep red (60A), throat yellowish white (155D); outside with outer tepals deep purplish red (61A), deep purplish red (71A) towards margins, midribs yellowish white (155D) and deep red (60A), strong yellow-green (144C) at top, margins at base pale greenish yellow (1D), inner tepals strong purplish red (61B), strong purplish red (71B) towards margins, at top picotee yellowish white (155D), midribs yellowish white (155D) with some moderate purplish red (64A), light yellow-green (144D) at base, margins pale greenish yellow (1D) at base; spots absent; papillae dark red (59A); nectaries strong yellow-green (144B) with brilliant greenish yellow (6A) edge; pollen deep orange-yellow (163A); stigma light purple (77C). Fls 195 mm wide, fragrant; tepals 135 × 50-68 mm, margins ruffled, tips slightly recurved. Lvs scattered, 215 × 45 mm. Stems 1.2-1.6 m, green, with c.5 fls.  
Trade designation: OPIUM.

### VII(a/b) 'Zanlortrofeo'

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2003), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside with inner tepals vivid purplish red (67B) towards tip and from halfway upwards at margins fading into very pale purple (69D), along midveins strong red (53B), outer tepals strong purplish red (67A) fading towards margins into very pale purple (69B), margins at tip very pale purple (69D), along midveins strong red (53B), throat yellowish white (155D); outside with outer tepals strong purplish red (64C), fading into moderate purplish pink (66D) towards margins, margins narrow, yellowish white (155D), midribs strong yellow-green (144C) at base, fading into light yellow-green (145D) at top with a strong purplish red (64C) flush, strong yellow-green (143B) at top, inner tepals deep purplish pink (64D), with yellowish white (155D) veins at base, picotee moderate purplish red (64A), fading into yellowish white (155D) at top, midribs light yellow-green (142D) at base; spots absent; papillae many, moderate purplish red (58A); nectaries strong yellow-green (144B), flushed with strong red (53B) and with light greenish yellow (1C) edge; pollen absent; stigma yellowish grey (194C). Fls 220 mm wide; tepals 115 × 44-55 mm, margins ruffled, tips slightly recurved. Lvs scattered, 150 × 43 mm. Stems 1-1.4 m, light green with darker markings, and with c.7 fls.  
Trade designation: TROFEO.

### VII(a/b) 'Zanlorvenna'

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2003), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside strong purplish red (64B), tepal-margins and tips greenish white (155C), along midveins deep red/strong purplish red (60A/B), throat greenish white (155C); outside with outer tepals moderate purplish red (64A/70A), margins greenish white (155C), base strong yellow-green (144C) and moderate red (185B), midribs with greenish white (155C) stripe flushed with moderate/strong purplish red (64A/B), at top strong yellow-green (144B) with some strong yellow-green (144A), inner tepals strong purplish red/strong reddish purple (64B/70B), at base and margins greenish white (155C), midribs greenish white (157D) at base, changing into moderate purplish red (64A), at top light yellow-green (144D); spots absent; papillae greenish white (155C) and deep purplish red (187D); nectaries strong yellow-green (144B) with light yellow (10C) edge; pollen moderate reddish brown (175A); stigma light grey/deep purple (191C/79C). Fls 270 mm wide, fragrant; tepals 165 × 52-75 mm, margins ruffled, tips slightly recurved. Lvs scattered, 200 × 47 mm. Stems 1.1-1.5 m, green, with 3-7 fls.  
Trade designation: RAVENNA.

### **VII(a-b/c) 'Zanlorwhite'**

cv. Parentage unknown

R: Van Zanten Flowerbulbs B.V., N: Van Zanten Flowerbulbs B.V. (2003), REG: Van Zanten Flowerbulbs B.V. (2004)

Inside yellowish white (155D), midveins pale yellow-green (155A); outside with inner tepals yellowish white (155D), midribs strong yellow-green (144C) at base, at top brilliant greenish yellow (151D) with a little strong yellow-green (143B), outer tepals similar but with a vivid purplish red (57D) patch next to base of midribs; spots absent; papillae yellowish white (155D); nectaries moderate yellow-green (143D); pollen strong reddish orange (169A); stigma deep reddish purple (77A). Fls 245 mm wide, fragrant; tepals 133 × 54-72 mm, margins ruffled, tips recurved. Lvs scattered, 215 × 49 mm, dark green. Stems 1.1-1.45 m, green, with c.4 fls.

Trade designation: TOPWHITE.

### **VI(b/b) 'Zobor'**

cv. Parentage: 'White Henryi' × ('Damson' × 'Mabel Violet')

R: A. Mego (1995), N: A. Mego (2005), I: A. Mego (2005), REG: A. Mego (2005)

Inside white suffused with pale pink, tepal-centres and throat yellow; outside dark pinkish purple, midribs green; spots absent; nectaries dark green, pollen yellow; stigma grey-green. Fls 140 mm wide, slightly scented; tepals 115 × 45 mm, slightly recurved, margins slightly wavy; pedicels 100 mm, thick, brown-dotted. Lvs scattered, 150 × 18 mm, dull mid green. Stems 1.5 m, thick, dark green. Early July.

### **VI(b-c/b) 'Žonglér'**

cv. Parentage unknown

R: B. Mičulka, N: B. Mičulka, I: B. Mičulka (2005), REG: B. Mičulka (2005)

Inside cream, throat pale beige-apricot; outside with pale green midribs; spots absent; nectaries dark green; pollen brown; stigma dark red-brown. Fls 155 mm wide; tepals 105 × 40 mm, slightly curved, wavy and twisted; pedicels 55 mm, dark green, slightly brownish-dotted. Lvs scattered, 1-5 × 19 mm, glossy dark green. Stems 0.75 m, thick, brownish-dotted. Late June.

## LIST OF RAISERS, REGISTRANTS AND INTRODUCERS

(† = deceased)

ARTHUR, Beth	Thorpdale South, Victoria, Australia
BARBER, L. Maureen	Oakville, Ontario, Canada
BAZETT, Richard H.	Kelowna, British Columbia, Canada
BISCHOFF TULLEKEN LELIES B.V.	Wieringerwerf, The Netherlands
BRANDENBURG-ELSHOF B.V.	Anna Paulowna, The Netherlands
BROWN, Judith J.	Red Deer, Alberta, Canada
†CORNEIL, E.A.	Lindsay, Ontario, Canada
DALE, Daniel	Stony Plain, Alberta, Canada
DE VRIES VEREDELING B.V.	Breezand, The Netherlands
DIERSSEN, Hermann	Lemgo, Germany
DUDEK, Jiří	Hlučín, Czech Republic
ESTATE PERENNIALS	Spruce Grove, Alberta, Canada
EVANS Dr Ieuan R.	Spruce Grove, Alberta, Canada
FLORA HOLLAND	Rijnsburg, The Netherlands
GRIESBACH, Dr Robert J.	Belavan, Wisconsin, USA
HAY, Donna J.	Saskatoon, Saskatchewan, Canada
HILLCREST HARMONY FLOWERS	Churchbridge, Saskatchewan, Canada
HOOGLAND B.V.	Dirkshorn, The Netherlands
HORNER, Estelle	see Horner Lilies
HORNER LILIES	Thorndale, Ontario, Canada
JONG LELIES BEHEER B.V., De	Andijk, The Netherlands
JONG LELIES HOLLAND B.V., De	Andijk, The Netherlands
JOSLIN, Marvin	see Estate Perennials
KOVÁŘ, J.	Volyně, Czech Republic
LAAN LELIE	Zwaagdijk, The Netherlands
LILIANDE B.V.	't Zand, The Netherlands
LILY COMPANY B.V.	Andijk, The Netherlands
†LISS, Mary	Onaway, Alberta, Canada
MAK 't ZAND B.V.	't Zand, The Netherlands
MAK BREEDING B.V.	't Zand, The Netherlands
MANITOBA REGIONAL LILY SOCIETY	Canada
MAREL B.V., A. van der	Gravenzande, The Netherlands
MAREL LELIE B.V., van der	Gravenzande, The Netherlands
MARKLILY C.V.	Lisse, The Netherlands
MEGO, Anton	Bratislava, Slovakia
MÍČULKA, Břetislav	Velehrad, Czech Republic
NELSON, Dolores	Regina, Saskatchewan, Canada
PROCHASKA, Richard A.	New Prague, Minnesota, USA
PUTLAND, Diann	see Hillcrest Harmony Flowers
REZEK, J.	Brno, Czech Republic
SANDE B.V.	't Zand, The Netherlands
TESTCENTRUM VOOR	
SIERGEWASSEN B.V.	Lisse, The Netherlands
TRIOR LELIE B.V.	Hillegom, The Netherlands
UNIVERSITY OF LATVIA	
BOTANICAL GARDEN	Rīga, Latvia
VALLEY K GREENHOUSES	Edberg, Alberta, Canada
VERDEGAAL & ZN EXPORT	
B.V., Fa G.A.	Noordwijkerhout, The Netherlands
VINK & ZN, Fa Q.J.	St. Maartensvlotbrug, The Netherlands
VLETTER & DEN HAAN BEHEER B.V.	Rijnsburg, The Netherlands

VLETTER & DEN HAAN, Gebr.  
†WADEKAMPER, Julius  
WERELD B.V., J.W.A. van der  
WIDERSKI, Roman  
WIONZEK, J.  
WORLD BREEDING B.V.  
ZANTEN FLOWERBULBS B.V., Van  
ZORGEVICS, Ādolfš

Rijnsburg, The Netherlands  
Parkdale, Oregon, USA  
Breezand, The Netherlands  
Rybnik, Poland  
Buchanan, Saskatchewan, Canada  
Breezand, The Netherlands  
Hillegom, The Netherlands  
Riga, Latvia



